

wwPDB X-ray Structure Validation Summary Report (i)

Oct 11, 2023 – 09:25 AM EDT

PDB ID	:	7L2C
Title	:	Crystallographic structure of neutralizing antibody 2-51 in complex with
		SARS-CoV-2 spike N-terminal domain (NTD)
Authors	:	Cerutti, G.; Reddem, E.R.; Shapiro, L.
Deposited on	:	2020-12-16
Resolution	:	3.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 3.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	1557 (3.82 - 3.50)
Clashscore	141614	1037 (3.80-3.52)
Ramachandran outliers	138981	1004 (3.80-3.52)
Sidechain outliers	138945	1002 (3.80-3.52)
RSRZ outliers	127900	1441 (3.82-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	342	% 71% 11%	·	16%	_
1	В	342	% 68% 11%		20%	_
2	С	227	% 8 5%		13%	·
2	Н	227	% 86 %		11%	·
3	D	215	85%		13%	·



Mol	Chain	Length	Quality of chain		
3	L	215	90%	9%	•
4	Е	2	100%		-

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	В	405	-	-	-	Х



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 11381 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	1 Δ	286	Total	С	Ν	Ο	S	0	0	0
	200	2301	1487	380	426	8	0	0	0	
1	В	973	Total	С	Ν	Ο	S	0	1	0
I D	213	2210	1433	363	406	8	0		0	

Chain	Residue	Modelled	Actual	Comment	Reference
А	335	GLY	-	expression tag	UNP P0DTC2
А	336	SER	-	expression tag	UNP P0DTC2
А	337	LEU	-	expression tag	UNP P0DTC2
А	338	GLU	-	expression tag	UNP P0DTC2
А	339	VAL	-	expression tag	UNP P0DTC2
А	340	LEU	-	expression tag	UNP P0DTC2
А	341	PHE	-	expression tag	UNP P0DTC2
А	342	GLN	-	expression tag	UNP P0DTC2
В	335	GLY	-	expression tag	UNP P0DTC2
В	336	SER	-	expression tag	UNP P0DTC2
В	337	LEU	-	expression tag	UNP P0DTC2
В	338	GLU	-	expression tag	UNP P0DTC2
В	339	VAL	-	expression tag	UNP P0DTC2
В	340	LEU	-	expression tag	UNP P0DTC2
В	341	PHE	-	expression tag	UNP P0DTC2
В	342	GLN	-	expression tag	UNP P0DTC2

There are 16 discrepancies between the modelled and reference sequences:

• Molecule 2 is a protein called 2-51 heavy chain.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace
2	9 Н	221	Total	С	Ν	0	S	0	2	0
	221	1687	1071	272	336	8	0	2		
2	C	222	Total	С	Ν	0	S	0	0	0
	U		1696	1078	274	336	8	0	Z	





• Molecule 3 is a protein called 2-51 light chain.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
9	5 I	012	Total	С	Ν	0	\mathbf{S}	0	0	0
о L	213	1573	979	263	325	6	0	0	0	
2	П	012	Total	С	Ν	0	S	0	0	0
	213	1573	979	263	325	6	0	0	U	

• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	Е	2	Total 28	C 16	N 2	O 10	0	0	0

• Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
5	А	1	Total 14	C 8	N 1	0 5	0	0
5	А	1	Total 14	0 C 8	N 1	0 5	0	0



Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
5	Δ	1	Total	С	Ν	0	0	0
5	A	L	14	8	1	5	0	0
5	Δ	1	Total	С	Ν	0	0	0
0	11	I	14	8	1	5	0	0
5	Δ	1	Total	С	Ν	Ο	0	0
0	11	T	14	8	1	5	0	0
5	Δ	1	Total	С	Ν	Ο	0	0
0	11	T	14	8	1	5	0	0
5	А	1	Total	С	Ν	Ο	0	0
0		1	14	8	1	5	0	
5	В	1	Total	С	Ν	Ο	0	0
	D	1	14	8	1	5	0	0
5	В	1	Total	С	Ν	0	0	0
	D	1	14	8	1	5	Ŭ	0
5	В	1	Total	С	Ν	Ο	0	0
		-	14	8	1	5		<u> </u>
5	В	1	Total	С	Ν	Ο	0	0
	2	-	14	8	1	5		
5	В	1	Total	С	Ν	Ο	0	0
		*	14	8	1	5	Ŭ	
5	В	1	Total	С	Ν	Ο	0	0
			14	8	1	5		

Continued from previous page...

• Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	6	Total Ca 6 6	0	0
6	В	6	Total Ca 6 6	0	0
6	Н	7	Total Ca 7 7	0	0
6	L	3	Total Ca 3 3	0	0
6	С	11	Total Ca 11 11	0	0
6	D	6	Total Ca 6 6	0	0

• Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
7	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
7	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
7	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
7	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
7	Н	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
7	Н	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
7	L	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
7	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 8 is CACODYLATE ION (three-letter code: CAC) (formula: $C_2H_6AsO_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	А	1	Total 5	As 1	$\begin{array}{c} \mathrm{C} \\ \mathrm{2} \end{array}$	O 2	0	0

• Molecule 9 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	С	1	Total 10	С 6	0 4	0	0

• Molecule 10 is water.



71	001
(]	L20

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	8	Total O 8 8	0	0
10	В	9	Total O 9 9	0	0
10	Н	5	Total O 5 5	0	0
10	L	4	Total O 4 4	0	0
10	С	7	Total O 7 7	0	0
10	D	8	Total O 8 8	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Spike glycoprotein



• Molecule 3: 2-51 light chain

Chain L:	90%	9%	
41 33 834 834 835 855 856 855 850 855 850 856 850 850 850 850 850 850 850 850 850 850	T145 Y172 Y173 Y174 Y175 Y175 Y196 Y196		
• Molecule 3: 2-51 light chain			
Chain D:	85%	13%	•
01 96 99 89 89 89 89 850 850 850 850 853 883 883 883 883 883	T102 K103 L106 L105 L132 L132 L135 L135 L135 L135 L136 L144 L144 L144 L144 L144 L144 L144 L14	N169 L178 K186	R189 T196 E210

CYS SER

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:

100%

NAG1 NAG2



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	66.79Å 115.78Å 137.61Å	Depositor
a, b, c, α , β , γ	90.00° 99.97° 90.00°	Depositor
Bosolution (Å)	88.03 - 3.65	Depositor
Resolution (A)	88.03 - 3.65	EDS
% Data completeness	98.3 (88.03-3.65)	Depositor
(in resolution range)	$98.3 \ (88.03 - 3.65)$	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.52 (at 3.67 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
P. P.	0.216 , 0.272	Depositor
n, n_{free}	0.216 , 0.271	DCC
R_{free} test set	1128 reflections (4.97%)	wwPDB-VP
Wilson B-factor $(Å^2)$	68.4	Xtriage
Anisotropy	0.528	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.34 , 79.0	EDS
L-test for $twinning^2$	$ \langle L \rangle = 0.43, \langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	11381	wwPDB-VP
Average B, all atoms $(Å^2)$	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, ACT, CA, PGE, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		Bond lengths		ond angles
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.37	0/2366	0.77	6/3222~(0.2%)
1	В	0.52	4/2274~(0.2%)	0.78	6/3096~(0.2%)
2	С	0.35	0/1740	0.66	0/2372
2	Н	0.45	2/1731~(0.1%)	0.74	2/2362~(0.1%)
3	D	0.40	1/1612~(0.1%)	0.65	1/2197~(0.0%)
3	L	0.39	0/1612	0.63	0/2197
All	All	0.42	7/11335~(0.1%)	0.71	15/15446~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	Н	0	3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
1	В	159	VAL	CB-CG2	-9.35	1.33	1.52
1	В	143	VAL	CB-CG2	-8.51	1.34	1.52
2	Н	196	CYS	CB-SG	-6.37	1.71	1.82
1	В	143	VAL	CB-CG1	-6.35	1.39	1.52
1	В	159	VAL	CB-CG1	-6.32	1.39	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	105	ILE	CG1-CB-CG2	-8.40	92.92	111.40
1	В	223	LEU	CB-CG-CD2	-8.07	97.28	111.00



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	В	176	LEU	CB-CG-CD1	-8.02	97.36	111.00
1	В	97	LYS	CB-CG-CD	7.74	131.72	111.60
1	А	126	VAL	CG1-CB-CG2	7.11	122.27	110.90

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	Н	97	ALA	Mainchain
2	Н	98[A]	TYR	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2301	0	2221	27	0
1	В	2210	0	2126	24	0
2	С	1696	0	1652	17	0
2	Н	1687	0	1635	10	0
3	D	1573	0	1516	18	0
3	L	1573	0	1516	11	0
4	Ε	28	0	25	0	0
5	А	98	0	91	0	0
5	В	84	0	78	0	0
6	А	6	0	0	0	0
6	В	6	0	0	0	0
6	С	11	0	0	0	0
6	D	6	0	0	0	0
6	Н	7	0	0	0	0
6	L	3	0	0	0	0
7	А	12	0	9	0	0
7	В	8	0	6	0	0
7	D	4	0	3	0	0
7	Н	8	0	6	0	0
7	L	4	0	3	0	0
8	А	5	0	0	0	0
9	С	10	0	14	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	А	8	0	0	0	0
10	В	9	0	0	0	0
10	С	7	0	0	0	0
10	D	8	0	0	0	0
10	Н	5	0	0	0	0
10	L	4	0	0	0	0
All	All	11381	0	10901	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 103 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:B:121:ASN:OD1	1:B:126:VAL:HG22	1.90	0.71	
1:A:33:THR:HB	1:A:219:GLY:HA3	1.73	0.70	
1:A:126:VAL:HG22	1:A:174:PRO:HA	1.75	0.69	
1:B:245[B]:HIS:H	1:B:260:ALA:HB2	1.57	0.68	
3:L:33:VAL:HG22	3:L:90:SER:HB3	1.77	0.65	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	282/342~(82%)	256 (91%)	26 (9%)	0	100	100
1	В	268/342~(78%)	254 (95%)	14 (5%)	0	100	100
2	С	222/227~(98%)	210 (95%)	12 (5%)	0	100	100
2	Н	221/227~(97%)	210 (95%)	11 (5%)	0	100	100
3	D	211/215~(98%)	201 (95%)	10 (5%)	0	100	100



001111	John precious page										
Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles				
3	L	211/215~(98%)	201 (95%)	10 (5%)	0	100	100				
All	All	1415/1568~(90%)	1332 (94%)	83 (6%)	0	100	100				

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	258/310~(83%)	258 (100%)	0	100	100
1	В	247/310~(80%)	247 (100%)	0	100	100
2	С	190/193~(98%)	190 (100%)	0	100	100
2	Н	189/193~(98%)	189 (100%)	0	100	100
3	D	176/178~(99%)	176 (100%)	0	100	100
3	L	176/178~(99%)	176 (100%)	0	100	100
All	All	1236/1362~(91%)	1236 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	115	GLN
2	Н	1	GLN
2	С	164	HIS
2	С	1	GLN
1	А	239	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dog	Tink	Bond lengths			Bond angles		
WIOI	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
4	NAG	Е	1	1,4	14,14,15	0.69	1 (7%)	17,19,21	1.04	1 (5%)
4	NAG	Е	2	4	14,14,15	1.62	2 (14%)	17,19,21	0.93	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	Е	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	Е	2	4	-	1/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
4	Ε	2	NAG	C1-C2	5.55	1.60	1.52
4	Е	2	NAG	O5-C1	2.17	1.47	1.43
4	Е	1	NAG	O5-C1	2.10	1.47	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	Е	2	NAG	C4-C3-C2	2.33	114.44	111.02
4	Е	1	NAG	C4-C3-C2	-2.11	107.93	111.02



There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Е	1	NAG	O5-C5-C6-O6
4	Е	1	NAG	C4-C5-C6-O6
4	Е	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry (i)

Of 63 ligands modelled in this entry, 39 are monoatomic - leaving 24 for Mogul analysis.



7L2C

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dog	Link	Bo	ond leng	ths	B	Bond angles		
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
7	ACT	Н	308	-	3,3,3	1.30	0	3,3,3	1.52	0	
5	NAG	В	404	1	14,14,15	0.45	0	17,19,21	1.55	2 (11%)	
7	ACT	D	307	-	3,3,3	1.36	0	3,3,3	1.29	0	
7	ACT	А	414	-	3,3,3	1.30	0	3,3,3	1.52	0	
5	NAG	А	401	1	$14,\!14,\!15$	0.43	0	17,19,21	0.56	0	
5	NAG	А	402	1	14,14,15	0.36	0	17,19,21	0.60	1 (5%)	
5	NAG	А	404	1	14,14,15	0.19	0	17,19,21	0.57	0	
7	ACT	В	414	-	3,3,3	1.28	0	3,3,3	1.37	0	
7	ACT	В	413	-	3,3,3	1.51	1 (33%)	3,3,3	1.29	0	
5	NAG	В	401	1	14,14,15	0.28	0	17,19,21	0.38	0	
5	NAG	В	405	1	14,14,15	0.56	0	17,19,21	1.30	2 (11%)	
7	ACT	А	415	-	3,3,3	1.31	0	3,3,3	1.53	0	
5	NAG	В	403	1	14,14,15	0.25	0	17,19,21	0.56	0	
5	NAG	В	402	1,6	14,14,15	0.53	0	17,19,21	0.46	0	
7	ACT	L	304	-	3,3,3	1.27	0	3,3,3	1.54	0	
5	NAG	А	406	1	14,14,15	0.43	0	17,19,21	1.24	2 (11%)	
5	NAG	А	403	1	14,14,15	0.91	1 (7%)	17,19,21	2.36	2 (11%)	
5	NAG	В	406	1	14,14,15	0.55	0	17,19,21	1.27	1 (5%)	
9	PGE	С	312	-	9,9,9	0.29	0	8,8,8	0.46	0	
5	NAG	А	407	1	14,14,15	1.16	1 (7%)	17,19,21	1.64	3 (17%)	
7	ACT	Н	309	-	3,3,3	1.30	0	3,3,3	1.38	0	
7	ACT	А	416	-	3,3,3	1.26	0	3,3,3	1.48	0	
8	CAC	A	417	-	0,4,4	-	-	0,6,6	-	-	
5	NAG	A	405	1	14,14,15	0.99	1 (7%)	17,19,21	1.34	3 (17%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	А	403	1	-	2/6/23/26	0/1/1/1
5	NAG	В	401	1	-	0/6/23/26	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10101	турс	Cham	ICCD	LIIIK	Cimais	TOTOTOTO	Tungs
5	NAG	А	407	1	-	4/6/23/26	0/1/1/1
5	NAG	В	405	1	-	0/6/23/26	0/1/1/1
5	NAG	В	406	1	-	2/6/23/26	0/1/1/1
5	NAG	В	404	1	-	5/6/23/26	0/1/1/1
5	NAG	В	403	1	-	1/6/23/26	0/1/1/1
5	NAG	А	405	1	-	3/6/23/26	0/1/1/1
5	NAG	В	402	1,6	-	1/6/23/26	0/1/1/1
5	NAG	А	402	1	-	0/6/23/26	0/1/1/1
5	NAG	А	401	1	-	3/6/23/26	0/1/1/1
5	NAG	А	404	1	-	0/6/23/26	0/1/1/1
5	NAG	A	406	1	-	2/6/23/26	0/1/1/1
9	PGE	С	312	-	-	1/7/7/7	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	А	407	NAG	O5-C1	-4.10	1.37	1.43
5	А	405	NAG	C1-C2	3.33	1.57	1.52
5	А	403	NAG	O5-C1	3.15	1.48	1.43
7	В	413	ACT	CH3-C	2.13	1.58	1.49

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	А	403	NAG	C2-N2-C7	6.62	132.33	122.90
5	А	403	NAG	C1-C2-N2	6.20	121.08	110.49
5	В	404	NAG	C1-O5-C5	4.09	117.73	112.19
5	А	407	NAG	C2-N2-C7	4.06	128.69	122.90
5	В	406	NAG	C2-N2-C7	4.01	128.62	122.90

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	А	403	NAG	C3-C2-N2-C7
5	А	406	NAG	C8-C7-N2-C2
5	А	406	NAG	O7-C7-N2-C2
5	А	407	NAG	C8-C7-N2-C2
5	В	404	NAG	C1-C2-N2-C7



There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	С	312	PGE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	286/342~(83%)	0.18	4 (1%) 75 63	47, 72, 115, 158	0
1	В	273/342~(79%)	0.17	2 (0%) 87 80	42, 73, 125, 158	0
2	С	222/227~(97%)	0.10	2 (0%) 84 74	40, 57, 85, 135	0
2	Н	221/227~(97%)	0.30	3 (1%) 75 63	42, 64, 93, 133	0
3	D	213/215~(99%)	0.01	0 100 100	40, 59, 79, 107	0
3	L	213/215~(99%)	0.11	0 100 100	45, 64, 94, 113	0
All	All	1428/1568~(91%)	0.15	11 (0%) 86 77	40, 65, 103, 158	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Н	130	SER	5.3
1	А	72	GLY	3.6
1	В	69	HIS	3.4
2	Н	129	LYS	3.1
2	С	131	THR	3.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
4	NAG	Е	2	14/15	0.80	0.32	66,101,115,120	0
4	NAG	Е	1	14/15	0.89	0.28	66, 84, 96, 98	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
6	CA	В	407	1/1	0.07	0.20	92,92,92,92	0
6	CA	А	412	1/1	0.45	0.18	$95,\!95,\!95,\!95$	0
5	NAG	В	405	14/15	0.56	0.54	84,103,114,119	0
6	CA	D	306	1/1	0.59	0.24	79,79,79,79	0
6	CA	В	409	1/1	0.66	0.18	88,88,88,88	0
6	CA	С	311	1/1	0.67	0.22	82,82,82,82	0
5	NAG	А	406	14/15	0.69	0.33	85,111,120,121	0

Continued on next page...



7	L2C	2

		Tueu 110	Chain	is page.	Atoma	DECC	DCD	D footoms (λ^2)	0 < 0.0
b CA D 303 1/1 0.69 0.23 84,84,84,84 0 5 NAG B 404 14/15 0.69 0.30 91,119,131,133 0 6 CA H 304 1/1 0.70 0.17 89,89,898 0 6 CA D 305 1/1 0.71 0.18 98,98,989 0 6 CA D 305 1/1 0.71 0.18 98,98,989 0 6 CA D 305 1/1 0.77 0.32 101,126,136,136 0 5 NAG B 402 14/15 0.77 0.28 77,910,0172 0 5 NAG B 401 14/15 0.78 0.30 95,109,172 0 5 NAG B 403 14/15 0.78 0.30 95,109,172 0 5 NAG B 403 14/15 0.84		Type	D	Res		RSCC	R5R	\mathbf{B} -factors(\mathbf{A}^{-})	Q<0.9
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	0 	CA		303		0.69	0.23	84,84,84,84	0
b CA II 304 $1/1$ 0.70 0.17 $89, 89, 89, 89$ 0 6 CA D 305 $1/1$ 0.71 0.18 $98, 98, 98, 98$ 0 7 ACT H 308 $4/4$ 0.72 0.39 $66, 66, 69, 76$ 0 6 CA C 303 $1/1$ 0.77 0.32 $101, 126, 136, 136$ 0 5 NAG A 401 $14/15$ 0.77 0.28 $65, 88, 104, 110$ 0 8 CAC A 417 $5/5$ 0.77 0.28 $67, 93, 109, 1172$ 0 5 NAG B 401 $14/15$ 0.78 0.38 $86, 107, 119, 130, 125$ 0 5 NAG B 401 $14/15$ 0.79 0.24 $107, 119, 130, 135$ 0 6 CA A 402 $14/15$ 0.81 0.23 $96, 107, 118, 119$	C C	NAG	В	404	$\frac{14}{10}$	0.69	0.30	91,119,131,133	0
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	0 	CA		304		0.70	0.17	89,89,89,89	0
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0	CA		305	1/1	0.71	0.18	98,98,98,98	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	ACT	H	308	4/4	0.72	0.39	66,66,69,76	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	C	303		0.77	0.17	85,85,85,85	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	5	NAG	A	401	14/15	0.77	0.32	101,126,136,136	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	5	NAG	B	402	14/15	0.77	0.28	65,88,104,110	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	8	CAC	A	417	5/5	0.77	0.28	77,95,109,172	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	5	NAG	A	405	14/15	0.78	0.38	86,107,120,125	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	5	NAG	В	401	14/15	0.78	0.30	95,109,117,120	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	NAG	В	403	14/15	0.79	0.24	107,119,130,135	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	6	CA	В	408	1/1	0.80	0.18	71,71,71,71	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	NAG	A	402	14/15	0.81	0.23	96,107,118,119	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	A	410	1/1	0.81	0.17	$66,\!66,\!66,\!66$	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	6	CA	С	307	1/1	0.82	0.34	90,90,90,90	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	7	ACT	А	416	4/4	0.83	0.25	70,75,80,86	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	NAG	В	406	14/15	0.83	0.30	$91,\!108,\!115,\!117$	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	С	301	1/1	0.83	0.15	73,73,73,73	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	7	ACT	L	304	4/4	0.84	0.27	$61,\!61,\!62,\!67$	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	С	305	1/1	0.86	0.10	72,72,72,72	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	L	301	1/1	0.86	0.14	81,81,81,81	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	NAG	А	403	14/15	0.86	0.28	91,102,106,107	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	С	302	1/1	0.86	0.14	77,77,77,77	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	Н	306	1/1	0.86	0.08	103,103,103,103	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	7	ACT	В	413	4/4	0.87	0.26	58,60,63,69	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	NAG	А	407	14/15	0.87	0.20	77,82,96,101	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	6	CA	D	302	1/1	0.87	0.14	80,80,80,80	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	NAG	А	404	14/15	0.87	0.22	71,79,86,94	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	Н	305	1/1	0.88	0.32	87,87,87,87	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	С	309	1/1	0.88	0.14	82,82,82,82	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	CA	С	306	1/1	0.88	0.14	73,73,73,73	0
6 CA H 307 1/1 0.89 0.20 72,72,72,72 0 6 CA A 409 1/1 0.89 0.11 60,60,60,60 0 0 DCF C 210 10/10 0.20 224 224,62,421 0	6	CA	L	303	1/1	0.89	0.21	83,83,83,83	0
6 CA A 409 1/1 0.89 0.11 60,60,60,60 0 0 DCD C 210 10/10 0.20 0.20 20.4054.61 0	6	CA	Н	307	1/1	0.89	0.20	72,72,72,72	0
	6	CA	А	409	1/1	0.89	0.11	60,60,60,60	0
9 PGE C 312 10/10 0.89 0.36 39,48,54,61 0	9	PGE	С	312	10/10	0.89	0.36	39,48,54,61	0
7 ACT D 307 4/4 0.90 0.27 47,54,54,57 0	7	ACT	D	307	4/4	0.90	0.27	47,54,54,57	0
6 CA B 412 1/1 0.90 0.15 100,100,100 0	6	CA	В	412	1/1	0.90	0.15	100,100,100,100	0
7 ACT A 415 4/4 0.90 0.19 42,52,55,58 0	7	ACT	А	415	4/4	0.90	0.19	42,52,55,58	0
7 ACT H 309 4/4 0.91 0.42 50,61,64,84 0	7	ACT	Н	309	4/4	0.91	0.42	50,61,64,84	0
6 CA C 308 1/1 0.91 0.16 84.84.84.84 0	6	CA	С	308	1/1	0.91	0.16	84,84,84.84	0
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	7	ACT	A	414	4/4	0.91	0.22	58,68,68,69	0

Continued from previous page...

Continued on next page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
7	ACT	В	414	4/4	0.91	0.21	71,77,80,82	0
6	CA	С	304	1/1	0.91	0.21	78,78,78,78	0
6	CA	Н	302	1/1	0.92	0.11	71,71,71,71	0
6	CA	А	411	1/1	0.92	0.12	74, 74, 74, 74	0
6	CA	L	302	1/1	0.94	0.10	$75,\!75,\!75,\!75$	0
6	CA	Н	301	1/1	0.94	0.16	64,64,64,64	0
6	CA	D	301	1/1	0.94	0.17	$65,\!65,\!65,\!65$	0
6	CA	В	411	1/1	0.94	0.20	83,83,83,83	0
6	CA	Н	303	1/1	0.94	0.23	74,74,74,74	0
6	CA	А	408	1/1	0.94	0.13	$61,\!61,\!61,\!61$	0
6	CA	А	413	1/1	0.95	0.18	80,80,80,80	0
6	CA	С	310	1/1	0.96	0.30	68,68,68,68	0
6	CA	D	304	1/1	0.97	0.24	66,66,66,66	0
6	CA	В	410	1/1	0.97	0.07	72,72,72,72	0

Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




















































































































































































































































6.5 Other polymers (i)

There are no such residues in this entry.

