



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2024 – 10:33 PM EST

PDB ID : 4L4Z  
Title : Crystal structures of the LsrR proteins complexed with phospho-AI-2 and its two different analogs reveal distinct mechanisms for ligand recognition  
Authors : Ryu, K.S.; Ha, J.H.; Eo, Y.  
Deposited on : 2013-06-10  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	 7% 73% 17% •• 6%
1	B	266	 8% 79% 13% •• 6%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Transcriptional regulator LsrR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	249	Total 1820	C 1138	N 324	O 348	S 10	0	0	0
1	B	250	Total 1856	C 1164	N 330	O 352	S 10	0	4	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	SER	-	expression tag	UNP P76141
B	52	SER	-	expression tag	UNP P76141

- Molecule 2 is (2S)-2,3,3-trihydroxy-4-oxopentyl dihydrogen phosphate (three-letter code: D5X) (formula: C<sub>5</sub>H<sub>11</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	Total 14	C 5	O 8	P 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	O	P	0	0
			14	5	8	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	74	Total	O	0	0
			74	74		
3	B	99	Total	O	0	0
			99	99		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional regulator LsrR



- Molecule 1: Transcriptional regulator LsrR



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.69Å 117.69Å 79.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.68 – 2.30 34.66 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.5 (34.68-2.30) 96.5 (34.66-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.172 , 0.233 0.180 , 0.238	Depositor DCC
$R_{free}$ test set	1357 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.3	Xtrriage
Anisotropy	0.008	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.057 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3877	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: D5X

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	0/1838	0.92	1/2483 (0.0%)
1	B	0.78	0/1887	0.90	0/2547
All	All	0.77	0/3725	0.91	1/5030 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	157	MET	N-CA-C	-9.00	86.71	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	156	TYR	Peptide
1	A	157	MET	Peptide
1	B	152	GLY	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1820	0	1874	40	0
1	B	1856	0	1921	32	0
2	A	14	0	11	0	0
2	B	14	0	11	0	0
3	A	74	0	0	2	0
3	B	99	0	0	4	0
All	All	3877	0	3817	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:ASN:HD22	1:A:258:ASN:H	1.09	0.99
1:B:164:ASN:HB2	1:B:165:ALA:HA	1.53	0.89
1:B:164:ASN:HB2	1:B:167:CYS:SG	2.15	0.86
1:B:157:MET:HA	1:B:160:ILE:HG13	1.63	0.81
1:A:258:ASN:H	1:A:258:ASN:ND2	1.77	0.79
1:B:164:ASN:CB	1:B:165:ALA:HA	2.12	0.78
1:B:153:VAL:O	1:B:156:TYR:HB2	1.86	0.74
1:B:129:MET:HB3	1:B:158:THR:HG21	1.72	0.71
1:A:153:VAL:O	1:A:156:TYR:HB2	1.93	0.68
1:A:143:GLN:NE2	1:A:165:ALA:O	2.27	0.68
1:B:133:GLN:HE21	1:B:158:THR:HG23	1.59	0.67
1:A:143:GLN:HA	1:A:167:CYS:SG	2.35	0.66
1:A:226:TYR:OH	3:A:635:HOH:O	2.10	0.66
1:B:108:HIS:HB2	3:B:574:HOH:O	1.96	0.65
1:B:165:ALA:HA	1:B:167:CYS:SG	2.42	0.60
1:A:258:ASN:ND2	1:A:258:ASN:N	2.47	0.60
1:A:221:ILE:HG23	1:A:227:ILE:CD1	2.32	0.59
1:B:157:MET:CA	1:B:160:ILE:HG13	2.31	0.59
1:A:162:GLN:CD	1:A:163:LEU:H	2.06	0.58
1:A:154:GLY:O	1:A:156:TYR:O	2.22	0.57
1:B:108:HIS:CD2	3:B:539:HOH:O	2.57	0.56

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:SER:H	1:B:157:MET:HG2	1.70	0.56
1:B:111:MET:HG2	1:B:138:PHE:CD2	2.41	0.56
1:B:164:ASN:CB	1:B:165:ALA:CA	2.84	0.56
1:A:155:SER:C	1:A:156:TYR:O	2.43	0.55
1:A:162:GLN:OE1	1:A:163:LEU:O	2.25	0.55
1:A:258:ASN:HD22	1:A:258:ASN:N	1.83	0.55
1:B:164:ASN:HB3	1:B:166:ALA:N	2.23	0.54
1:A:154:GLY:O	1:A:158:THR:HG21	2.08	0.54
1:A:170:ASN:OD1	1:A:196:VAL:HG22	2.08	0.54
1:B:160:ILE:C	1:B:160:ILE:HD12	2.29	0.53
1:A:76:THR:O	1:A:80:ARG:HG3	2.09	0.52
1:A:212:ALA:H	1:A:215:GLN:HE21	1.58	0.52
1:B:156:TYR:O	1:B:157:MET:HG2	2.09	0.52
1:A:164:ASN:O	1:A:165:ALA:HB3	2.09	0.51
1:B:103:GLY:HA3	1:B:130[A]:ASN:HD22	1.76	0.51
1:A:163:LEU:HD11	1:A:169:VAL:HG23	1.94	0.50
1:B:92:GLY:H	1:B:308:GLN:HE21	1.60	0.50
1:A:154:GLY:HA3	3:B:536:HOH:O	2.11	0.49
1:A:155:SER:HA	3:A:657:HOH:O	2.11	0.49
1:B:86:HIS:CD2	1:B:113:LEU:HD21	2.47	0.49
1:B:161:GLY:O	1:B:162:GLN:HG2	2.13	0.48
1:A:113:LEU:HD13	1:A:278:VAL:HG11	1.94	0.48
1:A:162:GLN:O	1:A:163:LEU:HB3	2.14	0.48
1:A:163:LEU:HG	1:A:164:ASN:N	2.29	0.48
1:B:133:GLN:HE21	1:B:158:THR:CG2	2.26	0.47
1:A:248:PHE:H	1:A:262:HIS:CE1	2.32	0.47
1:A:175:PRO:HB3	1:B:192:CYS:SG	2.56	0.46
1:B:76:THR:O	1:B:80:ARG:HG3	2.16	0.46
1:A:163:LEU:O	1:A:164:ASN:HB2	2.14	0.46
1:B:164:ASN:CB	1:B:167:CYS:SG	2.97	0.46
1:A:244:ILE:HG22	1:A:245:LEU:HG	1.97	0.46
1:B:133:GLN:NE2	1:B:158:THR:HG23	2.30	0.46
1:B:160:ILE:HD12	1:B:161:GLY:N	2.31	0.46
1:A:72:LEU:C	1:A:72:LEU:HD13	2.36	0.45
1:B:145:ARG:HG2	1:B:168:SER:HB3	1.98	0.45
1:A:103:GLY:HA3	1:A:130:ASN:ND2	2.31	0.45
1:A:222:ILE:HD12	1:A:227:ILE:CD1	2.47	0.45
1:A:117:GLN:HA	1:A:143:GLN:O	2.17	0.45
1:A:186:THR:HG21	1:B:183:ILE:HG12	2.00	0.44
1:B:76:THR:HA	1:B:79:ARG:NH1	2.33	0.43
1:A:222:ILE:HD12	1:A:227:ILE:HD11	2.01	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:GLN:O	1:A:216:GLN:C	2.56	0.43
1:A:266:ILE:CD1	1:A:266:ILE:N	2.83	0.42
1:B:120:LEU:HB3	1:B:146:LEU:HD12	2.01	0.42
1:A:161:GLY:O	1:A:162:GLN:CB	2.67	0.42
1:A:163:LEU:HD12	1:A:167:CYS:HB3	2.03	0.41
1:A:221:ILE:HG23	1:A:227:ILE:HD11	2.03	0.41
1:A:90:ILE:HD11	1:A:105:GLY:HA3	2.03	0.40
1:B:128:THR:HG23	3:B:508:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/266 (93%)	230 (93%)	10 (4%)	7 (3%)	5	3
1	B	252/266 (95%)	234 (93%)	13 (5%)	5 (2%)	7	6
All	All	499/532 (94%)	464 (93%)	23 (5%)	12 (2%)	6	4

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	156	TYR
1	A	160	ILE
1	A	163	LEU
1	A	164	ASN
1	B	155	SER
1	B	157	MET
1	A	162	GLN
1	B	158	THR
1	A	158	THR
1	B	168	SER

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	97	ASP
1	A	91	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/203 (93%)	167 (89%)	21 (11%)	6	6
1	B	193/203 (95%)	182 (94%)	11 (6%)	20	28
All	All	381/406 (94%)	349 (92%)	32 (8%)	11	13

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	LEU
1	A	88	ARG
1	A	101	ARG
1	A	112	SER
1	A	126	GLU
1	A	132	LEU
1	A	136	SER
1	A	140	SER
1	A	141	SER
1	A	142	GLN
1	A	157	MET
1	A	163	LEU
1	A	168	SER
1	A	170	ASN
1	A	194	LYS
1	A	222	ILE
1	A	227	ILE
1	A	243	ASP
1	A	258	ASN
1	A	266	ILE
1	A	270	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	130[A]	ASN
1	B	130[B]	ASN
1	B	136	SER
1	B	140	SER
1	B	146	LEU
1	B	155	SER
1	B	158	THR
1	B	170	ASN
1	B	217	ASP
1	B	243	ASP
1	B	287	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	A	130	ASN
1	A	142	GLN
1	A	191	ASN
1	A	201	GLN
1	A	215	GLN
1	A	258	ASN
1	A	262	HIS
1	B	133	GLN
1	B	142	GLN
1	B	164	ASN
1	B	191	ASN
1	B	287	ASN
1	B	308	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	D5X	A	501	-	10,13,13	1.66	3 (30%)	9,20,20	1.55	1 (11%)
2	D5X	B	401	-	10,13,13	1.62	2 (20%)	9,20,20	1.38	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	D5X	A	501	-	-	3/11/18/18	-
2	D5X	B	401	-	-	3/11/18/18	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	D5X	P-O5	-2.96	1.50	1.60
2	B	401	D5X	O3-C3	2.69	1.43	1.39
2	A	501	D5X	C1-C2	2.48	1.56	1.50
2	B	401	D5X	P-O3P	2.10	1.63	1.54
2	A	501	D5X	P-O3P	2.04	1.62	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	D5X	O3P-P-O5	3.82	116.89	106.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	D5X	O1P-P-O3P	2.42	116.89	107.64

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	D5X	C4-C5-O5-P
2	A	501	D5X	C4-C5-O5-P
2	A	501	D5X	O2-C2-C3-O3
2	A	501	D5X	O2-C2-C3-O3 <sup>3</sup>
2	B	401	D5X	O2-C2-C3-O3
2	B	401	D5X	O2-C2-C3-O3 <sup>3</sup>

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	249/266 (93%)	0.13	19 (7%) 13 18	22, 33, 102, 126	0
1	B	250/266 (93%)	0.13	20 (8%) 12 16	19, 32, 99, 139	0
All	All	499/532 (93%)	0.13	39 (7%) 13 17	19, 33, 102, 139	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	93	LEU	12.9
1	B	158	THR	11.0
1	B	95	ASP	11.0
1	B	94	ALA	10.5
1	A	165	ALA	10.3
1	A	96	ALA	9.3
1	B	96	ALA	7.9
1	A	95	ASP	6.1
1	B	159	GLY	5.8
1	B	162	GLN	5.6
1	B	166	ALA	5.6
1	A	94	ALA	5.4
1	B	167	CYS	5.4
1	B	317	SER	5.1
1	B	160	ILE	5.1
1	A	160	ILE	5.0
1	A	92	GLY	5.0
1	B	165	ALA	4.5
1	B	97	ASP	4.1
1	A	70	GLY	4.1
1	B	154	GLY	4.0
1	A	163	LEU	3.9
1	A	167	CYS	3.9
1	A	166	ALA	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	97	ASP	3.8
1	A	317	SER	3.7
1	B	68	PHE	3.6
1	A	164	ASN	3.5
1	B	164	ASN	3.1
1	A	159	GLY	3.0
1	A	71	CYS	2.8
1	B	155	SER	2.7
1	A	162	GLN	2.6
1	A	69	GLU	2.6
1	B	93	LEU	2.5
1	B	98	VAL	2.5
1	A	155	SER	2.4
1	B	163	LEU	2.4
1	B	161	GLY	2.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	D5X	B	401	14/14	0.98	0.14	23,40,57,59	0
2	D5X	A	501	14/14	0.99	0.12	26,38,58,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.