



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2024 – 04:38 PM EDT

PDB ID : 1L8R
Title : Structure of the Retinal Determination Protein Dachshund Reveals a DNA-Binding Motif
Authors : Kim, S.S.; Zhang, R.; Braunstein, S.E.; Joachimiak, A.; Cvekl, A.; Hegde, R.S.
Deposited on : 2002-03-21
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	101	
1	B	101	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dachshund.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	101	775	488	137	143	6	1	0	0	0
1	B	99	764	482	135	140	6	1	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	GLY	-	cloning artifact	UNP Q9UI36
A	183	SER	-	cloning artifact	UNP Q9UI36
A	190	MSE	MET	modified residue	UNP Q9UI36
B	182	GLY	-	cloning artifact	UNP Q9UI36
B	183	SER	-	cloning artifact	UNP Q9UI36
B	190	MSE	MET	modified residue	UNP Q9UI36

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	87	Total	O	0	0
			87	87		
2	B	66	Total	O	0	0
			66	66		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Dachshund

Chain A:  76% 22%



- Molecule 1: Dachshund

Chain B:  76% 18%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	28.80Å 47.73Å 50.44Å 60.75° 86.86° 84.55°	Depositor
Resolution (Å)	50.00 – 1.65	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-1.65)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.237 , 0.259	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1692	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/783	0.56	0/1054
1	B	0.32	0/772	0.56	0/1039
All	All	0.33	0/1555	0.56	0/2093

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	775	0	794	18	0
1	B	764	0	785	21	0
2	A	87	0	0	2	0
2	B	66	0	0	3	0
All	All	1692	0	1579	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:ASN:HD22	1:B:185:ASN:H	1.10	0.97

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:222:LEU:HD21	2:B:43:HOH:O	1.78	0.83
1:A:209:ILE:HD13	1:A:267:ILE:O	1.90	0.71
1:B:185:ASN:H	1:B:185:ASN:ND2	1.89	0.66
1:B:193:LEU:HD22	1:B:209:ILE:HD12	1.76	0.66
1:B:224:GLY:HA3	1:B:228:THR:OG1	1.98	0.64
1:A:209:ILE:HD13	1:A:209:ILE:H	1.63	0.64
1:A:252:ARG:HD3	1:A:261:VAL:O	1.98	0.63
1:B:190:MSE:HE1	1:B:254:LEU:HD22	1.81	0.62
1:B:185:ASN:HD22	1:B:185:ASN:N	1.91	0.61
1:B:193:LEU:HD22	1:B:209:ILE:CD1	2.35	0.57
1:B:188:CYS:HB3	2:B:129:HOH:O	2.05	0.56
1:B:262:ASN:H	1:B:262:ASN:HD22	1.55	0.55
1:A:211:LEU:HB3	1:A:212:PRO:HD3	1.90	0.53
1:A:222:LEU:HD13	1:A:280:THR:HG21	1.92	0.52
1:A:222:LEU:HD13	1:A:280:THR:CG2	2.40	0.51
1:A:209:ILE:HD13	1:A:209:ILE:N	2.27	0.50
1:B:262:ASN:H	1:B:262:ASN:ND2	2.10	0.49
1:A:230:TYR:O	1:A:234:LYS:HG3	2.12	0.49
1:B:244:ASN:H	1:B:247:GLN:NE2	2.11	0.48
1:B:240:PRO:HB2	1:B:265:LYS:HD3	1.94	0.48
1:B:206:CYS:HB3	2:B:95:HOH:O	2.13	0.48
1:A:252:ARG:HG2	1:A:257:ILE:HG13	1.96	0.47
1:A:234:LYS:HD2	2:A:118:HOH:O	2.14	0.47
1:B:188:CYS:HB2	1:B:201:PHE:CE1	2.51	0.46
1:B:184:GLN:HB2	1:B:185:ASN:H	1.64	0.45
1:B:189:LYS:HG2	1:B:190:MSE:N	2.33	0.43
1:A:209:ILE:HD12	1:A:269:ARG:HG3	2.00	0.43
1:B:262:ASN:HD22	1:B:262:ASN:N	2.12	0.42
1:B:262:ASN:HD21	1:B:263:ARG:NH1	2.17	0.42
1:B:229:VAL:HG13	1:B:275:LEU:HD21	2.01	0.41
1:A:269:ARG:O	1:A:273:GLU:HG2	2.20	0.41
1:B:262:ASN:ND2	1:B:262:ASN:N	2.69	0.41
1:A:219:LEU:HB3	1:A:223:VAL:HG12	2.02	0.41
1:A:269:ARG:HG3	2:A:108:HOH:O	2.20	0.41
1:A:215:PHE:HZ	1:A:225:GLY:H	1.68	0.40
1:A:235:ARG:NH2	1:A:278:ASP:OD1	2.54	0.40
1:A:189:LYS:C	1:A:190:MSE:HE2	2.41	0.40
1:A:233:LEU:HD13	1:A:275:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/101 (98%)	95 (96%)	3 (3%)	1 (1%)	13	2
1	B	97/101 (96%)	95 (98%)	2 (2%)	0	100	100
All	All	196/202 (97%)	190 (97%)	5 (3%)	1 (0%)	25	11

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/87 (99%)	84 (98%)	2 (2%)	45	22
1	B	85/87 (98%)	80 (94%)	5 (6%)	16	3
All	All	171/174 (98%)	164 (96%)	7 (4%)	26	7

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	209	ILE
1	A	233	LEU
1	B	184	GLN
1	B	185	ASN
1	B	190	MSE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	204	GLU
1	B	262	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	184	GLN
1	B	185	ASN
1	B	247	GLN
1	B	262	ASN
1	B	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.