



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 12:59 pm GMT

PDB ID : 5L9G  
Title : Crystal Structure of the PBP MotA in complex with mannopine from *A. tumefaciens* B6  
Authors : Marty, L.; Morera, S.  
Deposited on : 2016-06-10  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	
1	B	351	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	407	-	-	X	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5481 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called periplasmic binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	325	Total	C	N	O	S	0	0	0
			2499	1588	421	481	9			
1	B	324	Total	C	N	O	S	0	0	0
			2492	1585	419	479	9			

There are 54 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	10	MET	-	initiating methionine	UNP A0A109C781
A	11	GLY	-	expression tag	UNP A0A109C781
A	12	SER	-	expression tag	UNP A0A109C781
A	13	SER	-	expression tag	UNP A0A109C781
A	14	HIS	-	expression tag	UNP A0A109C781
A	15	HIS	-	expression tag	UNP A0A109C781
A	16	HIS	-	expression tag	UNP A0A109C781
A	17	HIS	-	expression tag	UNP A0A109C781
A	18	HIS	-	expression tag	UNP A0A109C781
A	19	HIS	-	expression tag	UNP A0A109C781
A	20	SER	-	expression tag	UNP A0A109C781
A	21	SER	-	expression tag	UNP A0A109C781
A	22	GLY	-	expression tag	UNP A0A109C781
A	23	LEU	-	expression tag	UNP A0A109C781
A	24	VAL	-	expression tag	UNP A0A109C781
A	25	PRO	-	expression tag	UNP A0A109C781
A	26	ARG	-	expression tag	UNP A0A109C781
A	27	GLY	-	expression tag	UNP A0A109C781
A	28	SER	-	expression tag	UNP A0A109C781
A	29	HIS	-	expression tag	UNP A0A109C781
A	30	MET	-	expression tag	UNP A0A109C781
A	355	HIS	-	expression tag	UNP A0A109C781
A	356	HIS	-	expression tag	UNP A0A109C781
A	357	HIS	-	expression tag	UNP A0A109C781
A	358	HIS	-	expression tag	UNP A0A109C781

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	359	HIS	-	expression tag	UNP A0A109C781
A	360	HIS	-	expression tag	UNP A0A109C781
B	10	MET	-	initiating methionine	UNP A0A109C781
B	11	GLY	-	expression tag	UNP A0A109C781
B	12	SER	-	expression tag	UNP A0A109C781
B	13	SER	-	expression tag	UNP A0A109C781
B	14	HIS	-	expression tag	UNP A0A109C781
B	15	HIS	-	expression tag	UNP A0A109C781
B	16	HIS	-	expression tag	UNP A0A109C781
B	17	HIS	-	expression tag	UNP A0A109C781
B	18	HIS	-	expression tag	UNP A0A109C781
B	19	HIS	-	expression tag	UNP A0A109C781
B	20	SER	-	expression tag	UNP A0A109C781
B	21	SER	-	expression tag	UNP A0A109C781
B	22	GLY	-	expression tag	UNP A0A109C781
B	23	LEU	-	expression tag	UNP A0A109C781
B	24	VAL	-	expression tag	UNP A0A109C781
B	25	PRO	-	expression tag	UNP A0A109C781
B	26	ARG	-	expression tag	UNP A0A109C781
B	27	GLY	-	expression tag	UNP A0A109C781
B	28	SER	-	expression tag	UNP A0A109C781
B	29	HIS	-	expression tag	UNP A0A109C781
B	30	MET	-	expression tag	UNP A0A109C781
B	355	HIS	-	expression tag	UNP A0A109C781
B	356	HIS	-	expression tag	UNP A0A109C781
B	357	HIS	-	expression tag	UNP A0A109C781
B	358	HIS	-	expression tag	UNP A0A109C781
B	359	HIS	-	expression tag	UNP A0A109C781
B	360	HIS	-	expression tag	UNP A0A109C781

- Molecule 2 is mannopine (three-letter code: MO0) (formula: C<sub>11</sub>H<sub>22</sub>N<sub>2</sub>O<sub>8</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	N	O	0	0
			21	11	2	8		
2	B	1	Total	C	N	O	0	0
			21	11	2	8		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Ca 2 2	0	0
4	B	2	Total Ca 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	196	Total O 196 196	0	0
5	B	200	Total O 200 200	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.22Å 71.22Å 135.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.00 – 1.75 49.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.00-1.75) 99.8 (49.00-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.37 (at 1.75Å)	Xtrriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.177 , 0.208 0.179 , 0.208	Depositor DCC
$R_{free}$ test set	3375 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.8	Xtrriage
Anisotropy	0.006	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.053 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5481	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CA, MOO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/2561	0.58	0/3482
1	B	0.52	1/2555 (0.0%)	0.59	0/3475
All	All	0.51	1/5116 (0.0%)	0.59	0/6957

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	57	LEU	C-N	-8.17	1.15	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2499	0	2421	11	0
1	B	2492	0	2409	15	0
2	A	21	0	0	0	0
2	B	21	0	0	0	0
3	A	24	0	36	8	0
3	B	24	0	34	8	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	196	0	0	0	0
5	B	200	0	0	1	0
All	All	5481	0	4900	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:ASN:HD21	3:A:406:EDO:H12	1.33	0.92
1:B:295:PHE:CD1	3:B:407:EDO:H12	2.09	0.87
1:A:301:ASN:ND2	3:A:406:EDO:H12	2.02	0.74
1:A:284:LYS:HA	3:A:406:EDO:H21	1.69	0.73
1:A:61:TYR:HE1	3:A:403:EDO:H22	1.53	0.73
1:B:75:GLN:HE22	1:B:83:TRP:H	1.40	0.67
1:B:167:GLN:HG2	3:B:405:EDO:H12	1.78	0.65
1:B:295:PHE:HA	3:B:407:EDO:H21	1.82	0.61
1:B:292:PRO:HG3	1:B:299:PRO:HD3	1.90	0.54
1:A:292:PRO:HG3	1:A:299:PRO:HD3	1.90	0.54
1:B:103:GLU:HG2	1:B:267:LYS:HA	1.93	0.51
1:B:75:GLN:NE2	1:B:82:THR:H	2.08	0.50
1:A:168:PRO:CD	3:A:402:EDO:H12	2.45	0.47
1:A:181:THR:HG23	3:A:407:EDO:H12	1.96	0.47
1:B:241:GLY:HA3	3:B:407:EDO:H11	1.98	0.45
1:B:167:GLN:CG	3:B:405:EDO:H12	2.45	0.44
1:A:163:ALA:HB2	1:A:229:VAL:HG11	2.00	0.42
1:B:163:ALA:HB2	1:B:229:VAL:HG11	2.00	0.42
1:A:167:GLN:HG2	3:A:402:EDO:H21	2.02	0.42
1:B:68:ASN:HD21	3:B:405:EDO:H11	1.84	0.42
3:B:407:EDO:H22	5:B:607:HOH:O	2.20	0.42
1:A:344:SER:OG	3:A:402:EDO:H11	2.20	0.42
1:B:289:LYS:HB2	1:B:308:ALA:HB2	2.03	0.41
1:B:241:GLY:CA	3:B:407:EDO:H11	2.50	0.41
1:B:164:LEU:CD2	1:B:233:ILE:HD12	2.51	0.41
1:B:138:LYS:HD2	1:B:250:ALA:HB2	2.02	0.41
1:A:289:LYS:HB2	1:A:308:ALA:HB2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/351 (92%)	317 (98%)	6 (2%)	0	100	100
1	B	322/351 (92%)	315 (98%)	7 (2%)	0	100	100
All	All	645/702 (92%)	632 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/274 (92%)	250 (100%)	1 (0%)	91	87
1	B	250/274 (91%)	248 (99%)	2 (1%)	81	72
All	All	501/548 (91%)	498 (99%)	3 (1%)	86	79

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	352	LEU
1	B	109	ILE
1	B	273	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	GLN
1	A	240	GLN
1	A	244	ASN
1	A	271	HIS
1	A	334	ASN
1	B	75	GLN
1	B	167	GLN
1	B	240	GLN
1	B	244	ASN
1	B	334	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	B	405	-	3,3,3	0.46	0	2,2,2	0.55	0
3	EDO	A	402	-	3,3,3	0.31	0	2,2,2	0.31	0
3	EDO	A	403	-	3,3,3	0.63	0	2,2,2	0.29	0
3	EDO	B	403	-	3,3,3	0.58	0	2,2,2	0.25	0
3	EDO	B	404	-	3,3,3	0.66	0	2,2,2	0.12	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	B	406	-	3,3,3	0.53	0	2,2,2	0.37	0
3	EDO	A	406	-	3,3,3	0.54	0	2,2,2	0.15	0
3	EDO	A	407	-	3,3,3	0.57	0	2,2,2	0.50	0
3	EDO	A	404	-	3,3,3	0.65	0	2,2,2	0.15	0
2	MO0	B	401	-	20,20,20	1.65	3 (15%)	23,26,26	1.37	3 (13%)
3	EDO	B	402	-	3,3,3	0.63	0	2,2,2	0.22	0
2	MO0	A	401	4	20,20,20	1.75	3 (15%)	23,26,26	1.24	2 (8%)
3	EDO	A	405	-	3,3,3	0.59	0	2,2,2	0.18	0
3	EDO	B	407	-	3,3,3	1.20	0	2,2,2	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	405	-	-	0/1/1/1	-
3	EDO	A	402	-	-	1/1/1/1	-
3	EDO	A	403	-	-	1/1/1/1	-
3	EDO	B	403	-	-	0/1/1/1	-
3	EDO	B	404	-	-	0/1/1/1	-
3	EDO	B	406	-	-	0/1/1/1	-
3	EDO	A	406	-	-	0/1/1/1	-
3	EDO	A	407	-	-	0/1/1/1	-
3	EDO	A	404	-	-	0/1/1/1	-
2	MO0	B	401	-	-	3/28/28/28	-
3	EDO	B	402	-	-	1/1/1/1	-
2	MO0	A	401	4	-	7/28/28/28	-
3	EDO	A	405	-	-	0/1/1/1	-
3	EDO	B	407	-	-	0/1/1/1	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	MO0	OE1-CD	5.55	1.40	1.24
2	B	401	MO0	OE1-CD	5.44	1.40	1.24
2	A	401	MO0	CD-NE2	-2.50	1.24	1.32
2	B	401	MO0	CD-NE2	-2.43	1.24	1.32
2	B	401	MO0	OAC-CAG	-2.23	1.38	1.43
2	A	401	MO0	OAC-CAG	-2.07	1.39	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	MO0	CG-CD-NE2	2.50	124.28	116.51
2	A	401	MO0	CG-CD-NE2	2.32	123.72	116.51
2	B	401	MO0	O-C-CA	2.20	120.72	113.40
2	A	401	MO0	O-C-OXT	-2.19	119.12	124.09
2	B	401	MO0	O-C-OXT	-2.08	119.37	124.09

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	MO0	CAI-CAJ-CAK-N
2	A	401	MO0	OAE-CAJ-CAK-N
2	B	401	MO0	CAI-CAJ-CAK-N
2	B	401	MO0	OAE-CAJ-CAK-N
2	A	401	MO0	OAB-CAF-CAG-OAC
3	A	403	EDO	O1-C1-C2-O2
3	B	402	EDO	O1-C1-C2-O2
2	A	401	MO0	O-C-CA-CB
2	A	401	MO0	CB-CA-N-CAK
2	B	401	MO0	CB-CA-N-CAK
2	A	401	MO0	OAB-CAF-CAG-CAH
3	A	402	EDO	O1-C1-C2-O2
2	A	401	MO0	OXT-C-CA-CB


There are no ring outliers.

6 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	405	EDO	3	0
3	A	402	EDO	3	0
3	A	403	EDO	1	0
3	A	406	EDO	3	0
3	A	407	EDO	1	0
3	B	407	EDO	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	57:LEU	C	58:ASN	N	1.15

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	325/351 (92%)	0.05	4 (1%) 79 84	28, 46, 78, 107	0
1	B	324/351 (92%)	-0.11	2 (0%) 89 92	27, 41, 66, 86	0
All	All	649/702 (92%)	-0.03	6 (0%) 84 89	27, 43, 74, 107	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	352	LEU	4.2
1	A	107	PRO	3.3
1	A	353	SER	3.1
1	B	107	PRO	2.9
1	A	354	ARG	2.7
1	B	352	LEU	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	A	404	4/4	0.64	0.27	64,64,65,65	0
3	EDO	B	403	4/4	0.70	0.23	77,77,77,77	0
3	EDO	B	404	4/4	0.73	0.19	54,55,57,59	0
3	EDO	B	402	4/4	0.79	0.21	71,71,72,72	0
3	EDO	A	403	4/4	0.81	0.27	54,57,60,61	0
3	EDO	A	405	4/4	0.83	0.16	69,69,69,70	0
3	EDO	A	406	4/4	0.87	0.23	45,46,47,48	0
3	EDO	B	406	4/4	0.87	0.30	47,50,51,54	0
3	EDO	A	407	4/4	0.89	0.22	58,59,61,62	0
3	EDO	A	402	4/4	0.93	0.22	44,45,47,50	0
2	MO0	A	401	21/21	0.94	0.09	28,33,37,37	0
3	EDO	B	407	4/4	0.94	0.18	30,38,38,39	0
4	CA	A	409	1/1	0.94	0.08	49,49,49,49	1
3	EDO	B	405	4/4	0.96	0.28	40,41,43,47	0
2	MO0	B	401	21/21	0.97	0.09	25,28,31,33	0
4	CA	B	409	1/1	0.97	0.06	35,35,35,35	1
4	CA	A	408	1/1	0.98	0.08	38,38,38,38	0
4	CA	B	408	1/1	1.00	0.08	37,37,37,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around MO0 A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around MO0 B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.