



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 06:33 PM EDT

PDB ID : 5LFD
Title : Crystal structure of allantoin racemase from *Pseudomonas fluorescens* AllR
Authors : Cendron, I.; Zanotti, G.; Percudani, R.; Ramazzina, I.; Puggioni, V.; Maccacaro, E.; Liuzzi, A.; Secchi, A.
Deposited on : 2016-07-01
Resolution : 2.15 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	 4% 81% 13% . .
1	B	242	 5% 83% 12% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Allantoin racemase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	235	1749	1096	301	342	10	0	0	0
1	B	235	1749	1096	301	342	10	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	145	145	145	0	0
2	B	143	143	143	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	109.72Å 109.72Å 109.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.83 – 2.15 44.79 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.8 (44.83-2.15) 96.8 (44.79-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	12.19 (at 2.16Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.202 , 0.264 0.202 , 0.262	Depositor DCC
R_{free} test set	1197 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.037 for l,-k,h	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3786	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.85	2/1770 (0.1%)	0.92	7/2406 (0.3%)
1	B	0.79	1/1770 (0.1%)	0.97	5/2406 (0.2%)
All	All	0.82	3/3540 (0.1%)	0.94	12/4812 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	CYS	CB-SG	7.79	1.95	1.82
1	B	75	GLY	C-O	6.60	1.34	1.23
1	A	180	CYS	N-CA	5.20	1.56	1.46

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	214	ARG	NE-CZ-NH2	-6.82	116.89	120.30
1	A	214	ARG	NE-CZ-NH1	6.64	123.62	120.30
1	B	180	CYS	N-CA-CB	5.95	121.32	110.60
1	B	214	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	A	180	CYS	N-CA-CB	5.72	120.90	110.60
1	A	214	ARG	NE-CZ-NH2	-5.66	117.47	120.30
1	A	84	LEU	CB-CG-CD1	5.46	120.29	111.00
1	B	93	VAL	CG1-CB-CG2	5.36	119.48	110.90
1	A	23	ARG	NE-CZ-NH2	-5.33	117.64	120.30
1	A	93	VAL	CG1-CB-CG2	5.32	119.41	110.90
1	B	118	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	A	84	LEU	CA-CB-CG	5.03	126.88	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1749	0	1783	21	0
1	B	1749	0	1783	17	0
2	A	145	0	0	3	0
2	B	143	0	0	4	0
All	All	3786	0	3566	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:GLY:O	2:B:301:HOH:O	1.60	1.20
1:B:58:MET:HG2	1:B:84:LEU:HD13	1.59	0.84
1:A:98:ALA:O	1:A:102:THR:HG23	1.79	0.81
1:A:75:GLY:O	2:A:301:HOH:O	2.02	0.77
1:B:191:ARG:NH2	2:B:303:HOH:O	2.22	0.73
1:B:102:THR:HG21	1:B:204:ALA:HB1	1.72	0.70
1:A:115:THR:HB	1:A:144:MET:O	1.95	0.65
1:B:64:TYR:CE2	1:B:66:GLN:HB2	2.32	0.64
1:B:98:ALA:O	1:B:102:THR:HG23	1.99	0.63
1:A:157:MET:HE2	1:A:186:LEU:HD22	1.80	0.63
1:A:102:THR:HG22	2:A:430:HOH:O	2.03	0.59
1:A:102:THR:HG21	1:A:204:ALA:HB1	1.85	0.59
1:A:58:MET:HG2	1:A:84:LEU:HD13	1.87	0.56
1:A:227:THR:O	2:A:303:HOH:O	2.18	0.56
1:A:64:TYR:CE2	1:A:66:GLN:HB2	2.42	0.55
1:B:98:ALA:O	1:B:102:THR:CG2	2.57	0.53
1:A:214:ARG:HD2	2:B:385:HOH:O	2.10	0.51
1:A:157:MET:CE	1:A:186:LEU:HD22	2.41	0.50
1:B:44:VAL:HG11	1:B:78:GLU:HG2	1.92	0.50
1:B:72:ILE:HA	1:B:93:VAL:O	2.14	0.48
1:A:188:GLU:OE1	1:A:191:ARG:NH1	2.41	0.48
1:B:62:MET:CE	1:B:88:LEU:HD21	2.44	0.48
1:B:49:GLU:OE2	2:B:302:HOH:O	2.20	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:VAL:HG12	1:A:62:MET:HE2	1.96	0.47
1:A:210:GLU:O	1:A:214:ARG:HG3	2.17	0.45
1:A:62:MET:CE	1:A:88:LEU:HD21	2.47	0.45
1:B:86:GLU:HG2	1:B:225:TYR:HA	1.99	0.45
1:A:61:VAL:HG12	1:A:62:MET:CE	2.47	0.44
1:A:86:GLU:HG2	1:A:225:TYR:HA	2.00	0.43
1:B:62:MET:CE	1:B:88:LEU:CD2	2.97	0.42
1:A:157:MET:O	1:A:161:VAL:HG12	2.19	0.42
1:A:76:TYR:CE1	1:A:96:THR:HA	2.54	0.42
1:B:157:MET:O	1:B:161:VAL:HG13	2.20	0.42
1:A:127:LEU:HA	1:A:127:LEU:HD23	1.81	0.41
1:B:64:TYR:HE2	1:B:66:GLN:HB2	1.80	0.41
1:B:181:GLY:C	1:B:183:MET:H	2.25	0.40
1:B:126:ARG:HD2	1:B:126:ARG:HA	1.86	0.40
1:A:110:TYR:CZ	1:A:136:CYS:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/242 (96%)	224 (96%)	7 (3%)	2 (1%)	17	11
1	B	233/242 (96%)	223 (96%)	9 (4%)	1 (0%)	34	29
All	All	466/484 (96%)	447 (96%)	16 (3%)	3 (1%)	25	18

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	151	GLU
1	A	223	ARG
1	B	146	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/189 (97%)	173 (94%)	11 (6%)	19	14
1	B	184/189 (97%)	172 (94%)	12 (6%)	17	12
All	All	368/378 (97%)	345 (94%)	23 (6%)	18	13

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	ILE
1	A	76	TYR
1	A	84	LEU
1	A	93	VAL
1	A	102	THR
1	A	158	GLU
1	A	161	VAL
1	A	175	VAL
1	A	180	CYS
1	A	187	ASP
1	A	233	VAL
1	B	65	ASP
1	B	76	TYR
1	B	84	LEU
1	B	93	VAL
1	B	102	THR
1	B	149	LEU
1	B	158	GLU
1	B	175	VAL
1	B	180	CYS
1	B	186	LEU
1	B	232	LYS
1	B	233	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	235/242 (97%)	-0.27	9 (3%) 40 49	8, 17, 49, 92	0
1	B	235/242 (97%)	-0.12	11 (4%) 31 41	8, 18, 52, 130	0
All	All	470/484 (97%)	-0.19	20 (4%) 35 45	8, 18, 51, 130	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	147	LEU	9.6
1	A	147	LEU	6.4
1	B	146	VAL	6.0
1	B	149	LEU	5.4
1	B	150	GLU	4.4
1	A	146	VAL	3.8
1	B	151	GLU	3.3
1	A	180	CYS	3.3
1	A	149	LEU	3.2
1	A	150	GLU	3.0
1	A	76	TYR	2.8
1	B	184	ALA	2.7
1	B	76	TYR	2.5
1	B	180	CYS	2.5
1	B	182	GLY	2.4
1	A	145	ALA	2.4
1	B	144	MET	2.2
1	A	151	GLU	2.1
1	B	153	PRO	2.1
1	A	154	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.