



## Full wwPDB EM Validation Report ⓘ

Apr 1, 2021 – 10:19 AM EDT

PDB ID : 7LJ0  
EMDB ID : EMD-9976  
Title : Linker 3 and scaffolded phycoerythrin beta subunit from the phycobilisome of  
Porphyridium purpureum  
Authors : Rathbone, H.W.; Landsberg, M.J.; Michie, K.A.; Green, B.R.; Curmi, P.M.G.  
Deposited on : 2021-01-28  
Resolution : 2.80 Å (reported)  
Based on initial models : 3V57, 4LMX

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev75  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.18


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	
2	B	177	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Linker 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	27	217	140	40	36	1	0	0

- Molecule 2 is a protein called B-phycoerythrin beta chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	145	1060	651	186	212	11	0	0

- Molecule 3 is PHYCOERYTHROBILIN (three-letter code: PEB) (formula:  $C_{33}H_{40}N_4O_6$ ).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
3	B	1	129	99	12	18	0
3	B	1	129	99	12	18	0
3	B	1	129	99	12	18	0



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	191825	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	48	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.143	Depositor
Minimum map value	-0.087	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.02	Depositor
Map size ( $\text{\AA}$ )	610.95996, 610.95996, 610.95996	wwPDB
Map dimensions	560, 560, 560	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.091, 1.091, 1.091	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEB, MEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/221	0.94	0/301
2	B	0.75	0/1060	0.85	0/1430
All	All	0.74	0/1281	0.86	0/1731

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	217	0	229	7	0
2	B	1060	0	1055	25	0
3	B	129	0	110	7	0
All	All	1406	0	1394	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:VAL:N	2:B:108:ARG:HD3	1.64	1.13
2:B:64:PRO:O	2:B:67:ILE:HG12	1.70	0.90
2:B:92:TYR:CD2	2:B:109:CYS:HB2	2.22	0.75
3:B:201:PEB:HNA	3:B:201:PEB:HMB2	1.52	0.73
1:A:268:VAL:N	2:B:108:ARG:HH11	1.91	0.68
1:A:268:VAL:N	2:B:108:ARG:CD	2.50	0.64
3:B:202:PEB:HNA	3:B:202:PEB:HMB2	1.64	0.62
2:B:92:TYR:HE1	2:B:108:ARG:HB2	1.65	0.62
2:B:92:TYR:CE1	2:B:108:ARG:HB2	2.36	0.60
1:A:274:VAL:HG21	2:B:95:TYR:CE1	2.37	0.60
2:B:64:PRO:O	2:B:67:ILE:CG1	2.49	0.58
2:B:92:TYR:CG	2:B:109:CYS:HB2	2.38	0.57
2:B:92:TYR:CE2	2:B:109:CYS:HB2	2.39	0.57
2:B:157:ASP:OD1	2:B:159:THR:HG23	2.09	0.53
3:B:202:PEB:HBA3	3:B:202:PEB:HHA1	1.90	0.52
2:B:52:VAL:HG21	2:B:87:GLU:HG2	1.90	0.52
3:B:202:PEB:HMB2	3:B:202:PEB:NA	2.27	0.48
3:B:202:PEB:HBD1	3:B:202:PEB:OD	2.14	0.47
1:A:290:ILE:CG2	2:B:91:ARG:NH1	2.79	0.46
1:A:274:VAL:CG2	2:B:95:TYR:CE1	2.99	0.45
2:B:92:TYR:CD2	2:B:109:CYS:SG	3.10	0.45
2:B:52:VAL:HG11	2:B:87:GLU:HG3	2.00	0.44
2:B:128:ILE:HG12	2:B:173:GLY:HA2	1.99	0.43
2:B:47:ASN:OD1	2:B:149:ARG:HD3	2.19	0.42
2:B:142:ILE:O	3:B:202:PEB:HAA2	2.20	0.42
2:B:110:LEU:HD23	2:B:110:LEU:HA	1.82	0.42
1:A:268:VAL:N	2:B:108:ARG:NH1	2.62	0.41
2:B:92:TYR:CD2	2:B:109:CYS:CB	2.98	0.41
2:B:72:MEN:HE22	3:B:201:PEB:HBB2	2.03	0.41
2:B:146:ALA:O	2:B:150:LYS:HE2	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	25/316 (8%)	25 (100%)	0	0	100	100
2	B	142/177 (80%)	141 (99%)	1 (1%)	0	100	100
All	All	167/493 (34%)	166 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	24/256 (9%)	24 (100%)	0	100	100
2	B	113/137 (82%)	113 (100%)	0	100	100
All	All	137/393 (35%)	137 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MEN	B	72	2	7,8,9	0.69	0	6,9,11	1.34	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MEN	B	72	2	-	3/7/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	72	MEN	CB-CG-ND2	2.45	118.78	115.48

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	72	MEN	CA-CB-CG-OD1
2	B	72	MEN	CA-CB-CG-ND2
2	B	72	MEN	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	72	MEN	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEB	B	201	2	37,46,46	3.29	10 (27%)	39,67,67	2.47	16 (41%)
3	PEB	B	203	2	37,46,46	3.84	8 (21%)	39,67,67	2.23	16 (41%)
3	PEB	B	202	2	37,46,46	3.37	11 (29%)	39,67,67	2.08	17 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEB	B	201	2	-	2/20/74/74	0/4/4/4
3	PEB	B	203	2	-	4/20/74/74	0/4/4/4
3	PEB	B	202	2	-	3/20/74/74	0/4/4/4

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	203	PEB	CHB-C4B	17.59	1.49	1.35
3	B	201	PEB	CHB-C4B	14.85	1.47	1.35
3	B	202	PEB	CHB-C4B	14.58	1.47	1.35
3	B	203	PEB	C2D-C3D	9.57	1.46	1.34
3	B	202	PEB	C2D-C3D	8.07	1.44	1.34
3	B	201	PEB	C2D-C3D	7.82	1.44	1.34
3	B	203	PEB	C3C-C4C	6.63	1.52	1.42
3	B	201	PEB	C3B-C2B	5.44	1.48	1.36
3	B	203	PEB	C3B-C2B	5.19	1.47	1.36
3	B	201	PEB	C3C-C4C	4.96	1.49	1.42
3	B	202	PEB	C3C-C4C	4.80	1.49	1.42
3	B	202	PEB	C1A-NA	-4.57	1.31	1.37
3	B	202	PEB	CHA-C1B	4.42	1.50	1.40
3	B	201	PEB	CHA-C1B	4.30	1.50	1.40
3	B	203	PEB	CHA-C1B	4.28	1.50	1.40
3	B	202	PEB	C3B-C2B	4.19	1.45	1.36
3	B	203	PEB	C1D-ND	4.03	1.51	1.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	203	PEB	C2C-C3C	3.63	1.48	1.37
3	B	203	PEB	OD-C4D	3.55	1.30	1.23
3	B	202	PEB	OD-C4D	3.54	1.30	1.23
3	B	202	PEB	C4D-ND	3.43	1.39	1.35
3	B	202	PEB	C2C-C3C	2.93	1.46	1.37
3	B	201	PEB	C2C-C3C	2.87	1.46	1.37
3	B	202	PEB	C2A-C1A	-2.67	1.49	1.52
3	B	201	PEB	OD-C4D	2.61	1.28	1.23
3	B	201	PEB	C1A-NA	-2.35	1.34	1.37
3	B	202	PEB	C1B-C2B	2.24	1.50	1.45
3	B	201	PEB	C2A-C1A	-2.14	1.50	1.52
3	B	201	PEB	C1C-CHB	2.08	1.49	1.41

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	201	PEB	CHC-C1D-ND	-5.97	107.02	113.95
3	B	201	PEB	OA-C1A-C2A	-5.62	121.70	126.17
3	B	203	PEB	OA-C1A-C2A	-5.47	121.82	126.17
3	B	201	PEB	C3D-C4D-ND	4.78	116.64	107.26
3	B	203	PEB	OD-C4D-C3D	-4.56	119.12	129.46
3	B	201	PEB	C2A-C1A-NA	4.26	111.94	108.27
3	B	201	PEB	CAB-CBB-CGB	-4.14	105.72	112.67
3	B	203	PEB	CBC-CAC-C2C	-4.01	105.08	112.49
3	B	201	PEB	C4B-C3B-C2B	-3.99	102.36	106.78
3	B	203	PEB	CHC-C1D-ND	-3.94	109.37	113.95
3	B	202	PEB	C3D-C4D-ND	3.64	114.40	107.26
3	B	201	PEB	OD-C4D-C3D	-3.51	121.52	129.46
3	B	202	PEB	OD-C4D-C3D	-3.49	121.56	129.46
3	B	202	PEB	C3B-C4B-NB	3.48	115.10	110.05
3	B	202	PEB	CHC-C4C-C3C	-3.46	124.43	130.34
3	B	202	PEB	OA-C1A-C2A	-3.35	123.51	126.17
3	B	203	PEB	C3D-C4D-ND	3.35	113.83	107.26
3	B	202	PEB	CMD-C2D-C3D	-3.31	125.39	130.06
3	B	202	PEB	CAC-CBC-CGC	-3.26	107.20	112.67
3	B	202	PEB	CBC-CAC-C2C	-3.18	106.62	112.49
3	B	201	PEB	C3B-C4B-NB	3.14	114.61	110.05
3	B	203	PEB	CHA-C4A-NA	3.10	128.89	125.20
3	B	202	PEB	CHB-C4B-NB	-3.09	124.53	128.83
3	B	203	PEB	C4B-C3B-C2B	-2.98	103.48	106.78
3	B	203	PEB	CMB-C2B-C1B	2.97	129.64	125.06
3	B	201	PEB	CHB-C4B-NB	-2.96	124.72	128.83

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	201	PEB	CHC-C4C-C3C	-2.95	125.31	130.34
3	B	201	PEB	OD-C4D-ND	-2.84	121.72	125.93
3	B	202	PEB	CHC-C1D-ND	-2.66	110.86	113.95
3	B	202	PEB	C1B-C2B-C3B	-2.63	103.49	106.51
3	B	202	PEB	CMB-C2B-C1B	2.62	129.10	125.06
3	B	202	PEB	C4B-C3B-C2B	-2.54	103.97	106.78
3	B	203	PEB	C2A-C1A-NA	2.53	110.45	108.27
3	B	203	PEB	CBD-CAD-C3D	-2.47	115.31	127.62
3	B	203	PEB	CAB-C3B-C4B	2.46	129.37	125.01
3	B	201	PEB	CHA-C1B-NB	-2.36	119.99	124.93
3	B	203	PEB	CHA-C1B-NB	-2.32	120.09	124.93
3	B	203	PEB	OA-C1A-NA	2.30	127.72	124.94
3	B	202	PEB	OA-C1A-NA	2.28	127.71	124.94
3	B	202	PEB	CHA-C1B-NB	-2.23	120.27	124.93
3	B	203	PEB	CHC-C4C-C3C	-2.23	126.54	130.34
3	B	203	PEB	CAA-C3A-C2A	2.22	119.81	114.26
3	B	201	PEB	CHB-C4B-C3B	-2.22	120.20	125.32
3	B	202	PEB	CHB-C4B-C3B	-2.18	120.29	125.32
3	B	201	PEB	C1C-CHB-C4B	2.10	131.32	128.81
3	B	203	PEB	CAA-C3A-C4A	-2.10	107.28	112.67
3	B	201	PEB	CMB-C2B-C1B	2.10	128.30	125.06
3	B	201	PEB	C2A-C3A-C4A	2.07	104.44	101.34
3	B	202	PEB	CBD-CAD-C3D	-2.05	117.43	127.62

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	202	PEB	NB-C1B-CHA-C4A
3	B	202	PEB	C2B-C1B-CHA-C4A
3	B	203	PEB	C2D-C3D-CAD-CBD
3	B	203	PEB	C4D-C3D-CAD-CBD
3	B	203	PEB	NB-C1B-CHA-C4A
3	B	203	PEB	C2B-C1B-CHA-C4A
3	B	201	PEB	NB-C1B-CHA-C4A
3	B	201	PEB	C2B-C1B-CHA-C4A
3	B	202	PEB	C4A-C3A-CAA-CBA


There are no ring outliers.


2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	201	PEB	2	0
3	B	202	PEB	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-9976. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y



Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 280



Y Index: 280




Z Index: 280




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 305



Y Index: 292




Z Index: 275

The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 2573  $\text{nm}^3$ ; this corresponds to an approximate mass of 2324 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum i



\*Reported resolution corresponds to spatial frequency of 0.357 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-9976 and PDB model 7LJ0. Per-residue inclusion information can be found in section 3 on page 4.

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Atom inclusion [i](#)



At the recommended contour level, 11% of all backbone atoms, 8% of all non-hydrogen atoms, are inside the map.