

Full wwPDB X-ray Structure Validation Report (i)

Jun 18, 2024 – 09:45 PM EDT

PDB ID	:	4LKG
Title	:	The structure of hemagglutinin from a avian-origin H7N9 influenza virus
		(A/Shanghai/1/2013) in complex with avian receptor analog 3'SLNLN
Authors	:	Shi, Y.; Zhang, W.; Wang, F.; Qi, J.; Song, H.; Wu, Y.; Gao, F.; Zhang, Y.;
		Fan, Z.; Gong, W.; Wang, D.; Shu, Y.; Wang, Y.; Yan, J.; Gao, G.F.
Deposited on	:	2013-07-07
Resolution	:	2.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
- 1		01.0	%		
	А	316	80%	19%	•
			3%		
1	С	316	79%	20%	•
			6%		
1	Ε	316	79%	20%	•
			8%		
2	В	170	85%	14%	•
			8%		
2	D	170	85%	15%	•



Mol	Chain	Length	Qua	ality of chain
2	F	170	21% 85 ^c	% <u>14%</u> .
3	G	3	33%	67%
4	Н	2	50%	50%
4	Ι	2	50%	50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	G	1	-	-	Х	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 11531 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	216	Total	С	Ν	0	\mathbf{S}	0	0	0
	A	510	2415	1497	435	468	15	0	0	0
1	С	216	Total	С	Ν	0	S	0	0	0
	C	510	2415	1497	435	468	15	0	0	0
1	Б	216	Total	С	Ν	0	S	0	0	0
	E	510	2415	1497	435	468	15	0	0	0

• Molecule 1 is a protein called hemagglutinin.

• Molecule 2 is a protein called hemagglutinin.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
9	Р	170	Total	С	Ν	0	S	0	0	0
	D	170	1378	851	240	280	7	0	0	0
0	Л	170	Total	С	Ν	0	S	0	0	0
	D	170	1378	851	240	280	7	0	0	0
0	Б	170	Total	С	Ν	0	S	0	0	0
	Г	170	1378	851	240	280	7	0		U

• Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galacto pyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	I	Aton	ns		ZeroOcc	AltConf	Trace
3	G	3	Total 46	C 25	N 2	0 19	0	0	0

• Molecule 4 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galacto pyranose.





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	Н	2	Total 32	C 17	N 1	0 14	0	0	0
4	Ι	2	Total 32	C 17	N 1	0 14	0	0	0

• Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C N O 14 8 1 5	0	0
5	D	1	Total C N O 14 8 1 5	0	0
5	F	1	Total C N O 14 8 1 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: hemagglutinin



V293 (2294 (2294 (2298 (2310 (3310 (3311) (3311) (3312) (3



• Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-de
oxy-beta-D-glucopyranose

Chain G:	33%	67%
NAG1 GAL2 SIA3		
• Molecule 4:	N-acetyl-alpha-ne	uraminic acid-(2-3)-beta-D-galactopyranose
Chain H:	50%	50%
GAL1 SIA2		
• Molecule 4:	N-acetyl-alpha-ne	uraminic acid-(2-3)-beta-D-galactopyranose
Chain I:	50%	50%
GAL1 SIA2		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 3	Depositor
Cell constants	110.76Å 110.76Å 133.23Å	Deneriten
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
$\mathbf{P}_{\mathrm{esolution}}(\mathbf{\hat{A}})$	42.59 - 2.99	Depositor
Resolution (A)	42.59 - 2.99	EDS
% Data completeness	99.7 (42.59-2.99)	Depositor
(in resolution range)	99.7(42.59-2.99)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.21 (at 3.01 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
B B.	0.246 , 0.278	Depositor
n, n_{free}	0.238 , 0.269	DCC
R_{free} test set	1838 reflections (5.00%)	wwPDB-VP
Wilson B-factor $(Å^2)$	72.7	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.23 , 4.2	EDS
L-test for twinning ²	$< L > = 0.45, < L^2 > = 0.27$	Xtriage
	0.029 for -h,-k,l	
Estimated twinning fraction	0.109 for h,-h-k,-l	Xtriage
	0.037 for -k,-h,-l	
F_o, F_c correlation	0.92	EDS
Total number of atoms	11531	wwPDB-VP
Average B, all atoms $(Å^2)$	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 15.49% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
INIOI	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.27	0/2460	0.46	0/3323	
1	С	0.26	0/2460	0.46	0/3323	
1	Е	0.26	0/2460	0.46	0/3323	
2	В	0.25	0/1402	0.41	0/1889	
2	D	0.25	0/1402	0.42	0/1889	
2	F	0.28	0/1402	0.43	0/1889	
All	All	0.26	0/11586	0.45	0/15636	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2415	0	2366	25	1
1	С	2415	0	2366	35	0
1	Е	2415	0	2366	26	1
2	В	1378	0	1278	12	0
2	D	1378	0	1278	13	0
2	F	1378	0	1278	13	0
3	G	46	0	40	14	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes	
4	Н	32	0	28	1	0	
4	Ι	32	0	28	1	0	
5	В	14	0	13	0	0	
5	D	14	0	13	0	0	
5	F	14	0	13	0	0	
All	All	11531	0	11067	119	1	

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:267:ASP:HB3	3:G:1:NAG:H82	1.32	1.12
1:C:267:ASP:HB3	3:G:1:NAG:C8	1.96	0.96
1:A:217:GLN:HE22	3:G:2:GAL:H3	1.33	0.93
1:C:40:ARG:HG2	1:C:263:GLN:HG3	1.53	0.88
1:C:267:ASP:HB3	3:G:1:NAG:C7	2.08	0.84
1:C:267:ASP:CB	3:G:1:NAG:H82	2.13	0.77
2:F:448:ARG:HD3	2:F:480:HIS:CD2	2.26	0.70
1:E:144:LEU:HD11	1:E:185:LEU:HD22	1.75	0.69
1:A:121:ARG:NH1	1:A:145:SER:O	2.26	0.68
1:A:144:LEU:HD11	1:A:185:LEU:HD22	1.76	0.68
1:E:121:ARG:HB3	1:E:144:LEU:HB3	1.75	0.68
1:A:121:ARG:HB3	1:A:144:LEU:HB3	1.74	0.68
1:C:121:ARG:HB3	1:C:144:LEU:HB3	1.74	0.68
1:E:121:ARG:NH1	1:E:145:SER:O	2.26	0.68
1:C:267:ASP:C	3:G:1:NAG:H82	2.15	0.67
1:C:144:LEU:HD11	1:C:185:LEU:HD22	1.76	0.66
1:C:121:ARG:NH1	1:C:145:SER:O	2.27	0.66
1:E:70:LEU:O	1:E:110:LYS:NZ	2.30	0.64
1:C:70:LEU:O	1:C:110:LYS:NZ	2.31	0.64
1:A:70:LEU:O	1:A:110:LYS:NZ	2.32	0.63
4:H:2:SIA:H6	4:H:2:SIA:O1B	1.99	0.63
1:E:316:GLU:HA	2:F:333:ASN:HD22	1.65	0.61
2:F:448:ARG:HD3	2:F:480:HIS:NE2	2.17	0.59
3:G:1:NAG:H62	3:G:2:GAL:O2	2.02	0.59
1:C:316:GLU:HA	2:D:333:ASN:HD22	1.68	0.59
1:A:316:GLU:HA	2:B:333:ASN:HD22	1.68	0.58
4:I:2:SIA:H6	4:I:2:SIA:O1B	2.03	0.58
1:A:222:ASP:HB3	1:A:224:HIS:CE1	2.40	0.57



	A A	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:E:222:ASP:HB3	1:E:224:HIS:CE1	2.40	0.57
1:C:222:ASP:HB3	1:C:224:HIS:CE1	2.40	0.56
1:A:40:ARG:HG2	1:A:263:GLN:HG3	1.87	0.56
1:E:40:ARG:HG2	1:E:263:GLN:HG3	1.87	0.56
1:C:111:GLU:OE2	1:C:163:ARG:NH1	2.40	0.55
1:C:160:LYS:HE3	1:C:231:ASN:HA	1.89	0.55
1:E:111:GLU:OE2	1:E:163:ARG:NH1	2.38	0.55
1:A:111:GLU:OE2	1:A:163:ARG:NH1	2.39	0.54
1:C:267:ASP:HB3	3:G:1:NAG:N2	2.21	0.54
1:C:267:ASP:O	3:G:1:NAG:H82	2.09	0.53
1:A:160:LYS:HE3	1:A:231:ASN:HA	1.90	0.53
1:E:160:LYS:HE3	1:E:231:ASN:HA	1.90	0.53
2:D:437:ASP:O	2:D:441:GLU:HG2	2.10	0.52
2:F:437:ASP:O	2:F:441:GLU:HG2	2.10	0.52
2:B:437:ASP:O	2:B:441:GLU:HG2	2.09	0.51
1:C:207:SER:O	1:C:211:ARG:NH2	2.42	0.51
3:G:3:SIA:O1B	3:G:3:SIA:H6	2.09	0.51
1:E:293:VAL:HG11	2:F:386:LEU:HD13	1.93	0.50
2:F:461:ILE:HG22	2:F:463:HIS:H	1.77	0.49
1:C:293:VAL:HG11	2:D:386:LEU:HD13	1.95	0.49
2:F:376:LEU:HD22	2:F:420:LEU:HD21	1.95	0.48
1:A:217:GLN:HE22	3:G:2:GAL:C3	2.14	0.48
1:A:207:SER:O	1:A:211:ARG:NH2	2.45	0.48
2:D:376:LEU:HD22	2:D:420:LEU:HD21	1.96	0.48
1:A:153:PRO:O	1:A:155:MET:HG3	2.14	0.48
2:B:376:LEU:HD22	2:B:420:LEU:HD21	1.95	0.48
1:C:153:PRO:O	1:C:155:MET:HG3	2.14	0.48
1:A:47:ARG:NH1	1:A:73:SER:HB3	2.29	0.47
1:C:163:ARG:HD3	1:C:250:PHE:CE2	2.50	0.47
2:D:461:ILE:HG22	2:D:463:HIS:H	1.78	0.47
2:B:461:ILE:HG22	2:B:463:HIS:H	1.78	0.47
1:E:124:GLY:HA3	1:E:142:TRP:HB3	1.97	0.47
1:E:207:SER:O	1:E:211:ARG:NH2	2.44	0.47
1:E:153:PRO:O	1:E:155:MET:HG3	2.15	0.46
1:E:35:ARG:NH1	1:E:303:ARG:O	2.47	0.46
1:A:163:ARG:HD3	1:A:250:PHE:CE2	2.50	0.46
1:A:35:ARG:NH1	1:A:303:ARG:O	2.48	0.46
1:E:106:GLY:HA2	1:E:255:SER:HB3	1.98	0.46
2:D:327:ILE:N	2:D:433:ASP:OD1	2.42	0.46
1:E:163:ARG:HD3	1:E:250:PHE:CE2	2.50	0.46
1:E:47:ARG:NH1	1:E:73:SER:HB3	2.30	0.45



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:106:GLY:HA2	1:A:255:SER:HB3	1.98	0.45
1:C:35:ARG:NH1	1:C:303:ARG:O	2.49	0.45
1:C:106:GLY:HA2	1:C:255:SER:HB3	1.99	0.45
2:F:327:ILE:N	2:F:433:ASP:OD1	2.42	0.45
1:E:7:HIS:CD2	2:F:327:ILE:HG12	2.51	0.45
2:D:330:PHE:CD1	2:D:331:ILE:HG13	2.52	0.45
2:D:448:ARG:HE	2:D:448:ARG:HB3	1.61	0.45
1:E:163:ARG:HD3	1:E:250:PHE:CZ	2.52	0.45
1:C:47:ARG:NH1	1:C:73:SER:HB3	2.32	0.45
1:A:293:VAL:HG11	2:B:386:LEU:HD13	1.97	0.45
2:F:330:PHE:CD1	2:F:331:ILE:HG13	2.52	0.45
1:C:117:TYR:HB3	1:C:120:ILE:HD11	1.99	0.44
1:C:267:ASP:CB	3:G:1:NAG:HN2	2.30	0.44
1:A:124:GLY:HA3	1:A:142:TRP:HB3	1.99	0.44
1:C:124:GLY:HA3	1:C:142:TRP:HB3	1.98	0.44
2:B:322:GLY:HA3	2:B:433:ASP:OD2	2.18	0.44
2:D:322:GLY:HA3	2:D:433:ASP:OD2	2.18	0.43
1:A:105:SER:HB2	1:A:251:LEU:HD22	2.00	0.43
1:A:117:TYR:HB3	1:A:120:ILE:HD11	2.00	0.43
2:F:379:LYS:HE3	2:F:379:LYS:HB2	1.82	0.43
1:A:163:ARG:HD3	1:A:250:PHE:CZ	2.53	0.43
2:B:330:PHE:CD1	2:B:331:ILE:HG13	2.52	0.43
1:E:117:TYR:HB3	1:E:120:ILE:HD11	2.00	0.43
1:C:105:SER:HB2	1:C:251:LEU:HD22	2.01	0.43
1:C:267:ASP:HB3	3:G:1:NAG:HN2	1.84	0.43
2:D:369:ILE:HD11	2:D:428:THR:HG23	2.01	0.43
2:B:448:ARG:HE	2:B:448:ARG:HB3	1.61	0.42
1:E:52:GLY:HA2	1:E:80:ARG:HG3	2.00	0.42
2:F:322:GLY:HA3	2:F:433:ASP:OD2	2.19	0.42
1:C:52:GLY:HA2	1:C:80:ARG:HG3	2.01	0.42
1:C:163:ARG:HD3	1:C:250:PHE:CZ	2.53	0.42
1:E:199:ASN:OD1	1:E:229:ASN:ND2	2.51	0.42
1:A:282:ASN:HB3	2:B:377:ILE:HG23	2.02	0.42
1:E:105:SER:HB2	1:E:251:LEU:HD22	2.02	0.42
1:C:199:ASN:OD1	1:C:229:ASN:ND2	2.49	0.41
2:F:369:ILE:HD11	2:F:428:THR:HG23	2.02	0.41
1:C:57:LEU:HD23	1:C:102:LEU:HD12	2.03	0.41
2:B:369:ILE:HD11	2:B:428:THR:HG23	2.01	0.41
1:C:267:ASP:CA	3:G:1:NAG:H82	2.51	0.41
1:A:52:GLY:HA2	1:A:80:ARG:HG3	2.02	0.41
1:C:35:ARG:HD3	1:C:303:ARG:HG2	2.03	0.41



4LKG

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:E:232:ASP:OD1	1:E:233:THR:N	2.50	0.41
2:B:379:LYS:HE3	2:B:379:LYS:HB2	1.82	0.41
1:E:35:ARG:HD3	1:E:303:ARG:HG2	2.02	0.41
1:E:57:LEU:HD23	1:E:102:LEU:HD12	2.03	0.41
1:A:57:LEU:HD23	1:A:102:LEU:HD12	2.03	0.41
2:D:360:LYS:HB2	2:D:360:LYS:HE3	1.86	0.41
2:B:326:ALA:HB3	2:B:433:ASP:OD1	2.22	0.40
2:D:331:ILE:HG13	2:D:331:ILE:H	1.74	0.40
1:C:7:HIS:CD2	2:D:327:ILE:HG12	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:GLU:OE1	$1:E:164:LYS:NZ[2_545]$	2.11	0.09

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	314/316~(99%)	292 (93%)	22 (7%)	0	100	100
1	С	314/316~(99%)	291 (93%)	23~(7%)	0	100	100
1	Е	314/316~(99%)	291 (93%)	23~(7%)	0	100	100
2	В	168/170~(99%)	159~(95%)	9~(5%)	0	100	100
2	D	168/170~(99%)	159~(95%)	9~(5%)	0	100	100
2	F	168/170~(99%)	159(95%)	9(5%)	0	100	100
All	All	1446/1458~(99%)	1351 (93%)	95~(7%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	265/265~(100%)	240~(91%)	25~(9%)	8 32
1	С	265/265~(100%)	238~(90%)	27 (10%)	7 28
1	Ε	265/265~(100%)	239~(90%)	26 (10%)	8 30
2	В	145/145~(100%)	135~(93%)	10 (7%)	15 48
2	D	145/145~(100%)	135~(93%)	10 (7%)	15 48
2	F	145/145~(100%)	135~(93%)	10 (7%)	15 48
All	All	1230/1230~(100%)	1122 (91%)	108 (9%)	10 36

All (108) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	8	HIS
1	А	19	LEU
1	А	24	VAL
1	А	32	THR
1	А	65	GLN
1	А	86	VAL
1	А	118	SER
1	А	121	ARG
1	А	122	THR
1	А	123	ASN
1	А	141	LYS
1	А	147	THR
1	А	148	ASP
1	А	179	THR
1	А	194	THR
1	А	202	GLN
1	А	203	SER
1	А	212	THR
1	А	237	SER
1	А	247	ARG
1	А	256	MET
1	А	271	ASP



Mol	Chain	Res	Type
1	А	295	LYS
1	А	298	ARG
1	А	316	GLU
2	В	340	ASP
2	В	379	LYS
2	В	386	LEU
2	В	392	THR
2	В	434	SER
2	В	436	MET
2	В	465	CYS
2	В	488	MET
2	В	489	GLN
2	В	491	ARG
1	С	8	HIS
1	С	19	LEU
1	С	24	VAL
1	С	32	THR
1	С	47	ARG
1	С	65	GLN
1	С	86	VAL
1	С	118	SER
1	С	121	ARG
1	С	122	THR
1	С	123	ASN
1	С	141	LYS
1	С	147	THR
1	С	148	ASP
1	С	179	THR
1	С	182	GLN
1	С	194	THR
1	С	202	GLN
1	C	203	SER
1	С	212	THR
1	C	237	SER
1	С	247	ARG
1	С	256	MET
1	С	271	ASP
1	C	295	LYS
1	С	298	ARG
1	C	316	GLU
2	D	340	ASP
2	D	379	LYS



Mol	Chain	Res	Type
2	D	386	LEU
2	D	392	THR
2	D	434	SER
2	D	436	MET
2	D	465	CYS
2	D	488	MET
2	D	489	GLN
2	D	491	ARG
1	Е	8	HIS
1	Е	19	LEU
1	Е	24	VAL
1	Е	32	THR
1	Е	47	ARG
1	Е	65	GLN
1	Е	86	VAL
1	E	118	SER
1	E	121	ARG
1	Ε	122	THR
1	Ε	123	ASN
1	Ε	141	LYS
1	Ε	147	THR
1	Е	148	ASP
1	Ε	179	THR
1	Е	194	THR
1	Е	202	GLN
1	Е	203	SER
1	Е	212	THR
1	E	237	SER
1	Ε	247	ARG
1	Е	256	MET
1	E	271	ASP
1	Е	295	LYS
1	E	298	ARG
1	Е	316	GLU
2	F	340	ASP
2	F	379	LYS
2	F	386	LEU
2	F	392	THR
2	F	434	SER
2	F	436	MET
2	F	465	CYS
2	F	488	MET



Continued from previous page...

Mol	Chain	Res	Type
2	F	489	GLN
2	F	491	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	217	GLN
1	А	224	HIS
2	В	383	GLN
1	С	224	HIS
1	Е	224	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tink	Bo	ond leng	$_{\rm ths}$	B	ond ang	gles
MOI	туре	Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	NAG	G	1	3	$15,\!15,\!15$	0.48	0	21,21,21	1.32	3 (14%)
3	GAL	G	2	3	11,11,12	0.64	0	15,15,17	0.90	0
3	SIA	G	3	3	20,20,21	0.55	0	21,28,31	0.81	1 (4%)
4	GAL	Н	1	4	12,12,12	0.50	0	17,17,17	0.54	0
4	SIA	Н	2	4	20,20,21	0.56	0	21,28,31	0.81	1 (4%)
4	GAL	Ι	1	4	12,12,12	0.46	0	17,17,17	0.52	0



Mal	Tuno	Chain	Dog	Tink	Bo	ond leng	$_{\rm ths}$	B	ond ang	les
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SIA	Ι	2	4	20,20,21	0.56	0	21,28,31	0.82	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3	-	2/6/26/26	0/1/1/1
3	GAL	G	2	3	-	0/2/19/22	0/1/1/1
3	SIA	G	3	3	-	4/18/34/38	0/1/1/1
4	GAL	Н	1	4	-	2/2/22/22	0/1/1/1
4	SIA	Н	2	4	-	0/18/34/38	0/1/1/1
4	GAL	Ι	1	4	-	2/2/22/22	0/1/1/1
4	SIA	Ι	2	4	-	3/18/34/38	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
3	G	1	NAG	C4-C3-C2	4.02	116.26	110.40
3	G	1	NAG	C1-C2-C3	2.32	113.71	110.54
3	G	3	SIA	01B-C1-C2	2.22	118.50	112.71
4	Ι	2	SIA	O1B-C1-C2	2.22	118.49	112.71
4	Н	2	SIA	O1B-C1-C2	2.20	118.44	112.71
3	G	1	NAG	C3-C2-N2	-2.02	106.90	110.62
4	Ι	2	SIA	O1A-C1-C2	-2.01	118.51	122.85

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Н	1	GAL	O5-C5-C6-O6
3	G	1	NAG	C8-C7-N2-C2
4	Н	1	GAL	C4-C5-C6-O6
3	G	1	NAG	O7-C7-N2-C2
3	G	3	SIA	O7-C7-C8-C9
3	G	3	SIA	C6-C7-C8-O8
4	Ι	1	GAL	O5-C5-C6-O6
3	G	3	SIA	C6-C7-C8-C9



Mol	Chain	Res	Type	Atoms
3	G	3	SIA	O7-C7-C8-O8
4	Ι	2	SIA	C6-C7-C8-O8
4	Ι	1	GAL	C4-C5-C6-O6
4	Ι	2	SIA	O7-C7-C8-C9
4	Ι	2	SIA	O7-C7-C8-O8

There are no ring outliers.

5 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1	NAG	11	0
3	G	2	GAL	3	0
3	G	3	SIA	1	0
4	Н	2	SIA	1	0
4	Ι	2	SIA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Typ	Turne	Chain	Dec	Bog Link Bond lengths				B	ond ang	gles
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	В	601	2	14,14,15	0.49	0	17,19,21	0.93	1 (5%)
5	NAG	D	601	2	14,14,15	0.52	0	17,19,21	0.62	0
5	NAG	F	601	2	14,14,15	0.47	0	17,19,21	0.86	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	В	601	2	-	0/6/23/26	0/1/1/1
5	NAG	D	601	2	-	0/6/23/26	0/1/1/1
5	NAG	F	601	2	-	0/6/23/26	0/1/1/1

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	В	601	NAG	C1-O5-C5	3.22	116.50	112.19
5	F	601	NAG	C1-O5-C5	2.54	115.59	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	316/316~(100%)	-0.05	2 (0%) 89 72	43, 73, 111, 140	0
1	С	316/316~(100%)	0.13	8 (2%) 57 29	46, 91, 145, 232	0
1	Е	316/316~(100%)	0.31	18 (5%) 23 8	59, 101, 169, 236	0
2	В	170/170~(100%)	0.31	13 (7%) 13 4	39, 80, 148, 277	0
2	D	170/170~(100%)	0.31	14 (8%) 11 3	41, 99, 156, 229	0
2	F	170/170~(100%)	1.31	35 (20%) 1 0	55, 158, 221, 306	0
All	All	1458/1458~(100%)	0.31	90 (6%) 20 7	39, 91, 179, 306	0

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	327	ILE	19.2
2	F	326	ALA	17.6
2	F	322	GLY	12.5
2	В	322	GLY	12.0
2	В	323	LEU	11.7
2	F	330	PHE	9.9
2	F	328	ALA	9.7
2	В	324	PHE	9.1
2	F	329	GLY	8.0
2	F	323	LEU	7.7
2	В	328	ALA	7.0
1	Е	5	LEU	6.8
2	F	482	LYS	6.8
2	В	330	PHE	6.4
2	F	343	TYR	6.1
1	E	143	LEU	5.9
2	В	326	ALA	5.9
2	D	322	GLY	5.7
2	F	481	SER	5.7



4LKG

Mol	Chain	Res	Type	RSRZ
2	F	459	PHE	5.6
2	F	325	GLY	5.5
1	Е	310	GLY	5.5
1	С	316	GLU	5.5
1	Е	142	TRP	5.3
2	D	327	ILE	5.2
2	D	324	PHE	5.0
2	D	323	LEU	5.0
2	F	353	GLU	4.9
2	F	485	GLU	4.9
2	F	344	GLY	4.8
2	D	328	ALA	4.8
2	F	324	PHE	4.7
1	С	190	ASN	4.6
2	F	443	VAL	4.6
1	Е	240	GLY	4.5
2	D	353	GLU	4.2
2	F	461	ILE	4.1
2	F	447	LEU	4.0
1	С	164	LYS	3.8
1	Е	3	ILE	3.7
2	F	357	ALA	3.6
2	F	473	ILE	3.6
1	С	200	TYR	3.5
2	В	327	ILE	3.5
2	В	325	GLY	3.4
2	D	379	LYS	3.4
1	Е	316	GLU	3.4
2	D	325	GLY	3.4
1	Е	242	PHE	3.3
2	F	440	TYR	3.3
1	С	236	PHE	3.2
1	Е	6	GLY	3.1
2	F	455	GLY	3.1
2	F	480	HIS	3.1
2	D	329	GLY	3.0
1	Е	7	HIS	3.0
2	В	329	GLY	2.9
1	Е	311	MET	2.9
1	А	2	LYS	2.9
2	F	345	PHE	2.9
1	С	202	GLN	2.8



Mol	Chain	Res	Type	RSRZ
2	F	478	TYR	2.8
2	D	326	ALA	2.8
2	D	354	GLY	2.7
2	F	350	ALA	2.6
1	Е	241	ALA	2.6
1	Е	269	GLU	2.5
2	D	485	GLU	2.5
1	Е	312	LYS	2.5
2	F	358	ASP	2.5
2	D	330	PHE	2.5
1	Е	205	VAL	2.5
1	Е	236	PHE	2.5
2	В	345	PHE	2.4
2	В	376	LEU	2.4
2	F	474	ARG	2.3
1	А	223	PHE	2.3
1	С	315	PRO	2.3
2	F	340	ASP	2.3
1	Е	144	LEU	2.3
2	D	437	ASP	2.2
1	С	117	TYR	2.2
2	F	446	GLN	2.2
2	F	433	ASP	2.1
2	В	378	GLU	2.1
2	В	440	TYR	2.1
2	F	462	PHE	2.1
2	F	453	GLU	2.1
1	Е	48	THR	2.1
2	F	442	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	NAG	G	1	15/15	0.75	0.24	133,154,172,175	0
4	GAL	Ι	1	12/12	0.76	0.25	123,142,151,156	0
4	GAL	Н	1	12/12	0.81	0.20	125,148,178,196	0
4	SIA	Ι	2	20/21	0.90	0.26	130,136,145,147	0
3	SIA	G	3	20/21	0.91	0.19	90,106,115,115	0
4	SIA	Н	2	20/21	0.91	0.15	92,102,113,115	0
3	GAL	G	2	11/12	0.93	0.21	84,89,95,106	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.











6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
5	NAG	В	601	14/15	0.86	0.22	70,84,88,89	0
5	NAG	F	601	14/15	0.88	0.19	96,120,140,140	0
5	NAG	D	601	14/15	0.92	0.13	83,90,107,114	0

6.5 Other polymers (i)

There are no such residues in this entry.

