



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 2, 2024 – 08:23 pm GMT

PDB ID : 5LLB
Title : Structure of Polyphosphate Kinase 2 from *Francisella tularensis* with AMP-PCH2PPP and polyphosphate
Authors : Roach, P.L.; Parnell, A.E.
Deposited on : 2016-07-27
Resolution : 1.92 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	 4% 86% 5% 11%
1	B	269	 4% 81% 8% 11%
2	C	263	 2% 86% 5% 9%
2	D	263	 3% 86% 5% 9%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 16918 atoms, of which 7943 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyphosphate kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	240	3969	1294	1959	351	358	7	0	0	0
1	B	240	3960	1293	1951	351	358	7	0	0	0

- Molecule 2 is a protein called Polyphosphate kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	C	240	3998	1294	1988	351	358	7	0	0	0
2	D	240	3999	1294	1989	351	358	7	0	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0

- Molecule 5 is [[(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl]oxy-[[oxidanyl-[oxidanyl(phosphonooxy)phosphoryl]oxy-phosphoryl]methyl]phosphinic acid (three-letter code: 6YZ) (formula: C₁₁H₂₀N₅O₁₈P₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C H N O P 53 11 14 5 18 5	0	0
5	B	1	Total C H N O P 53 11 14 5 18 5	0	0
5	C	1	Total C H N O P 53 11 14 5 18 5	0	0
5	D	1	Total C H N O P 53 11 14 5 18 5	0	0

- Molecule 6 is [oxidanyl-[oxidanyl-[oxidanyl(phosphonooxy)phosphoryl]oxy-phosphoryl]oxy-phosphoryl] phosphono hydrogen phosphate (three-letter code: 6YW) (formula: H₈O₁₉P₆).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	O P	0	0
			25	19 6		
6	B	1	Total	O P	0	0
			25	19 6		
6	C	1	Total	O P	0	0
			25	19 6		
6	D	1	Total	O P	0	0
			25	19 6		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	164	Total	O	0	0
			164	164		
7	B	163	Total	O	0	0
			163	163		
7	C	201	Total	O	0	0
			201	201		
7	D	144	Total	O	0	0
			144	144		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Polyphosphate kinase 2




- Molecule 1: Polyphosphate kinase 2



- Molecule 2: Polyphosphate kinase 2



- Molecule 2: Polyphosphate kinase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.70Å 144.91Å 70.59Å 90.00° 113.04° 90.00°	Depositor
Resolution (Å)	72.46 – 1.92 43.31 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.2 (72.46-1.92) 93.3 (43.31-1.92)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.92Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.179 , 0.231 0.180 , 0.232	Depositor DCC
R_{free} test set	3971 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	25.8	Xtrriage
Anisotropy	0.704	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 78.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.046 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16918	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 6YW, CL, 6YZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/2062	0.63	0/2781
1	B	0.56	0/2061	0.64	0/2780
2	C	0.57	0/2062	0.64	0/2781
2	D	0.54	0/2062	0.63	0/2781
All	All	0.56	0/8247	0.64	0/11123

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2010	1959	2000	9	0
1	B	2009	1951	1998	12	0
2	C	2010	1988	2000	7	0
2	D	2010	1989	2000	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	39	14	0	1	0
5	B	39	14	0	0	0
5	C	39	14	0	1	0
5	D	39	14	0	0	0
6	A	25	0	0	2	0
6	B	25	0	0	1	0
6	C	25	0	0	0	0
6	D	25	0	0	1	0
7	A	164	0	0	3	0
7	B	163	0	0	1	1
7	C	201	0	0	5	1
7	D	144	0	0	0	0
All	All	8975	7943	7998	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:ARG:NH2	6:B:304:6YW:O15	1.97	0.97
2:D:228:LYS:NZ	6:D:304:6YW:O18	2.14	0.79
2:C:31:LYS:NZ	7:C:401:HOH:O	2.18	0.76
1:B:172:LYS:NZ	7:B:401:HOH:O	2.21	0.73
5:C:303:6YZ:N31	7:C:403:HOH:O	2.26	0.68
5:A:303:6YZ:P02	7:A:401:HOH:O	2.59	0.60
1:A:228:LYS:NZ	6:A:304:6YW:O25	2.32	0.56
2:D:43:PHE:HE1	2:D:243:VAL:HG21	1.71	0.55
1:A:48:LYS:NZ	1:A:80:ARG:O	2.40	0.54
2:C:150:LYS:NZ	7:C:408:HOH:O	2.41	0.53
1:B:185:GLN:O	1:B:188:LEU:HG	2.09	0.53
1:B:26:ARG:NH2	1:B:30:GLU:OE2	2.42	0.52
1:B:117:ASP:O	1:B:118:ARG:HB2	2.12	0.49
1:B:225:SER:HA	1:B:231:ALA:HB2	1.94	0.49
1:A:83:LYS:NZ	7:A:402:HOH:O	2.45	0.49
2:C:117:ASP:O	2:C:118:ARG:HB2	2.14	0.47
1:A:25:PRO:HG2	1:A:28:VAL:HG12	1.96	0.47
1:B:163:TRP:HB3	1:B:222:ILE:HD13	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:LYS:HG2	7:A:487:HOH:O	2.16	0.46
1:B:68:GLY:O	1:B:72:ARG:HD3	2.17	0.45
1:B:175:PHE:HD2	1:B:196:LEU:HD21	1.81	0.44
1:A:184:LYS:NZ	6:A:304:6YW:O09	2.37	0.44
2:D:43:PHE:CE1	2:D:243:VAL:HG21	2.51	0.44
2:C:43:PHE:CE1	2:C:243:VAL:HG11	2.53	0.44
2:C:191:ILE:CD1	7:C:559:HOH:O	2.65	0.44
1:A:25:PRO:O	1:A:28:VAL:HG12	2.18	0.43
1:B:93:GLN:O	1:B:97:GLN:HG3	2.18	0.43
2:D:117:ASP:O	2:D:118:ARG:HB2	2.18	0.43
2:C:52:LYS:HE3	2:C:158:MET:HB2	2.00	0.42
1:A:24:ILE:CG1	1:A:25:PRO:HD2	2.50	0.42
1:B:24:ILE:CG1	1:B:25:PRO:HD2	2.50	0.41
2:C:232:ARG:NE	7:C:402:HOH:O	2.25	0.41
1:A:117:ASP:O	1:A:118:ARG:HB2	2.21	0.41
2:D:93:GLN:O	2:D:97:GLN:HG3	2.21	0.41
2:D:82:ALA:O	2:D:83:LYS:HD3	2.21	0.40
1:B:190:PRO:O	1:B:193:LYS:HB2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:448:HOH:O	7:C:457:HOH:O[2_655]	1.98	0.22

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/269 (88%)	233 (98%)	5 (2%)	0	100	100
1	B	238/269 (88%)	234 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	238/263 (90%)	234 (98%)	4 (2%)	0	100	100
2	D	238/263 (90%)	230 (97%)	8 (3%)	0	100	100
All	All	952/1064 (90%)	931 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/244 (88%)	214 (100%)	0	100	100
1	B	214/244 (88%)	210 (98%)	4 (2%)	57	51
2	C	214/238 (90%)	210 (98%)	4 (2%)	57	51
2	D	214/238 (90%)	209 (98%)	5 (2%)	50	43
All	All	856/964 (89%)	843 (98%)	13 (2%)	65	61

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26	ARG
1	B	188	LEU
1	B	191	ILE
1	B	195	SER
2	C	79	PRO
2	C	135	GLU
2	C	166	VAL
2	C	191	ILE
2	D	95	ARG
2	D	188	LEU
2	D	197	ASP
2	D	246	ASP
2	D	247	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	76	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	6YW	C	304	-	14,24,24	0.55	0	21,41,41	1.80	5 (23%)
6	6YW	D	304	-	14,24,24	0.66	0	21,41,41	1.89	5 (23%)
5	6YZ	A	303	4	32,41,41	4.23	10 (31%)	38,66,66	1.38	4 (10%)
6	6YW	A	304	-	14,24,24	0.64	0	21,41,41	2.29	4 (19%)
5	6YZ	C	303	4	32,41,41	3.78	11 (34%)	38,66,66	1.49	8 (21%)
6	6YW	B	304	-	14,24,24	0.67	0	21,41,41	1.83	5 (23%)
5	6YZ	D	303	4	32,41,41	4.11	11 (34%)	38,66,66	1.25	3 (7%)
5	6YZ	B	303	4	32,41,41	4.07	10 (31%)	38,66,66	1.28	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	6YW	C	304	-	-	4/30/30/30	-
6	6YW	D	304	-	-	4/30/30/30	-
5	6YZ	A	303	4	-	2/24/50/50	0/3/3/3
6	6YW	A	304	-	-	1/30/30/30	-
5	6YZ	C	303	4	-	2/24/50/50	0/3/3/3
6	6YW	B	304	-	-	2/30/30/30	-
5	6YZ	D	303	4	-	0/24/50/50	0/3/3/3
5	6YZ	B	303	4	-	4/24/50/50	0/3/3/3

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	303	6YZ	O20-C21	14.62	1.61	1.41
5	D	303	6YZ	O20-C21	14.27	1.61	1.41
5	B	303	6YZ	C32-C21	-14.21	1.32	1.53
5	D	303	6YZ	C32-C21	-14.11	1.32	1.53
5	A	303	6YZ	C32-C21	-13.68	1.33	1.53
5	B	303	6YZ	O20-C21	13.63	1.60	1.41
5	C	303	6YZ	C32-C21	-13.08	1.33	1.53
5	C	303	6YZ	O20-C21	12.52	1.58	1.41
5	D	303	6YZ	O20-C19	-7.50	1.28	1.45
5	A	303	6YZ	O20-C19	-7.42	1.28	1.45
5	B	303	6YZ	O20-C19	-7.22	1.28	1.45
5	C	303	6YZ	O20-C19	-6.93	1.29	1.45
5	A	303	6YZ	P09-O08	6.25	1.65	1.58
5	B	303	6YZ	P09-O08	4.00	1.62	1.58
5	A	303	6YZ	C26-N27	3.87	1.41	1.35
5	C	303	6YZ	P09-O08	3.81	1.62	1.58
5	D	303	6YZ	P09-O08	3.54	1.62	1.58
5	B	303	6YZ	C26-N27	3.54	1.40	1.35
5	B	303	6YZ	P12-O14	3.52	1.62	1.58
5	C	303	6YZ	C26-N27	3.45	1.40	1.35
5	A	303	6YZ	O35-C34	-3.42	1.34	1.43
5	D	303	6YZ	C26-N27	3.37	1.40	1.35
5	D	303	6YZ	C30-N31	3.29	1.46	1.34
5	A	303	6YZ	C30-N31	3.20	1.45	1.34
5	B	303	6YZ	C30-N31	3.15	1.45	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	303	6YZ	O33-C32	3.09	1.50	1.43
5	C	303	6YZ	O33-C32	3.05	1.50	1.43
5	D	303	6YZ	P12-O14	3.05	1.61	1.58
5	C	303	6YZ	C30-N31	3.02	1.45	1.34
5	B	303	6YZ	O35-C34	-2.99	1.35	1.43
5	C	303	6YZ	P12-O14	2.91	1.61	1.58
5	A	303	6YZ	O33-C32	2.85	1.49	1.43
5	A	303	6YZ	P12-O14	2.78	1.61	1.58
5	D	303	6YZ	O35-C34	-2.70	1.36	1.43
5	D	303	6YZ	O33-C32	2.62	1.49	1.43
5	C	303	6YZ	C28-N27	2.47	1.36	1.32
5	C	303	6YZ	O35-C34	-2.31	1.37	1.43
5	C	303	6YZ	P12-O13	-2.31	1.51	1.56
5	D	303	6YZ	C28-N27	2.16	1.35	1.32
5	A	303	6YZ	P12-O13	-2.15	1.51	1.56
5	D	303	6YZ	P12-O13	-2.13	1.51	1.56
5	B	303	6YZ	P12-O13	-2.08	1.51	1.56

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	304	6YW	P11-O10-P08	-6.41	110.82	132.83
6	A	304	6YW	P05-O04-P02	-6.37	110.97	132.83
6	D	304	6YW	P14-O16-P17	-5.66	113.41	132.83
6	B	304	6YW	P11-O10-P08	-4.79	116.38	132.83
6	B	304	6YW	P05-O04-P02	-4.42	117.66	132.83
5	C	303	6YZ	N27-C28-N29	-4.00	122.43	128.68
5	A	303	6YZ	O10-P09-C11	-3.90	98.76	109.07
5	D	303	6YZ	N27-C28-N29	-3.89	122.60	128.68
6	C	304	6YW	P08-O07-P05	-3.88	119.53	132.83
5	C	303	6YZ	C34-C32-C21	3.69	106.54	100.98
5	A	303	6YZ	N27-C28-N29	-3.56	123.12	128.68
5	B	303	6YZ	N27-C28-N29	-3.48	123.23	128.68
6	C	304	6YW	P05-O04-P02	-3.47	120.92	132.83
6	C	304	6YW	P11-O10-P08	-3.41	121.12	132.83
5	A	303	6YZ	O07-P04-O03	3.32	115.78	104.64
6	D	304	6YW	P11-O10-P08	-3.25	121.67	132.83
5	C	303	6YZ	O07-P04-O03	3.19	115.32	104.64
6	A	304	6YW	P14-O13-P11	-3.15	122.01	132.83
6	D	304	6YW	P14-O13-P11	-3.13	122.08	132.83
5	D	303	6YZ	C26-C25-N24	-3.11	106.16	109.40
6	D	304	6YW	P05-O04-P02	-2.86	123.01	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	303	6YZ	O10-P09-C11	-2.76	101.78	109.07
6	B	304	6YW	P14-O13-P11	-2.74	123.41	132.83
5	D	303	6YZ	C34-C32-C21	2.67	105.00	100.98
6	C	304	6YW	P14-O13-P11	-2.63	123.81	132.83
5	B	303	6YZ	O10-P09-C11	-2.48	102.51	109.07
6	C	304	6YW	O18-P17-O16	2.46	112.88	104.64
5	B	303	6YZ	C18-C19-C34	-2.44	106.03	115.18
5	B	303	6YZ	O06-P04-O03	2.37	112.57	104.64
6	B	304	6YW	P14-O16-P17	-2.36	124.72	132.83
5	C	303	6YZ	O38-P09-C11	2.33	116.10	106.58
5	C	303	6YZ	O37-P12-C11	2.26	115.04	109.07
6	A	304	6YW	O19-P17-O16	2.23	112.12	104.64
5	C	303	6YZ	O20-C21-C32	-2.23	103.66	106.93
5	C	303	6YZ	C18-C19-C34	-2.23	106.84	115.18
6	D	304	6YW	P08-O07-P05	-2.17	125.39	132.83
6	B	304	6YW	O03-P02-O04	2.14	111.80	104.64
5	A	303	6YZ	C18-C19-C34	-2.10	107.32	115.18

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	303	6YZ	P12-C11-P09-O08
5	B	303	6YZ	P12-C11-P09-O10
5	B	303	6YZ	P12-C11-P09-O38
5	B	303	6YZ	P02-O03-P04-O06
5	A	303	6YZ	P09-O08-P02-O39
6	C	304	6YW	P05-O07-P08-O09
6	C	304	6YW	P17-O16-P14-O15
6	D	304	6YW	P05-O07-P08-O09
5	C	303	6YZ	P12-C11-P09-O38
6	B	304	6YW	P11-O10-P08-O07
6	C	304	6YW	P11-O10-P08-O09
5	A	303	6YZ	P12-C11-P09-O10
5	C	303	6YZ	P12-C11-P09-O10
6	D	304	6YW	P11-O13-P14-O21
6	A	304	6YW	P11-O10-P08-O23
6	B	304	6YW	P11-O10-P08-O09
6	C	304	6YW	P11-O10-P08-O23
6	D	304	6YW	P08-O10-P11-O22
6	D	304	6YW	P11-O13-P14-O15


There are no ring outliers.




5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	304	6YW	1	0
5	A	303	6YZ	1	0
6	A	304	6YW	2	0
5	C	303	6YZ	1	0
6	B	304	6YW	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/269 (89%)	0.32	12 (5%) 28 32	37, 49, 73, 89	0
1	B	240/269 (89%)	0.30	11 (4%) 32 35	38, 50, 72, 123	0
2	C	240/263 (91%)	0.09	4 (1%) 70 72	37, 48, 66, 82	0
2	D	240/263 (91%)	0.41	9 (3%) 40 43	38, 53, 77, 90	0
All	All	960/1064 (90%)	0.28	36 (3%) 40 43	37, 49, 73, 123	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	LEU	5.5
1	B	191	ILE	5.0
1	B	188	LEU	4.6
2	D	24	ILE	4.2
2	D	166	VAL	4.0
1	A	28	VAL	3.5
1	A	190	PRO	3.5
1	A	191	ILE	3.3
2	D	191	ILE	3.3
1	B	25	PRO	3.3
1	B	193	LYS	3.2
1	A	196	LEU	3.1
2	D	188	LEU	2.6
1	A	24	ILE	2.6
1	A	192	ASP	2.5
1	B	28	VAL	2.4
2	C	27	ASN	2.4
1	A	251	GLU	2.4
1	A	27	ASN	2.4
1	B	196	LEU	2.3
1	A	132	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	175	PHE	2.3
1	A	263	THR	2.2
1	B	192	ASP	2.2
2	D	246	ASP	2.2
2	C	191	ILE	2.2
1	B	186	TRP	2.2
1	B	139	PHE	2.1
1	B	24	ILE	2.1
2	D	263	THR	2.1
2	C	244	ASP	2.1
1	A	181	HIS	2.1
1	B	32	GLN	2.1
2	D	210	PHE	2.0
2	C	24	ILE	2.0
2	D	211	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
6	6YW	A	304	25/25	0.67	0.17	75,106,151,161	0
6	6YW	D	304	25/25	0.68	0.18	83,115,139,147	0
6	6YW	C	304	25/25	0.73	0.17	69,92,118,150	0
6	6YW	B	304	25/25	0.73	0.17	68,96,112,122	0
5	6YZ	B	303	39/39	0.89	0.15	49,67,106,123	0
3	CL	A	301	1/1	0.91	0.13	57,57,57,57	0
4	MG	A	302	1/1	0.92	0.26	62,62,62,62	0
5	6YZ	D	303	39/39	0.92	0.12	53,63,75,79	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	6YZ	A	303	39/39	0.92	0.12	50,64,79,88	0
4	MG	D	302	1/1	0.93	0.10	58,58,58,58	0
3	CL	B	301	1/1	0.94	0.10	62,62,62,62	0
4	MG	B	302	1/1	0.94	0.18	73,73,73,73	0
5	6YZ	C	303	39/39	0.95	0.11	40,52,62,64	0
3	CL	D	301	1/1	0.96	0.08	72,72,72,72	0
3	CL	C	301	1/1	0.97	0.15	50,50,50,50	0
4	MG	C	302	1/1	0.98	0.15	45,45,45,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 6YW D 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 6YW C 304:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 6YW B 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 6YZ B 303:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 6YZ D 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 6YZ A 303:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.