



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2023 – 06:33 AM EDT

PDB ID : 7LMJ
Title : SARS-CoV-1 3CLPro in complex with 2-(1H-benzo[d][1,2,3]triazol-1-yl)-N-(3-chlorobenzyl)-N-(4-(2-oxo-1,2-dihydropyridin-3-yl)phenyl)acetamide
Authors : Arya, T.; Goins, C.M.; Macdonald, J.D.; Stauffer, S.R.
Deposited on : 2021-02-05
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	306	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	AAA	406	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2696 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	306	2390	1509	408	449	24	0	3	0

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	AAA	1	7	4	3	0	0
2	AAA	1	7	4	3	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total C O 6 3 3	0	0
3	AAA	1	Total C O 6 3 3	0	0
3	AAA	1	Total C O 6 3 3	0	0

- Molecule 4 is 2-(benzotriazol-1-yl)- {N}-[(3-chlorophenyl)methyl]- {N}-[4-(2-oxidanylidene-1 {H}-pyridin-3-yl)phenyl]ethanamide (three-letter code: Y67) (formula: C₂₆H₂₀ClN₅O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
4	AAA	1	34	26	1	5	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	240	Total	O	0	0
			240	240		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	107.54Å 82.71Å 53.20Å 90.00° 105.30° 90.00°	Depositor
Resolution (Å)	32.36 – 1.69 32.33 – 1.69	Depositor EDS
% Data completeness (in resolution range)	98.8 (32.36-1.69) 98.9 (32.33-1.69)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.97 (at 1.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.146 , 0.192 0.157 , 0.197	Depositor DCC
R_{free} test set	2506 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtrriage
Anisotropy	0.238	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2696	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: Y67, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.92	3/2443 (0.1%)	1.00	4/3318 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	290	GLU	CD-OE2	14.86	1.42	1.25
1	AAA	215	GLY	C-O	5.88	1.33	1.23
1	AAA	11	GLY	C-O	5.43	1.32	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	4	ARG	NE-CZ-NH2	-7.22	116.69	120.30
1	AAA	4	ARG	NE-CZ-NH1	6.52	123.56	120.30
1	AAA	153	ASP	CB-CA-C	5.58	121.56	110.40
1	AAA	222	ARG	NE-CZ-NH1	5.41	123.00	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2390	0	2336	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	AAA	14	0	20	7	0
3	AAA	18	0	24	1	0
4	AAA	34	0	0	1	0
5	AAA	240	0	0	5	0
All	All	2696	0	2380	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:AAA:401:PEG:H11	5:AAA:531:HOH:O	1.86	0.75
2:AAA:401:PEG:O1	5:AAA:501:HOH:O	2.06	0.72
1:AAA:188:ARG:HD3	5:AAA:614:HOH:O	1.99	0.62
1:AAA:304:THR:HG21	5:AAA:680:HOH:O	2.03	0.58
1:AAA:59:ILE:HD13	1:AAA:59:ILE:O	2.05	0.57
1:AAA:304:THR:CG2	5:AAA:680:HOH:O	2.54	0.54
1:AAA:297:VAL:HG21	2:AAA:406:PEG:H32	1.91	0.52
4:AAA:404:Y67:C14	3:AAA:405:GOL:H2	2.39	0.51
1:AAA:87:LEU:HD21	1:AAA:89:LEU:HD21	1.96	0.47
1:AAA:252:PRO:HB3	2:AAA:406:PEG:C1	2.44	0.47
1:AAA:222:ARG:O	1:AAA:222:ARG:HG2	2.15	0.47
1:AAA:297:VAL:HG11	2:AAA:406:PEG:H22	1.96	0.47
1:AAA:276:MET:HE2	1:AAA:279:ARG:O	2.15	0.46
1:AAA:115:LEU:HD11	1:AAA:122:PRO:HB3	1.97	0.46
1:AAA:66:PHE:CE1	1:AAA:87:LEU:HD21	2.52	0.45
1:AAA:252:PRO:HB3	2:AAA:406:PEG:H12	1.99	0.44
1:AAA:142:ASN:HD22	1:AAA:142:ASN:HA	1.58	0.44
1:AAA:113:SER:O	1:AAA:149:GLY:HA2	2.19	0.43
1:AAA:252:PRO:HG2	2:AAA:406:PEG:H41	2.01	0.42
1:AAA:69:GLN:HE21	1:AAA:69:GLN:HB2	1.60	0.42
1:AAA:114:VAL:O	1:AAA:125:VAL:HA	2.19	0.41
1:AAA:86:LEU:HG	1:AAA:179:GLY:CA	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	307/306 (100%)	302 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	266/263 (101%)	256 (96%)	10 (4%)	33	12

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	27	LEU
1	AAA	59	ILE
1	AAA	142	ASN
1	AAA	153	ASP
1	AAA	180	LYS
1	AAA	198[A]	THR
1	AAA	198[B]	THR
1	AAA	222	ARG
1	AAA	235	MET
1	AAA	279	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	AAA	403	-	5,5,5	0.39	0	5,5,5	0.64	0
2	PEG	AAA	406	-	6,6,6	1.17	0	5,5,5	0.74	0
4	Y67	AAA	404	-	35,38,38	2.46	8 (22%)	45,53,53	2.25	13 (28%)
3	GOL	AAA	402	-	5,5,5	0.32	0	5,5,5	0.63	0
2	PEG	AAA	401	-	6,6,6	0.63	0	5,5,5	0.66	0
3	GOL	AAA	405	-	5,5,5	0.11	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	AAA	403	-	-	0/4/4/4	-
2	PEG	AAA	406	-	-	4/4/4/4	-
4	Y67	AAA	404	-	-	0/20/20/20	0/5/5/5
3	GOL	AAA	402	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	AAA	401	-	-	3/4/4/4	-
3	GOL	AAA	405	-	-	3/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	AAA	404	Y67	C29-C28	9.36	1.51	1.35
4	AAA	404	Y67	C29-N30	6.71	1.46	1.36
4	AAA	404	Y67	C31-N30	3.62	1.46	1.39
4	AAA	404	Y67	C27-C26	3.61	1.45	1.38
4	AAA	404	Y67	C02-N13	3.50	1.41	1.36
4	AAA	404	Y67	C26-C31	2.78	1.50	1.46
4	AAA	404	Y67	C19-CL20	2.57	1.80	1.74
4	AAA	404	Y67	C22-N13	2.04	1.47	1.43

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	AAA	404	Y67	C29-N30-C31	-8.37	118.83	124.47
4	AAA	404	Y67	C02-C03-N04	-5.05	104.75	110.76
4	AAA	404	Y67	C25-C26-C27	-4.19	115.25	121.36
4	AAA	404	Y67	O01-C02-N13	-3.96	115.91	121.66
4	AAA	404	Y67	C33-C25-C26	-3.05	116.68	120.98
4	AAA	404	Y67	C34-C33-C25	-2.99	117.31	120.78
4	AAA	404	Y67	C03-N04-N12	2.93	124.26	117.11
4	AAA	404	Y67	C23-C22-N13	-2.87	115.98	120.18
4	AAA	404	Y67	C23-C24-C25	-2.86	117.45	120.78
4	AAA	404	Y67	C33-C25-C24	2.61	122.31	118.59
4	AAA	404	Y67	C14-C15-C16	-2.45	116.15	120.77
4	AAA	404	Y67	C28-C29-N30	2.39	122.95	120.19
4	AAA	404	Y67	C08-C09-C10	-2.04	117.14	120.08

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	AAA	402	GOL	O1-C1-C2-C3
3	AAA	405	GOL	C1-C2-C3-O3
2	AAA	401	PEG	O1-C1-C2-O2
2	AAA	401	PEG	O2-C3-C4-O4
2	AAA	406	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	AAA	405	GOL	O2-C2-C3-O3
3	AAA	402	GOL	O1-C1-C2-O2
2	AAA	406	PEG	O1-C1-C2-O2
2	AAA	406	PEG	C4-C3-O2-C2
2	AAA	401	PEG	C4-C3-O2-C2
2	AAA	406	PEG	C1-C2-O2-C3
3	AAA	405	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	406	PEG	5	0
4	AAA	404	Y67	1	0
2	AAA	401	PEG	2	0
3	AAA	405	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	306/306 (100%)	-0.04	12 (3%) 39 42	26, 38, 60, 88	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	154	TYR	7.7
1	AAA	277	ASN	4.8
1	AAA	72	ASN	4.6
1	AAA	191	ALA	3.4
1	AAA	303	VAL	3.0
1	AAA	302	GLY	2.9
1	AAA	153	ASP	2.8
1	AAA	255	ALA	2.7
1	AAA	115	LEU	2.5
1	AAA	278	GLY	2.4
1	AAA	47	GLU	2.1
1	AAA	125	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	AAA	406	7/7	0.83	0.44	41,47,63,64	0
3	GOL	AAA	405	6/6	0.85	0.19	66,68,80,82	0
2	PEG	AAA	401	7/7	0.92	0.25	46,57,70,74	0
3	GOL	AAA	403	6/6	0.93	0.09	41,59,67,75	0
3	GOL	AAA	402	6/6	0.97	0.11	37,44,57,61	0
4	Y67	AAA	404	34/34	0.97	0.07	28,37,50,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.