



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 03:41 pm BST

PDB ID : 1LO7
Title : X-ray structure of 4-Hydroxybenzoyl CoA Thioesterase complexed with 4-hydroxyphenacyl CoA
Authors : Thoden, J.B.; Holden, H.M.; Zhuang, Z.; Dunaway-Mariano, D.
Deposited on : 2002-05-06
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1375 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 4-hydroxybenzoyl-CoA Thioesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	140	1136	715	202	209	10	0	5	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	TYR	PHE	CONFLICT	UNP P56653

- Molecule 2 is 4-HYDROXYPHENACYL COENZYME A (three-letter code: 4CO) (formula: $C_{29}H_{42}N_7O_{18}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
2	A	1	58	29	7	18	3	1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	173	Total O 173 173	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-hydroxybenzoyl-CoA Thioesterase

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	51.50Å 55.20Å 93.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.50 18.83 – 1.50	Depositor EDS
% Data completeness (in resolution range)	94.1 (30.00-1.50) 96.2 (18.83-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 1.50Å)	Xtrriage
Refinement program	TNT 5F	Depositor
R, R_{free}	0.161 , 0.198 0.162 , 0.153	Depositor DCC
R_{free} test set	2076 reflections (9.85%)	wwPDB-VP
Wilson B-factor (Å ²)	48.8	Xtrriage
Anisotropy	0.657	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 74.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1375	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 4CO, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.97	8/1182 (0.7%)	1.37	22/1600 (1.4%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	53	GLU	CD-OE2	6.23	1.32	1.25
1	A	125	GLU	CD-OE2	6.16	1.32	1.25
1	A	12	GLU	CD-OE2	6.13	1.32	1.25
1	A	81	GLU	CD-OE2	5.89	1.32	1.25
1	A	138	GLU	CD-OE2	5.60	1.31	1.25
1	A	115	GLU	CD-OE2	5.50	1.31	1.25
1	A	131	GLU	CD-OE2	5.37	1.31	1.25
1	A	86	GLU	CD-OE2	5.12	1.31	1.25

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	17	ASP	CB-CG-OD2	-8.24	110.88	118.30
1	A	106	ASP	CB-CG-OD2	-7.21	111.81	118.30
1	A	17	ASP	CB-CG-OD1	6.77	124.39	118.30
1	A	117	ARG	NE-CZ-NH2	-6.64	116.98	120.30
1	A	3	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	A	106	ASP	CB-CG-OD1	6.09	123.78	118.30
1	A	126	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	A	75	ASP	CB-CG-OD2	-5.92	112.97	118.30
1	A	36	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	A	126	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	A	100	ARG	NE-CZ-NH1	5.85	123.23	120.30
1	A	75	ASP	CB-CG-OD1	5.75	123.48	118.30
1	A	121[A]	MET	CA-CB-CG	-5.69	103.63	113.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121[B]	MET	CA-CB-CG	-5.69	103.63	113.30
1	A	10	ARG	NE-CZ-NH1	5.67	123.13	120.30
1	A	117	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	A	112	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	A	114	ASP	CB-CG-OD1	5.26	123.04	118.30
1	A	111	MET	CG-SD-CE	5.24	108.58	100.20
1	A	54	ARG	NE-CZ-NH1	5.20	122.90	120.30
1	A	48	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	A	123	ASP	CB-CG-OD1	5.11	122.90	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1136	0	1112	23	0
2	A	58	0	38	1	0
3	A	8	0	12	0	0
4	A	173	0	0	6	4
All	All	1375	0	1162	24	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:THR:HG21	1:A:106:ASP:HB2	1.63	0.81
1:A:102:THR:HG22	1:A:106:ASP:H	1.50	0.74
1:A:102:THR:HG23	1:A:104:GLY:H	1.57	0.70
1:A:101:THR:HG22	1:A:102:THR:O	1.93	0.69
1:A:100:ARG:HG2	1:A:100:ARG:HH11	1.57	0.67
1:A:55:GLY:O	1:A:121[A]:MET:HG2	1.94	0.67
1:A:102:THR:CG2	1:A:106:ASP:HB2	2.27	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:ARG:HH11	1:A:100:ARG:CG	2.16	0.59
1:A:100:ARG:HG2	1:A:100:ARG:NH1	2.15	0.59
1:A:126:ARG:HD3	4:A:371:HOH:O	2.02	0.58
1:A:48:ARG:HD2	4:A:322:HOH:O	2.03	0.58
1:A:17:ASP:HB2	1:A:18:PRO:CD	2.34	0.57
1:A:17:ASP:HB2	1:A:18:PRO:HD2	1.86	0.57
1:A:102:THR:HG22	1:A:106:ASP:N	2.19	0.56
1:A:128:ARG:HB2	4:A:348:HOH:O	2.04	0.55
1:A:85:LYS:NZ	4:A:331:HOH:O	2.38	0.55
1:A:102:THR:OG1	1:A:108:GLN:NE2	2.42	0.52
2:A:170:4CO:O9P	2:A:170:4CO:H131	2.12	0.50
1:A:102:THR:HG23	1:A:104:GLY:N	2.25	0.48
1:A:100:ARG:NH2	4:A:241:HOH:O	2.50	0.45
1:A:11:ILE:HD13	1:A:22:VAL:CG2	2.47	0.45
1:A:59:THR:HA	1:A:118:VAL:O	2.18	0.44
1:A:85:LYS:HD3	4:A:331:HOH:O	2.19	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:352:HOH:O	4:A:352:HOH:O[2_665]	0.93	1.27
4:A:363:HOH:O	4:A:363:HOH:O[2_665]	1.16	1.04
4:A:353:HOH:O	4:A:353:HOH:O[2_655]	1.34	0.86
4:A:362:HOH:O	4:A:362:HOH:O[3_655]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	142/141 (101%)	138 (97%)	4 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	127/123 (103%)	122 (96%)	5 (4%)	32 7

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31[A]	LEU
1	A	31[B]	LEU
1	A	100	ARG
1	A	102	THR
1	A	128	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	GLN
1	A	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	401	-	3,3,3	0.27	0	2,2,2	0.27	0
3	EDO	A	400	-	3,3,3	0.41	0	2,2,2	0.11	0
2	4CO	A	170	-	53,61,61	1.50	6 (11%)	68,90,90	1.57	11 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	401	-	-	1/1/1/1	-
3	EDO	A	400	-	-	1/1/1/1	-
2	4CO	A	170	-	-	6/52/72/72	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	170	4CO	C9P-N8P	5.43	1.45	1.33
2	A	170	4CO	C5P-N4P	5.02	1.44	1.33
2	A	170	4CO	CB-C1B	-3.30	1.48	1.51
2	A	170	4CO	O4D-C1D	-2.50	1.37	1.41
2	A	170	4CO	P1A-O2A	-2.21	1.45	1.55
2	A	170	4CO	P3D-O8A	-2.02	1.47	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	170	4CO	C7P-N8P-C9P	-5.71	112.40	122.59
2	A	170	4CO	CDP-CBP-CAP	-4.87	100.38	108.82
2	A	170	4CO	C3P-N4P-C5P	-3.98	115.45	122.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	170	4CO	O4D-C1D-C2D	-3.35	102.03	106.93
2	A	170	4CO	O6A-CCP-CBP	-3.04	105.67	110.55
2	A	170	4CO	C2D-C3D-C4D	-2.40	98.98	103.22
2	A	170	4CO	C7B-C2B-C3B	2.31	121.88	118.59
2	A	170	4CO	C7P-C6P-C5P	-2.18	108.72	112.36
2	A	170	4CO	C6P-C5P-N4P	-2.13	112.84	116.42
2	A	170	4CO	C2P-S1P-CB	2.09	105.20	101.71
2	A	170	4CO	C6B-C7B-C2B	-2.00	118.45	120.78

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	170	4CO	S1P-C2P-C3P-N4P
2	A	170	4CO	O1B-C1B-CB-S1P
2	A	170	4CO	C2B-C1B-CB-S1P
2	A	170	4CO	P1A-O3A-P2A-O6A
3	A	401	EDO	O1-C1-C2-O2
3	A	400	EDO	O1-C1-C2-O2
2	A	170	4CO	C1B-CB-S1P-C2P
2	A	170	4CO	C3D-O3D-P3D-O9A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	170	4CO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	140/141 (99%)	-0.57	3 (2%) 63 68	9, 14, 46, 75	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	104	GLY	5.4
1	A	103	PRO	3.6
1	A	102	THR	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	400	4/4	0.90	0.11	23,33,36,43	0
3	EDO	A	401	4/4	0.91	0.12	31,33,49,55	0
2	4CO	A	170	58/58	0.98	0.05	7,14,20,35	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.