



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 3, 2024 – 04:15 PM EST

PDB ID : 1LPE  
Title : THREE-DIMENSIONAL STRUCTURE OF THE LDL RECEPTOR-BINDING DOMAIN OF HUMAN APOLIPOPROTEIN E  
Authors : Wilson, C.; Agard, D.A.  
Deposited on : 1991-08-22  
Resolution : 2.25 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	144	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1293 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APOLIPOPROTEIN E3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	144	1172	726	218	223	5	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	121	Total	O	0	0
			121	121		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: APOLIPOPROTEIN E3

Chain A:  68% 27% ..



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.65Å 53.96Å 85.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.25	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.25)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.175 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1293	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.90	0/1184	1.95	38/1589 (2.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	ARG	NE-CZ-NH2	-12.48	114.06	120.30
1	A	145	ARG	NE-CZ-NH1	12.47	126.53	120.30
1	A	147	ARG	NE-CZ-NH1	11.10	125.85	120.30
1	A	114	ARG	NE-CZ-NH2	-10.81	114.89	120.30
1	A	34	TRP	CD1-CG-CD2	10.21	114.47	106.30
1	A	103	ARG	NE-CZ-NH1	10.14	125.37	120.30
1	A	26	TRP	CD1-CG-CD2	10.14	114.41	106.30
1	A	145	ARG	NE-CZ-NH2	-9.08	115.76	120.30
1	A	134	ARG	NE-CZ-NH1	9.04	124.82	120.30
1	A	158	ARG	NE-CZ-NH1	8.55	124.57	120.30
1	A	26	TRP	CE2-CD2-CG	-8.46	100.53	107.30
1	A	34	TRP	CE2-CD2-CG	-8.41	100.57	107.30
1	A	38	ARG	NE-CZ-NH2	-8.28	116.16	120.30
1	A	158	ARG	NE-CZ-NH2	-7.43	116.58	120.30
1	A	103	ARG	CA-CB-CG	7.40	129.68	113.40
1	A	114	ARG	NE-CZ-NH1	6.90	123.75	120.30
1	A	39	TRP	CE2-CD2-CG	-6.72	101.92	107.30
1	A	34	TRP	CB-CG-CD1	-6.70	118.29	127.00

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	MET	CG-SD-CE	-6.41	89.94	100.20
1	A	34	TRP	CG-CD1-NE1	-6.40	103.70	110.10
1	A	142	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	A	39	TRP	CD1-CG-CD2	6.35	111.38	106.30
1	A	34	TRP	CG-CD2-CE3	6.29	139.56	133.90
1	A	26	TRP	CG-CD1-NE1	-6.24	103.86	110.10
1	A	47	VAL	CG1-CB-CG2	-6.16	101.05	110.90
1	A	92	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	A	125	MET	CG-SD-CE	6.01	109.82	100.20
1	A	65	ASP	CB-CG-OD1	6.01	123.71	118.30
1	A	135	VAL	CG1-CB-CG2	-5.67	101.82	110.90
1	A	90	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	84	PRO	N-CA-C	5.44	126.25	112.10
1	A	163	GLN	CA-CB-CG	5.29	125.05	113.40
1	A	128	GLN	CA-CB-CG	5.16	124.75	113.40
1	A	38	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	A	26	TRP	CB-CG-CD1	-5.10	120.36	127.00
1	A	108	MET	CG-SD-CE	5.10	108.35	100.20
1	A	90	ARG	CA-CB-CG	5.08	124.57	113.40
1	A	128	GLN	OE1-CD-NE2	-5.02	110.36	121.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162	TYR	Sidechain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1172	0	1188	13	0
2	A	121	0	0	1	0
All	All	1293	0	1188	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:LEU:HG	1:A:145:ARG:HD3	1.63	0.78
1:A:23:GLY:N	1:A:74:TYR:HH	1.96	0.64
1:A:85:VAL:HG11	1:A:166:ALA:HB2	1.83	0.60
1:A:33:PHE:HA	1:A:63:LEU:HD13	1.88	0.55
1:A:37:LEU:HG	1:A:145:ARG:CD	2.41	0.49
1:A:64:MET:HG3	1:A:108:MET:SD	2.53	0.49
1:A:141:LEU:O	1:A:145:ARG:HG3	2.14	0.48
1:A:54:SER:O	1:A:58:GLN:HG2	2.14	0.47
1:A:52:LEU:HA	1:A:119:ARG:HG3	1.97	0.46
1:A:32:ARG:HH11	1:A:32:ARG:HD2	1.68	0.44
1:A:114:ARG:HD3	1:A:140:HIS:CD2	2.52	0.44
1:A:38:ARG:HD2	2:A:215:HOH:O	2.18	0.43
1:A:83:THR:O	1:A:85:VAL:HG22	2.20	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	142/144 (99%)	136 (96%)	5 (4%)	1 (1%)	<b>22</b> <b>21</b>

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	PRO



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	122/122 (100%)	112 (92%)	10 (8%)	<b>11</b> <b>10</b>

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	SER
1	A	79	GLU
1	A	84	PRO
1	A	88	GLU
1	A	90	ARG
1	A	97	LEU
1	A	98	GLN
1	A	114	ARG
1	A	131	GLU
1	A	137	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	A	128	GLN
1	A	140	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.