



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 03:11 AM EDT

PDB ID : 3LQZ  
Title : Crystal Structure of HLA-DP2  
Authors : Dai, S.  
Deposited on : 2010-02-10  
Resolution : 3.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1482 (3.30-3.22)
Clashscore	180529	1546 (3.30-3.22)
Ramachandran outliers	177936	1536 (3.30-3.22)
Sidechain outliers	177891	1535 (3.30-3.22)
RSRZ outliers	164620	1483 (3.30-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	
2	B	212	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	B	190	X	-	-	-

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 3156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DP alpha 1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	181	1483	958	240	280	5	0	0	0

- Molecule 2 is a protein called HLA-DP2 beta chain linked with DRa peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	198	1631	1027	288	308	8	0	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	ARG	-	SEE REMARK 999	UNP Q5EP54
B	-21	LYS	-	SEE REMARK 999	UNP Q5EP54
B	-20	PHE	-	SEE REMARK 999	UNP Q5EP54
B	-19	HIS	-	SEE REMARK 999	UNP Q5EP54
B	-18	TYR	-	SEE REMARK 999	UNP Q5EP54
B	-17	LEU	-	SEE REMARK 999	UNP Q5EP54
B	-16	PRO	-	SEE REMARK 999	UNP Q5EP54
B	-15	PHE	-	SEE REMARK 999	UNP Q5EP54
B	-14	LEU	-	SEE REMARK 999	UNP Q5EP54
B	-13	PRO	-	SEE REMARK 999	UNP Q5EP54
B	-12	SER	-	SEE REMARK 999	UNP Q5EP54
B	-11	THR	-	SEE REMARK 999	UNP Q5EP54
B	-10	GLY	-	SEE REMARK 999	UNP Q5EP54
B	-9	GLY	-	SEE REMARK 999	UNP Q5EP54
B	-8	SER	-	SEE REMARK 999	UNP Q5EP54
B	-7	LEU	-	SEE REMARK 999	UNP Q5EP54
B	-6	VAL	-	SEE REMARK 999	UNP Q5EP54
B	-5	PRO	-	SEE REMARK 999	UNP Q5EP54
B	-4	ARG	-	SEE REMARK 999	UNP Q5EP54
B	-3	GLY	-	SEE REMARK 999	UNP Q5EP54

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	SER	-	SEE REMARK 999	UNP Q5EP54
B	-1	GLY	-	SEE REMARK 999	UNP Q5EP54
B	0	GLY	-	SEE REMARK 999	UNP Q5EP54
B	1	GLY	-	SEE REMARK 999	UNP Q5EP54
B	2	GLY	-	SEE REMARK 999	UNP Q5EP54
B	3	SER	-	SEE REMARK 999	UNP Q5EP54

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DP alpha 1 chain

Chain A: 



- Molecule 2: HLA-DP2 beta chain linked with DRa peptide

Chain B: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.28Å 157.28Å 61.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 3.25 50.00 – 3.25	Depositor EDS
% Data completeness (in resolution range)	86.2 (50.00-3.25) 99.7 (50.00-3.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 3.25Å)	Xtrriage
Refinement program	REFMAC 5.3.0040	Depositor
R, $R_{free}$	0.217 , 0.251 0.228 , 0.241	Depositor DCC
$R_{free}$ test set	703 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	81.6	Xtrriage
Anisotropy	0.219	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 79.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.058 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	3156	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	93.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/1531	0.71	2/2088 (0.1%)
2	B	0.58	0/1672	0.72	0/2263
All	All	0.59	0/3203	0.71	2/4351 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	LEU	CA-CB-CG	5.42	127.77	115.30
1	A	136	LEU	CA-CB-CG	5.07	126.95	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1483	0	1382	43	0
2	B	1631	0	1542	48	0
3	A	28	0	26	0	0
3	B	14	0	13	0	0
All	All	3156	0	2963	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 13.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:101:PRO:O	2:B:102:SER:HB2	1.82	0.78
2:B:161:MET:HG3	2:B:163:PRO:HD3	1.65	0.77
2:B:31:ASN:O	2:B:32:ARG:HB2	1.83	0.77
2:B:36:VAL:HG22	2:B:37:ARG:H	1.54	0.72
1:A:22:PHE:HB2	1:A:63:ILE:HD13	1.71	0.72
1:A:142:ASP:O	2:B:32:ARG:NH1	2.24	0.69
1:A:17:ARG:HB2	1:A:18:PRO:HD3	1.77	0.66
2:B:101:PRO:O	2:B:102:SER:CB	2.44	0.65
1:A:164:ARG:HG3	1:A:175:LEU:HD12	1.82	0.62
2:B:115:CYS:HB2	2:B:129:TRP:CZ2	2.34	0.62
2:B:63:LYS:HD3	2:B:63:LYS:H	1.65	0.60
1:A:69:ASN:ND2	2:B:-12:SER:H	2.00	0.60
2:B:95:PRO:HD2	2:B:178:LEU:HD21	1.84	0.59
1:A:75:GLN:O	1:A:77:SER:N	2.33	0.57
1:A:76:ARG:NH2	2:B:55:ASP:OD2	2.31	0.57
1:A:89:VAL:O	1:A:176:LYS:HG3	2.05	0.57
1:A:64:ALA:HA	1:A:67:ASN:HD22	1.71	0.56
1:A:69:ASN:HD21	2:B:-13:PRO:HA	1.70	0.55
1:A:62:ASN:HB3	2:B:-15:PHE:CE1	2.41	0.54
2:B:94:GLN:HE21	2:B:177:SER:HB2	1.71	0.54
1:A:17:ARG:CB	1:A:18:PRO:HD3	2.37	0.54
1:A:32:PHE:HB3	1:A:43:TRP:CE3	2.43	0.54
1:A:73:LEU:O	1:A:77:SER:OG	2.23	0.54
1:A:148:PHE:HB3	1:A:150:TYR:CE2	2.44	0.52
1:A:112:PHE:CE2	1:A:146:HIS:HA	2.44	0.52
1:A:57:GLN:HA	1:A:60:LEU:HD12	1.92	0.52
1:A:73:LEU:HD21	2:B:35:PHE:CZ	2.45	0.51
2:B:18:PHE:HB3	2:B:21:THR:CG2	2.41	0.51
2:B:36:VAL:HG22	2:B:37:ARG:N	2.25	0.51
1:A:36:LEU:HD23	1:A:60:LEU:HD23	1.92	0.51
1:A:105:LEU:N	1:A:151:LEU:O	2.41	0.51
2:B:38:PHE:HB2	2:B:45:PHE:CE1	2.45	0.51
2:B:111:ASN:HB3	2:B:161:MET:HB3	1.92	0.51
2:B:161:MET:CG	2:B:163:PRO:HD3	2.40	0.50
1:A:73:LEU:HD13	2:B:9:PHE:HE1	1.76	0.50
2:B:22:GLN:HG2	2:B:77:CYS:HB3	1.93	0.49
1:A:109:ILE:O	1:A:146:HIS:HB2	2.13	0.49
1:A:62:ASN:OD1	2:B:-16:PRO:HA	2.11	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:ARG:NH1	1:A:166:GLU:OE2	2.46	0.49
1:A:13:VAL:HG11	1:A:67:ASN:HA	1.95	0.48
1:A:12:PHE:CE1	1:A:21:GLU:HG2	2.49	0.48
2:B:124:SER:O	2:B:175:HIS:HD2	1.98	0.47
1:A:81:GLN:NE2	2:B:6:ASN:H	2.12	0.47
2:B:174:GLU:HG3	2:B:181:PRO:HB3	1.97	0.46
1:A:62:ASN:HB3	2:B:-15:PHE:HE1	1.80	0.46
2:B:12:ARG:HB2	2:B:27:ARG:HG3	1.98	0.46
2:B:127:VAL:HG22	2:B:173:VAL:HG12	1.98	0.45
2:B:95:PRO:HD2	2:B:178:LEU:CD2	2.46	0.45
2:B:25:LEU:HA	2:B:38:PHE:O	2.16	0.45
2:B:19:ASN:C	2:B:21:THR:H	2.19	0.45
1:A:109:ILE:HD12	1:A:147:LYS:HB3	1.98	0.45
1:A:62:ASN:HA	1:A:65:ILE:HD12	1.99	0.45
1:A:73:LEU:HD13	2:B:9:PHE:CE1	2.52	0.45
1:A:13:VAL:HG12	1:A:70:LEU:HD22	1.99	0.45
1:A:26:PHE:CE2	2:B:88:THR:HB	2.51	0.45
2:B:110:HIS:HD2	2:B:112:LEU:HD13	1.82	0.45
2:B:130:PHE:CE2	2:B:135:GLU:HG2	2.52	0.44
1:A:171:ASP:OD1	1:A:171:ASP:N	2.50	0.44
2:B:144:ASN:HD22	2:B:145:LEU:N	2.16	0.44
1:A:4:ASP:HB2	2:B:17:ALA:O	2.18	0.44
2:B:146:ILE:HG13	2:B:154:GLN:HB3	1.99	0.44
2:B:44:GLU:HB2	2:B:60:ASN:OD1	2.18	0.43
1:A:71:ASN:HA	1:A:74:ILE:HD12	2.00	0.43
1:A:105:LEU:HD23	1:A:105:LEU:HA	1.87	0.42
1:A:96:PRO:HG3	2:B:116:HIS:CG	2.54	0.42
1:A:22:PHE:HB2	1:A:63:ILE:CD1	2.45	0.42
1:A:11:ALA:HB2	1:A:22:PHE:HA	2.02	0.41
2:B:14:GLU:HB2	2:B:16:TYR:HE1	1.85	0.41
1:A:32:PHE:CD2	1:A:43:TRP:CZ3	3.09	0.41
2:B:19:ASN:N	2:B:21:THR:HG22	2.36	0.41
2:B:164:GLN:HG3	2:B:165:GLN:N	2.36	0.41
2:B:169:TYR:O	2:B:185:GLU:HB2	2.21	0.41
1:A:85:ASP:HA	1:A:86:PRO:HD2	1.98	0.40
2:B:22:GLN:HG2	2:B:77:CYS:CB	2.50	0.40
1:A:91:VAL:HA	1:A:106:ILE:O	2.21	0.40
2:B:-17:LEU:HA	2:B:-16:PRO:HD2	1.81	0.40
2:B:53:ARG:N	2:B:54:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/181 (99%)	160 (89%)	13 (7%)	6 (3%)	3	18
2	B	192/212 (91%)	173 (90%)	13 (7%)	6 (3%)	3	19
All	All	371/393 (94%)	333 (90%)	26 (7%)	12 (3%)	3	19

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	17	ARG
2	B	102	SER
1	A	76	ARG
1	A	77	SER
2	B	85	GLY
1	A	137	PHE
1	A	155	PRO
1	A	171	ASP
2	B	101	PRO
2	B	75	ARG
2	B	104	LYS
2	B	73	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/162 (100%)	145 (90%)	17 (10%)	5	21

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	179/188 (95%)	155 (87%)	24 (13%)	3	13
All	All	341/350 (97%)	300 (88%)	41 (12%)	4	17

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LYS
1	A	6	VAL
1	A	16	HIS
1	A	19	THR
1	A	23	MET
1	A	44	HIS
1	A	52	PHE
1	A	53	SER
1	A	57	GLN
1	A	72	THR
1	A	80	THR
1	A	84	ASN
1	A	136	LEU
1	A	140	ARG
1	A	146	HIS
1	A	175	LEU
1	A	179	GLU
2	B	-11	THR
2	B	12	ARG
2	B	25	LEU
2	B	32	ARG
2	B	49	THR
2	B	63	LYS
2	B	76	MET
2	B	89	LEU
2	B	96	ARG
2	B	103	LYS
2	B	110	HIS
2	B	111	ASN
2	B	114	VAL
2	B	118	THR
2	B	125	ILE
2	B	131	LEU
2	B	144	ASN
2	B	146	ILE
2	B	158	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	162	THR
2	B	170	THR
2	B	176	THR
2	B	178	LEU
2	B	182	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	HIS
1	A	67	ASN
1	A	69	ASN
1	A	81	GLN
1	A	146	HIS
1	A	181	GLN
2	B	6	ASN
2	B	22	GLN
2	B	62	GLN
2	B	94	GLN
2	B	110	HIS
2	B	144	ASN
2	B	148	ASN
2	B	154	GLN
2	B	164	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	182	1	14,14,15	0.62	0	17,19,21	1.62	1 (5%)
3	NAG	B	190	2	14,14,15	1.03	1 (7%)	17,19,21	2.03	4 (23%)
3	NAG	A	183	1	14,14,15	0.66	0	17,19,21	1.74	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	182	1	-	1/6/23/26	0/1/1/1
3	NAG	B	190	2	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	A	183	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	190	NAG	C1-C2	2.54	1.55	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	182	NAG	C1-O5-C5	5.66	119.77	112.19
3	A	183	NAG	C1-O5-C5	5.28	119.27	112.19
3	B	190	NAG	C2-N2-C7	4.83	129.38	122.90
3	B	190	NAG	C4-C3-C2	4.65	117.83	111.02
3	A	183	NAG	C4-C3-C2	3.13	115.60	111.02
3	A	183	NAG	C3-C4-C5	2.28	114.36	110.23
3	B	190	NAG	O5-C1-C2	-2.25	107.80	111.29
3	B	190	NAG	C3-C4-C5	2.24	114.29	110.23

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	190	NAG	C1

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	183	NAG	O5-C5-C6-O6
3	B	190	NAG	C4-C5-C6-O6
3	A	183	NAG	C4-C5-C6-O6
3	A	182	NAG	O5-C5-C6-O6
3	B	190	NAG	O5-C5-C6-O6
3	B	190	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/181 (100%)	0.26	2 (1%) 77 66	85, 93, 98, 105	0
2	B	198/212 (93%)	0.63	8 (4%) 43 32	85, 93, 99, 104	0
All	All	379/393 (96%)	0.45	10 (2%) 57 43	85, 93, 99, 105	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	-8	SER	3.0
1	A	15	THR	2.6
2	B	31	ASN	2.6
2	B	103	LYS	2.4
2	B	61	SER	2.4
2	B	186	TRP	2.3
1	A	158	GLU	2.2
2	B	-20	PHE	2.1
2	B	144	ASN	2.0
2	B	112	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	183	14/15	0.53	0.18	103,106,107,107	0
3	NAG	B	190	14/15	0.57	0.18	112,113,116,117	0
3	NAG	A	182	14/15	0.72	0.16	95,96,97,97	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.