



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 12, 2023 – 03:43 AM EDT

PDB ID : 4LQI
Title : Yeast 20S Proteasome in complex with Vibralactone
Authors : List, A.; Zeiler, E.; Gallastegui, N.; Rusch, M.; Hedberg, C.; Sieber, S.A.; Groll, M.
Deposited on : 2013-07-18
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	 2% 76% 23% .
1	O	250	 2% 74% 25% .
2	B	244	 2% 64% 31% 5%
2	P	244	 3% 66% 30% 5%
3	C	241	 2% 60% 37% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	241	
4	D	242	
4	R	242	
5	E	233	
5	S	233	
6	F	244	
6	T	244	
7	G	243	
7	U	243	
8	H	222	
8	V	222	
9	I	204	
9	W	204	
10	J	198	
10	X	198	
11	K	212	
11	Y	212	
12	L	222	
12	Z	222	
13	M	233	
13	a	233	
14	N	196	
14	b	196	

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 50942 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	Total 1915	C 1219	N 315	O 377	S 4	0	0	0
1	O	250	Total 1915	C 1219	N 315	O 377	S 4	0	0	0

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	Total 1905	C 1201	N 321	O 380	S 3	0	0	0
2	P	244	Total 1905	C 1201	N 321	O 380	S 3	0	0	0

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	241	Total 1891	C 1181	N 331	O 375	S 4	0	0	0
3	Q	241	Total 1891	C 1181	N 331	O 375	S 4	0	0	0

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	242	Total 1862	C 1162	N 314	O 379	S 7	0	0	0
4	R	242	Total 1862	C 1162	N 314	O 379	S 7	0	0	0

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	233	Total 1795	C 1129	N 312	O 350	S 4	0	0	0
5	S	233	Total 1795	C 1129	N 312	O 350	S 4	0	0	0

- Molecule 6 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	244	Total 1897	C 1205	N 330	O 358	S 4	0	0	0
6	T	244	Total 1897	C 1205	N 330	O 358	S 4	0	0	0

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	243	Total 1921	C 1221	N 322	O 370	S 8	0	0	0
7	U	243	Total 1921	C 1221	N 322	O 370	S 8	0	0	0

- Molecule 8 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	222	Total 1685	C 1061	N 293	O 324	S 7	0	0	0
8	V	222	Total 1685	C 1061	N 293	O 324	S 7	0	0	0

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	204	Total 1581	C 1010	N 258	O 305	S 8	0	0	0
9	W	204	Total 1581	C 1010	N 258	O 305	S 8	0	0	0

- Molecule 10 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	J	198	Total 1585	C 1005	N 269	O 305	S 6	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

- Molecule 11 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			


- Molecule 13 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	a	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is vibrallactone, bound form (three-letter code: 1Y9) (formula: C₁₂H₁₈O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	K	1	Total	C O	0	0
			15	12 3		
15	N	1	Total	C O	0	0
			15	12 3		
15	Y	1	Total	C O	0	0
			15	12 3		
15	b	1	Total	C O	0	0
			15	12 3		

- Molecule 16 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	56	Total	O	0	0
			56	56		
16	B	36	Total	O	0	0
			36	36		
16	C	42	Total	O	0	0
			42	42		
16	D	38	Total	O	0	0
			38	38		
16	E	23	Total	O	0	0
			23	23		
16	F	49	Total	O	0	0
			49	49		
16	G	61	Total	O	0	0
			61	61		
16	H	49	Total	O	0	0
			49	49		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	I	70	Total 70	O 70	0	0
16	J	53	Total 53	O 53	0	0
16	K	46	Total 46	O 46	0	0
16	L	56	Total 56	O 56	0	0
16	M	73	Total 73	O 73	0	0
16	N	57	Total 57	O 57	0	0
16	O	31	Total 31	O 31	0	0
16	P	28	Total 28	O 28	0	0
16	Q	28	Total 28	O 28	0	0
16	R	31	Total 31	O 31	0	0
16	S	20	Total 20	O 20	0	0
16	T	38	Total 38	O 38	0	0
16	U	62	Total 62	O 62	0	0
16	V	46	Total 46	O 46	0	0
16	W	61	Total 61	O 61	0	0
16	X	52	Total 52	O 52	0	0
16	Y	46	Total 46	O 46	0	0
16	Z	52	Total 52	O 52	0	0
16	a	75	Total 75	O 75	0	0
16	b	55	Total 55	O 55	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha type-2



- Molecule 1: Proteasome subunit alpha type-2




- Molecule 2: Proteasome subunit alpha type-3




- Molecule 2: Proteasome subunit alpha type-3

Chain P:  3% 66% 30% 5%




• Molecule 3: Proteasome subunit alpha type-4

Chain C:  2% 60% 37% .



• Molecule 3: Proteasome subunit alpha type-4


Chain Q:  7% 60% 37% .




• Molecule 4: Proteasome subunit alpha type-5

Chain D:  3% 70% 28% .







• Molecule 4: Proteasome subunit alpha type-5




• Molecule 5: Proteasome subunit alpha type-6




• Molecule 5: Proteasome subunit alpha type-6




- Molecule 6: Proteasome subunit alpha type-7




- Molecule 6: Proteasome subunit alpha type-7




- Molecule 7: Proteasome subunit alpha type-1




- Molecule 7: Proteasome subunit alpha type-1




- Molecule 8: Proteasome subunit beta type-2




- Molecule 8: Proteasome subunit beta type-2




- Molecule 9: Proteasome subunit beta type-3



- Molecule 9: Proteasome subunit beta type-3




- Molecule 10: Proteasome subunit beta type-4






- Molecule 10: Proteasome subunit beta type-4




- Molecule 11: Proteasome subunit beta type-5



- Molecule 11: Proteasome subunit beta type-5



- Molecule 12: Proteasome subunit beta type-6




- Molecule 12: Proteasome subunit beta type-6








• Molecule 13: Proteasome subunit beta type-7



• Molecule 13: Proteasome subunit beta type-7



• Molecule 14: Proteasome subunit beta type-1



• Molecule 14: Proteasome subunit beta type-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	135.86Å 299.40Å 144.99Å 90.00° 113.07° 90.00°	Depositor
Resolution (Å)	14.99 – 2.70 48.95 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.4 (14.99-2.70) 99.5 (48.95-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.69Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.202 , 0.227 0.200 , 0.223	Depositor DCC
R_{free} test set	14347 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	53.8	Xtrriage
Anisotropy	0.619	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	50942	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1Y9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1952	0.63	0/2642
1	O	0.38	0/1952	0.63	0/2642
2	B	0.38	0/1935	0.62	0/2618
2	P	0.39	0/1935	0.62	0/2618
3	C	0.36	0/1920	0.62	1/2598 (0.0%)
3	Q	0.36	0/1920	0.62	0/2598
4	D	0.35	0/1887	0.62	0/2541
4	R	0.37	0/1887	0.62	0/2541
5	E	0.36	0/1823	0.60	0/2463
5	S	0.36	0/1823	0.60	0/2463
6	F	0.38	0/1937	0.62	1/2614 (0.0%)
6	T	0.38	0/1937	0.62	1/2614 (0.0%)
7	G	0.40	0/1959	0.62	0/2652
7	U	0.40	0/1959	0.62	0/2652
8	H	0.40	0/1716	0.66	0/2326
8	V	0.39	0/1716	0.66	0/2326
9	I	0.40	0/1611	0.66	0/2174
9	W	0.40	0/1611	0.67	0/2174
10	J	0.40	0/1613	0.64	0/2173
10	X	0.40	0/1613	0.65	0/2173
11	K	0.41	0/1681	0.65	1/2274 (0.0%)
11	Y	0.41	0/1681	0.65	0/2274
12	L	0.40	0/1795	0.66	1/2420 (0.0%)
12	Z	0.40	0/1795	0.66	1/2420 (0.0%)
13	M	0.39	0/1855	0.66	0/2514
13	a	0.39	0/1855	0.66	1/2514 (0.0%)
14	N	0.40	0/1541	0.63	0/2087
14	b	0.39	0/1541	0.62	0/2087
All	All	0.39	0/50450	0.64	7/68192 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	L	95	TYR	N-CA-C	-5.26	96.80	111.00
12	Z	95	TYR	N-CA-C	-5.23	96.89	111.00
6	T	135	SER	N-CA-C	-5.21	96.94	111.00
6	F	135	SER	N-CA-C	-5.20	96.96	111.00
13	a	95	LEU	N-CA-C	-5.18	97.02	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1926	54	0
1	O	1915	0	1926	58	0
2	B	1905	0	1901	80	0
2	P	1905	0	1901	75	0
3	C	1891	0	1900	95	0
3	Q	1891	0	1900	99	0
4	D	1862	0	1836	51	0
4	R	1862	0	1836	53	0
5	E	1795	0	1797	88	0
5	S	1795	0	1797	85	0
6	F	1897	0	1886	62	0
6	T	1897	0	1886	61	0
7	G	1921	0	1910	66	0
7	U	1921	0	1910	74	0
8	H	1685	0	1688	40	0
8	V	1685	0	1688	42	0
9	I	1581	0	1574	36	0
9	W	1581	0	1574	39	0
10	J	1585	0	1590	64	0
10	X	1585	0	1590	68	0
11	K	1644	0	1594	49	0
11	Y	1644	0	1594	51	0
12	L	1757	0	1711	52	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	Z	1757	0	1711	54	0
13	M	1824	0	1832	52	0
13	a	1824	0	1832	0	0
14	N	1512	0	1480	36	0
14	b	1512	0	1480	0	0
15	K	15	0	17	2	0
15	N	15	0	17	3	0
15	Y	15	0	17	2	0
15	b	15	0	17	0	0
16	A	56	0	0	0	0
16	B	36	0	0	2	0
16	C	42	0	0	2	0
16	D	38	0	0	2	0
16	E	23	0	0	0	0
16	F	49	0	0	2	0
16	G	61	0	0	3	0
16	H	49	0	0	4	0
16	I	70	0	0	3	0
16	J	53	0	0	3	0
16	K	46	0	0	2	0
16	L	56	0	0	1	0
16	M	73	0	0	1	0
16	N	57	0	0	4	0
16	O	31	0	0	2	0
16	P	28	0	0	2	0
16	Q	28	0	0	4	0
16	R	31	0	0	1	0
16	S	20	0	0	0	0
16	T	38	0	0	3	0
16	U	62	0	0	3	0
16	V	46	0	0	3	0
16	W	61	0	0	2	0
16	X	52	0	0	3	0
16	Y	46	0	0	1	0
16	Z	52	0	0	3	0
16	a	75	0	0	0	0
16	b	55	0	0	0	0
All	All	50942	0	49318	1418	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 1418 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:202:THR:HG22	2:P:204:SER:H	1.13	1.09
2:B:202:THR:HG22	2:B:204:SER:H	1.14	1.07
7:G:96:ALA:HA	7:G:107:MET:HE2	1.40	1.03
7:U:96:ALA:HA	7:U:107:MET:HE2	1.43	1.00
11:K:105(B):LYS:H	11:K:105(B):LYS:HD2	1.26	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	235 (95%)	11 (4%)	2 (1%)	19	43
1	O	248/250 (99%)	235 (95%)	11 (4%)	2 (1%)	19	43
2	B	242/244 (99%)	219 (90%)	16 (7%)	7 (3%)	4	10
2	P	242/244 (99%)	219 (90%)	17 (7%)	6 (2%)	5	14
3	C	239/241 (99%)	226 (95%)	10 (4%)	3 (1%)	12	30
3	Q	239/241 (99%)	225 (94%)	11 (5%)	3 (1%)	12	30
4	D	240/242 (99%)	223 (93%)	13 (5%)	4 (2%)	9	23
4	R	240/242 (99%)	221 (92%)	15 (6%)	4 (2%)	9	23
5	E	231/233 (99%)	208 (90%)	16 (7%)	7 (3%)	4	10
5	S	231/233 (99%)	208 (90%)	16 (7%)	7 (3%)	4	10
6	F	242/244 (99%)	231 (96%)	9 (4%)	2 (1%)	19	43
6	T	242/244 (99%)	231 (96%)	9 (4%)	2 (1%)	19	43
7	G	241/243 (99%)	227 (94%)	13 (5%)	1 (0%)	34	60
7	U	241/243 (99%)	227 (94%)	13 (5%)	1 (0%)	34	60
8	H	220/222 (99%)	209 (95%)	9 (4%)	2 (1%)	17	40
8	V	220/222 (99%)	209 (95%)	8 (4%)	3 (1%)	11	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
9	I	202/204 (99%)	193 (96%)	9 (4%)	0	100	100
9	W	202/204 (99%)	193 (96%)	9 (4%)	0	100	100
10	J	196/198 (99%)	186 (95%)	9 (5%)	1 (0%)	29	54
10	X	196/198 (99%)	187 (95%)	8 (4%)	1 (0%)	29	54
11	K	210/212 (99%)	202 (96%)	8 (4%)	0	100	100
11	Y	210/212 (99%)	203 (97%)	7 (3%)	0	100	100
12	L	220/222 (99%)	206 (94%)	13 (6%)	1 (0%)	29	54
12	Z	220/222 (99%)	207 (94%)	12 (6%)	1 (0%)	29	54
13	M	231/233 (99%)	219 (95%)	11 (5%)	1 (0%)	34	60
13	a	231/233 (99%)	219 (95%)	11 (5%)	1 (0%)	34	60
14	N	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
14	b	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
All	All	6312/6368 (99%)	5946 (94%)	304 (5%)	62 (1%)	15	37

5 of 62 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	54	VAL
2	B	218(C)	ASP
3	C	58	LEU
3	C	203	THR
4	D	123(G)	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	205 (98%)	4 (2%)	57	82
1	O	209/209 (100%)	205 (98%)	4 (2%)	57	82
2	B	203/203 (100%)	193 (95%)	10 (5%)	25	52
2	P	203/203 (100%)	192 (95%)	11 (5%)	22	47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	213/213 (100%)	203 (95%)	10 (5%)	26	54
3	Q	213/213 (100%)	203 (95%)	10 (5%)	26	54
4	D	198/198 (100%)	186 (94%)	12 (6%)	18	41
4	R	198/198 (100%)	186 (94%)	12 (6%)	18	41
5	E	192/192 (100%)	176 (92%)	16 (8%)	11	25
5	S	192/192 (100%)	177 (92%)	15 (8%)	12	29
6	F	201/201 (100%)	186 (92%)	15 (8%)	13	31
6	T	201/201 (100%)	186 (92%)	15 (8%)	13	31
7	G	207/207 (100%)	195 (94%)	12 (6%)	20	43
7	U	207/207 (100%)	195 (94%)	12 (6%)	20	43
8	H	181/181 (100%)	174 (96%)	7 (4%)	32	61
8	V	181/181 (100%)	174 (96%)	7 (4%)	32	61
9	I	172/172 (100%)	169 (98%)	3 (2%)	60	84
9	W	172/172 (100%)	169 (98%)	3 (2%)	60	84
10	J	175/175 (100%)	169 (97%)	6 (3%)	37	66
10	X	175/175 (100%)	169 (97%)	6 (3%)	37	66
11	K	169/169 (100%)	159 (94%)	10 (6%)	19	43
11	Y	169/169 (100%)	160 (95%)	9 (5%)	22	48
12	L	185/185 (100%)	174 (94%)	11 (6%)	19	43
12	Z	185/185 (100%)	173 (94%)	12 (6%)	17	38
13	M	199/199 (100%)	191 (96%)	8 (4%)	31	60
13	a	199/199 (100%)	191 (96%)	8 (4%)	31	60
14	N	162/162 (100%)	155 (96%)	7 (4%)	29	57
14	b	162/162 (100%)	155 (96%)	7 (4%)	29	57
All	All	5332/5332 (100%)	5070 (95%)	262 (5%)	25	52

5 of 262 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	Y	69	ARG
12	Z	39	ASN
14	b	119	VAL
11	K	9	GLN
10	J	121	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 201 such sidechains are listed below:

Mol	Chain	Res	Type
3	Q	125	GLN
7	U	87	ASN
14	b	145	ASN
4	R	23	GLN
5	S	125	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
15	1Y9	K	301	11	12,15,15	2.43	2 (16%)	7,21,21	1.71	1 (14%)
15	1Y9	N	201	14	12,15,15	2.31	1 (8%)	7,21,21	2.07	1 (14%)
15	1Y9	b	201	14	12,15,15	2.30	1 (8%)	7,21,21	1.91	1 (14%)
15	1Y9	Y	301	11	12,15,15	2.41	2 (16%)	7,21,21	1.64	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	1Y9	K	301	11	-	5/10/26/26	0/1/1/1
15	1Y9	N	201	14	-	5/10/26/26	0/1/1/1
15	1Y9	b	201	14	-	5/10/26/26	0/1/1/1
15	1Y9	Y	301	11	-	4/10/26/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	K	301	1Y9	O5-C4	7.59	1.44	1.19
15	N	201	1Y9	O5-C4	7.37	1.44	1.19
15	b	201	1Y9	O5-C4	7.27	1.43	1.19
15	Y	301	1Y9	O5-C4	7.23	1.43	1.19
15	Y	301	1Y9	C15-C2	2.99	1.55	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	N	201	1Y9	C3-C6-C7	5.02	120.66	113.47
15	b	201	1Y9	C3-C6-C7	4.53	119.95	113.47
15	K	301	1Y9	C3-C6-C7	4.13	119.38	113.47
15	Y	301	1Y9	C3-C6-C7	3.85	118.98	113.47

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
15	K	301	1Y9	C2-C3-C4-O5
15	K	301	1Y9	C6-C3-C4-O5
15	N	201	1Y9	C2-C3-C4-O5
15	N	201	1Y9	C6-C3-C4-O5
15	Y	301	1Y9	C2-C3-C4-O5

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	K	301	1Y9	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	N	201	1Y9	3	0
15	Y	301	1Y9	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.26	3 (1%) 79 80	44, 58, 89, 114	0
1	O	250/250 (100%)	-0.22	5 (2%) 65 67	44, 59, 91, 114	0
2	B	244/244 (100%)	-0.15	4 (1%) 72 74	42, 62, 102, 124	0
2	P	244/244 (100%)	-0.06	8 (3%) 46 46	42, 63, 103, 124	0
3	C	241/241 (100%)	-0.06	5 (2%) 63 65	45, 67, 117, 132	0
3	Q	241/241 (100%)	0.13	16 (6%) 18 16	47, 69, 118, 133	0
4	D	242/242 (100%)	-0.02	7 (2%) 51 52	45, 66, 98, 133	0
4	R	242/242 (100%)	-0.10	6 (2%) 57 59	46, 67, 97, 133	0
5	E	233/233 (100%)	0.01	7 (3%) 50 51	49, 69, 95, 121	0
5	S	233/233 (100%)	-0.02	6 (2%) 56 57	49, 70, 95, 120	0
6	F	244/244 (100%)	-0.25	4 (1%) 72 74	45, 62, 99, 114	0
6	T	244/244 (100%)	-0.07	5 (2%) 65 67	46, 63, 99, 115	0
7	G	243/243 (100%)	-0.32	4 (1%) 72 74	40, 57, 87, 123	0
7	U	243/243 (100%)	-0.34	3 (1%) 79 80	39, 57, 86, 124	0
8	H	222/222 (100%)	-0.33	0 100 100	41, 55, 77, 108	0
8	V	222/222 (100%)	-0.40	2 (0%) 84 85	42, 56, 78, 110	0
9	I	204/204 (100%)	-0.40	0 100 100	39, 53, 71, 84	0
9	W	204/204 (100%)	-0.30	0 100 100	40, 53, 72, 85	0
10	J	198/198 (100%)	-0.33	3 (1%) 73 76	42, 55, 74, 133	0
10	X	198/198 (100%)	-0.32	5 (2%) 57 59	43, 56, 73, 133	0
11	K	212/212 (100%)	-0.41	0 100 100	39, 54, 73, 83	0
11	Y	212/212 (100%)	-0.43	0 100 100	40, 54, 74, 84	0
12	L	222/222 (100%)	-0.41	1 (0%) 91 92	38, 55, 78, 97	0
12	Z	222/222 (100%)	-0.36	1 (0%) 91 92	39, 55, 76, 97	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	233/233 (100%)	-0.44	1 (0%) 92 93	39, 56, 71, 79	0
13	a	233/233 (100%)	-0.48	1 (0%) 92 93	39, 55, 71, 79	0
14	N	196/196 (100%)	-0.49	0 100 100	39, 51, 72, 85	0
14	b	196/196 (100%)	-0.41	0 100 100	40, 51, 72, 85	0
All	All	6368/6368 (100%)	-0.25	97 (1%) 73 76	38, 58, 91, 133	0

The worst 5 of 97 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
7	U	6	ALA	11.1
4	D	123(E)	SER	10.9
4	D	123(D)	ALA	10.9
4	D	123(F)	GLY	9.2
10	X	193	GLN	9.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	1Y9	Y	301	15/15	0.90	0.25	67,75,78,80	0
15	1Y9	K	301	15/15	0.91	0.23	68,74,78,79	0
15	1Y9	N	201	15/15	0.92	0.26	65,71,74,75	0
15	1Y9	b	201	15/15	0.92	0.26	65,73,76,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.