



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2020 – 02:16 PM JST

PDB ID : 6LTR
Title : Crystal structure of Cas12i2 ternary complex with single Mg²⁺ bound in catalytic pocket
Authors : Huang, X.; Sun, W.; Cheng, Z.; Chen, M.; Li, X.; Wang, J.; Sheng, G.; Gong, W.; Wang, Y.
Deposited on : 2020-01-23
Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)
RNA backbone	3102	1008 (2.84-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1055	
2	B	56	
3	D	13	
4	C	35	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	6	 33% 50% 17%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 10800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cas12i2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1037	8327	5299	1470	1531	27	0	0	0

- Molecule 2 is a RNA chain called RNA (56-mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	49	1045	469	191	337	48	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(*GP*CP*CP*GP*CP*TP*TP*TP*CP*TP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	11	217	106	32	69	10	0	0	0

- Molecule 4 is a DNA chain called DNA (35-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	C	35	715	341	127	212	35	0	0	0

- Molecule 5 is a DNA chain called trans ssDNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	E	5	99	47	19	28	5	0	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Mg 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	259	Total O 259 259	0	0
8	B	65	Total O 65 65	0	0
8	D	20	Total O 20 20	0	0
8	C	47	Total O 47 47	0	0
8	E	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cas12i2



- Molecule 2: RNA (56-mer)




- Molecule 3: DNA (5'-D(*GP*CP*CP*GP*CP*TP*TP*TP*CP*TP*TP*T)-3')



- Molecule 4: DNA (35-MER)



- Molecule 5: trans ssDNA



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	93.26Å 123.03Å 280.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.74 – 2.51 46.74 – 2.51	Depositor EDS
% Data completeness (in resolution range)	98.4 (46.74-2.51) 98.4 (46.74-2.51)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.61 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874, PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.198 , 0.226 0.198 , 0.226	Depositor DCC
R_{free} test set	2746 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtrriage
Anisotropy	0.015	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10800	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	2/8496 (0.0%)	0.61	4/11469 (0.0%)
2	B	0.53	3/1170 (0.3%)	0.90	2/1822 (0.1%)
3	D	0.55	0/240	1.11	0/368
4	C	0.71	1/800 (0.1%)	1.23	8/1232 (0.6%)
5	E	1.98	5/110 (4.5%)	0.88	0/166
All	All	0.50	11/10816 (0.1%)	0.74	14/15057 (0.1%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	23	DC	O3'-P	-12.04	1.46	1.61
5	E	24	DC	O3'-P	-9.43	1.49	1.61
1	A	949	GLU	CD-OE1	7.55	1.33	1.25
5	E	24	DC	P-OP2	-7.30	1.36	1.49
2	B	45	U	O3'-P	-6.58	1.53	1.61
2	B	46	U	O3'-P	-6.23	1.53	1.61
1	A	949	GLU	CD-OE2	5.71	1.31	1.25
2	B	44	A	O3'-P	-5.61	1.54	1.61
4	C	4	DA	O3'-P	-5.61	1.54	1.61
5	E	23	DC	P-OP2	-5.52	1.39	1.49
5	E	23	DC	P-OP1	-5.47	1.39	1.49

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	47	A	O5'-P-OP2	-9.36	97.28	105.70
4	C	5	DA	O5'-P-OP1	-8.79	97.79	105.70
4	C	4	DA	C2'-C3'-O3'	-8.49	84.59	112.60
4	C	4	DA	C4'-C3'-O3'	-7.77	90.27	109.70
4	C	5	DA	O5'-P-OP2	-7.15	99.27	105.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	430	LYS	CD-CE-NZ	6.35	126.30	111.70
1	A	430	LYS	CA-CB-CG	6.11	126.85	113.40
4	C	5	DA	OP1-P-OP2	6.08	128.72	119.60
1	A	795	LEU	CB-CG-CD2	-5.76	101.21	111.00
2	B	47	A	C3'-C2'-C1'	5.65	106.02	101.50
4	C	4	DA	C8-N9-C1'	5.27	137.18	127.70
4	C	4	DA	C4-N9-C1'	-5.16	117.02	126.30
4	C	7	DG	O4'-C4'-C3'	-5.09	102.47	104.50
1	A	922	ARG	NE-CZ-NH2	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8327	0	8308	93	0
2	B	1045	0	529	10	0
3	D	217	0	128	4	0
4	C	715	0	396	12	0
5	E	99	0	56	1	0
6	A	4	0	6	0	0
7	A	1	0	0	0	0
8	A	259	0	0	6	0
8	B	65	0	0	0	0
8	C	47	0	0	0	0
8	D	20	0	0	0	0
8	E	1	0	0	0	0
All	All	10800	0	9423	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:46:U:H6	2:B:46:U:H5''	1.18	1.07
2:B:46:U:C6	2:B:46:U:H5''	2.02	0.93
1:A:184:LEU:HG	1:A:188:LYS:HE2	1.54	0.88
1:A:374:PRO:HB3	1:A:408:ARG:HH12	1.42	0.84
4:C:5:DA:H2''	4:C:6:DT:H5'	1.63	0.80
1:A:688:GLU:O	1:A:690:LYS:N	2.19	0.76
1:A:602:GLN:HB2	1:A:751:LEU:HD22	1.74	0.70
1:A:906:VAL:HA	1:A:909:ILE:HD12	1.75	0.68
1:A:533:LYS:O	1:A:536:VAL:HG13	1.94	0.67
1:A:580:PHE:HZ	1:A:869:MET:HG2	1.59	0.67
1:A:524:THR:HG23	1:A:526:GLN:H	1.59	0.66
1:A:945:LEU:HD11	1:A:972:LEU:HD21	1.76	0.66
1:A:178:GLU:HB3	3:D:8:DT:H5''	1.81	0.63
1:A:428:ARG:NH2	8:A:1204:HOH:O	2.31	0.62
1:A:963:ILE:CG1	1:A:967:VAL:HG21	2.29	0.62
1:A:374:PRO:HB3	1:A:408:ARG:NH1	2.14	0.62
1:A:612:GLU:HB2	1:A:630:ILE:HD11	1.81	0.61
1:A:711:ALA:HA	1:A:763:ILE:HD11	1.84	0.59
1:A:313:PHE:HE1	1:A:432:GLN:HG2	1.67	0.59
4:C:5:DA:H2'	4:C:6:DT:C6	2.39	0.57
1:A:421:LYS:HE2	1:A:425:GLN:NE2	2.20	0.56
1:A:277:ARG:NH2	8:A:1203:HOH:O	2.30	0.56
1:A:306:LYS:HG3	1:A:437:SER:HA	1.86	0.56
1:A:130:ASP:HA	1:A:133:ARG:HD2	1.87	0.55
1:A:368:LYS:O	1:A:372:ASP:N	2.39	0.55
1:A:688:GLU:O	1:A:688:GLU:HG3	2.07	0.55
1:A:688:GLU:C	1:A:690:LYS:H	2.10	0.54
2:B:46:U:C6	2:B:46:U:C5'	2.83	0.54
1:A:703:LEU:HD21	1:A:741:CYS:HB3	1.90	0.52
1:A:580:PHE:CZ	1:A:869:MET:HG2	2.43	0.52
1:A:964:PRO:O	1:A:967:VAL:HG22	2.10	0.51
1:A:967:VAL:HG23	1:A:968:LEU:N	2.25	0.51
1:A:185:ARG:HH11	1:A:185:ARG:HG2	1.76	0.51
4:C:32:DC:H2'	4:C:33:DG:C8	2.45	0.51
1:A:365:LYS:O	1:A:368:LYS:HG2	2.11	0.51
3:D:7:DT:H2'	3:D:8:DT:C6	2.45	0.51
1:A:255:THR:HG22	1:A:259:LYS:HE3	1.93	0.50
1:A:54:PHE:CZ	8:A:1341:HOH:O	2.55	0.50
1:A:54:PHE:CE1	8:A:1341:HOH:O	2.64	0.50
1:A:471:LEU:HB3	1:A:473:LEU:HD21	1.93	0.50
1:A:782:ASP:HB2	4:C:16:DC:OP1	2.12	0.50
1:A:991:ARG:NH1	1:A:1019:ASP:OD2	2.46	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:ILE:O	1:A:190:ILE:HG23	2.12	0.49
1:A:327:LEU:HD21	1:A:421:LYS:HD3	1.94	0.49
1:A:479:ASP:OD2	1:A:481:ARG:HD2	2.13	0.49
1:A:679:ASN:HD21	4:C:5:DA:P	2.36	0.49
1:A:588:THR:O	1:A:590:GLN:NE2	2.46	0.48
1:A:77:VAL:O	1:A:119:TYR:HA	2.12	0.48
1:A:764:ILE:HG23	1:A:792:LEU:HG	1.95	0.48
1:A:845:LYS:HE3	2:B:35:C:OP2	2.14	0.48
1:A:253:LEU:O	1:A:257:LEU:HG	2.14	0.48
1:A:56:THR:HG21	1:A:63:GLN:HB2	1.95	0.48
1:A:375:ALA:O	1:A:377:PRO:HD3	2.14	0.47
1:A:603:THR:HG23	5:E:25:DC:H5''	1.96	0.47
1:A:504:SER:OG	1:A:507:ASP:HB2	2.13	0.47
1:A:332:VAL:HG12	1:A:411:CYS:SG	2.55	0.47
2:B:48:U:H2'	2:B:49:U:C6	2.49	0.47
1:A:104:SER:O	1:A:108:LYS:HG2	2.14	0.47
1:A:921:LEU:HD13	1:A:936:VAL:CG1	2.45	0.46
1:A:507:ASP:HB3	1:A:555:PRO:HG3	1.97	0.46
1:A:968:LEU:HD23	1:A:972:LEU:HD11	1.97	0.46
1:A:403:TYR:O	1:A:406:THR:HB	2.16	0.46
1:A:678:LYS:NZ	1:A:684:ILE:HD11	2.30	0.46
1:A:940:LEU:CD1	1:A:949:GLU:HG3	2.46	0.46
1:A:845:LYS:HE2	1:A:845:LYS:HB2	1.68	0.45
1:A:378:GLU:O	1:A:382:VAL:HG13	2.16	0.45
4:C:1:DA:N6	4:C:2:DA:N6	2.65	0.45
1:A:614:VAL:O	1:A:626:PHE:HB2	2.16	0.45
1:A:191:LEU:HA	1:A:194:ILE:HG22	1.98	0.45
1:A:177:LYS:NZ	3:D:7:DT:OP1	2.48	0.45
1:A:367:TYR:CD1	1:A:413:ALA:HB2	2.52	0.45
1:A:365:LYS:HG2	1:A:368:LYS:HZ2	1.82	0.45
3:D:5:DC:H2'	3:D:6:DT:C6	2.52	0.44
1:A:656:TYR:HB3	1:A:657:PRO:HD3	1.98	0.44
1:A:190:ILE:HG22	1:A:214:VAL:HG11	1.98	0.44
1:A:841:ASN:N	2:B:33:A:OP1	2.50	0.44
1:A:524:THR:HG23	1:A:526:GLN:N	2.30	0.44
1:A:223:ARG:HH21	4:C:31:DG:H5'	1.82	0.44
1:A:953:LEU:O	1:A:956:ARG:NH1	2.45	0.44
1:A:493:ARG:NH1	8:A:1211:HOH:O	2.46	0.44
1:A:728:GLN:HA	1:A:731:GLN:OE1	2.18	0.44
1:A:848:ASN:O	1:A:852:MET:HG3	2.17	0.44
1:A:741:CYS:HA	1:A:746:LEU:HD12	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:45:U:H2'	2:B:46:U:C6	2.53	0.43
1:A:369:ALA:HA	1:A:372:ASP:HB3	2.01	0.43
1:A:880:LEU:HG	1:A:881:TYR:CD2	2.54	0.43
1:A:909:ILE:HG23	1:A:913:VAL:HG11	2.00	0.43
1:A:831:ARG:NH2	8:A:1207:HOH:O	2.40	0.43
1:A:956:ARG:HB2	1:A:959:ARG:HG3	2.00	0.43
2:B:35:C:H2'	2:B:36:A:C8	2.53	0.43
1:A:664:LYS:HE2	1:A:664:LYS:HB2	1.73	0.43
1:A:670:VAL:HG21	1:A:713:LEU:HD21	2.00	0.43
1:A:355:VAL:HG12	1:A:360:GLY:HA2	2.00	0.43
4:C:18:DT:H2''	4:C:19:DG:H8	1.83	0.42
1:A:666:ALA:HB1	1:A:710:TYR:CE1	2.53	0.42
1:A:656:TYR:HA	1:A:659:TYR:CE1	2.54	0.42
4:C:13:DC:H2'	4:C:14:DT:C6	2.54	0.42
1:A:730:ARG:NH1	1:A:789:ASP:OD2	2.50	0.42
4:C:13:DC:H2''	4:C:14:DT:H5'	2.01	0.42
1:A:906:VAL:HA	1:A:909:ILE:CD1	2.47	0.42
4:C:1:DA:N6	4:C:2:DA:C6	2.87	0.42
1:A:493:ARG:HB2	2:B:7:C:C4	2.55	0.42
1:A:807:LYS:HB3	1:A:855:LEU:HD11	2.02	0.41
1:A:934:GLN:O	1:A:937:LYS:HB2	2.20	0.41
4:C:3:DT:H2''	4:C:4:DA:C5'	2.50	0.41
1:A:313:PHE:CE1	1:A:432:GLN:HG2	2.50	0.41
1:A:370:TRP:CZ3	1:A:405:PHE:HA	2.56	0.41
1:A:670:VAL:HG13	1:A:690:LYS:HD3	2.03	0.40
1:A:94:LYS:HG3	1:A:98:TYR:CE2	2.57	0.40
1:A:515:ARG:HD3	2:B:23:G:OP1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1035/1055 (98%)	1001 (97%)	31 (3%)	3 (0%)	41 61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	689	ALA
1	A	377	PRO
1	A	680	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	888/930 (96%)	886 (100%)	2 (0%)	93 98

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	686	THR
1	A	735	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	679	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	48/56 (85%)	15 (31%)	1 (2%)

All (15) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	2	G
2	B	3	A
2	B	4	A
2	B	6	U
2	B	7	C
2	B	13	U
2	B	18	U
2	B	24	A
2	B	33	A
2	B	39	G
2	B	42	A
2	B	46	U
2	B	47	A
2	B	48	U
2	B	49	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	46	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	1101	-	3,3,3	0.43	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	1101	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1037/1055 (98%)	0.10	32 (3%) 49 52	15, 38, 73, 109	0
2	B	49/56 (87%)	-0.41	2 (4%) 37 40	19, 35, 77, 132	0
3	D	11/13 (84%)	-0.12	1 (9%) 9 9	26, 30, 64, 89	0
4	C	35/35 (100%)	0.06	4 (11%) 5 4	24, 35, 117, 120	0
5	E	5/6 (83%)	-0.03	0 100 100	35, 35, 42, 76	0
All	All	1137/1165 (97%)	0.07	39 (3%) 45 48	15, 37, 76, 132	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	49	U	5.7
1	A	328	ASN	4.1
1	A	366	LEU	3.9
1	A	371	GLU	3.6
1	A	380	ALA	3.5
1	A	375	ALA	3.4
1	A	329	ASN	3.3
1	A	376	ASP	3.2
4	C	1	DA	3.0
1	A	327	LEU	3.0
1	A	1034	GLY	2.9
1	A	966	TRP	2.9
1	A	405	PHE	2.9
1	A	963	ILE	2.9
1	A	379	ASN	2.9
1	A	953	LEU	2.8
4	C	2	DA	2.8
1	A	193	ALA	2.7
1	A	506	VAL	2.5
3	D	11	DT	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	C	5	DA	2.5
1	A	679	ASN	2.4
1	A	923	ALA	2.4
1	A	411	CYS	2.3
1	A	505	PRO	2.3
1	A	368	LYS	2.3
1	A	1033	ILE	2.3
4	C	3	DT	2.2
1	A	685	VAL	2.2
1	A	409	GLN	2.2
1	A	387	ASP	2.2
1	A	925	ASN	2.2
1	A	365	LYS	2.1
1	A	682	LYS	2.1
1	A	951	GLU	2.1
2	B	48	U	2.1
1	A	373	ASP	2.1
1	A	678	LYS	2.0
1	A	976	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	MG	A	1102	1/1	0.95	0.14	15,15,15,15	0
6	EDO	A	1101	4/4	0.98	0.18	26,26,27,28	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.