

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 10, 2023 – 04:16 AM EDT

PDB ID : 7LVX

Title : The internal aldimine form of the wild-type Salmonella Typhimurium Trypto-

phan Synthase in complex with N-(4'-trifluoromethoxybenzenesulfonyl)-2-a mino-1-ethylphosphate (F9F) at the enzyme alpha-site and cesium ion at the

metal coordination site at 1.55 Angstrom resolution

Authors: Hilario, E.; Dunn, M.F.; Mueller, L.J.

Deposited on : 2021-02-26

Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$ 

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1




## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	268	91%	8%
2	В	397	92%	7% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	304	-	-	X	-
5	CL	A	306	-	-	X	-



## 2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 5991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Tryptophan synthase alpha chain.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	267	Total	С	N	О	S	0	6	0
_			2041	1293	353	387	8			

• Molecule 2 is a protein called Tryptophan synthase beta chain.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	В	395	Total	С	N	О	S	0	12	0
_	D	300	3092	1939	543	589	21		1-	


• Molecule 3 is 2-({[4-(TRIFLUOROMETHOXY)PHENYL]SULFONYL}AMINO)ETHYL DIHYDROGEN PHOSPHATE (three-letter code: F9F) (formula: C<sub>9</sub>H<sub>11</sub>F<sub>3</sub>NO<sub>7</sub>PS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	$\mathbf{Atoms}$					ZeroOcc	AltConf		
3	A	1	Total 22			N 1			S 1	0	0



• Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	В	1	Total C O 4 2 2	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


I	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	5	A	3	Total Cl 3 3	0	0
	5	В	5	Total Cl 5 5	0	0

• Molecule 6 is CESIUM ION (three-letter code: CS) (formula: Cs) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total Cs 4 4	0	1
6	В	3	Total Cs 5 5	0	1




• Molecule 7 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula:  $C_8H_{10}NO_6P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Atoms				ZeroOcc	AltConf
7	D	1	Total	С	N	О	Р	0	0
1	Б	1	15	8	1	5	1	0	U

 $\bullet$  Molecule 8 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula:  $C_2H_6OS$ ).



Ι	Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
	8	В	1	Total	C	0	S	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	Total C O S 4 2 1 1	0	0
8	В	1	Total C O S 4 2 1 1	0	0

## • Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	301	Total O 308 308	0	7
9	В	453	Total O 468 468	0	15



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan synthase alpha chain





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	182.10Å 59.09Å 67.16Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $94.47^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	29.54 - 1.55	Depositor
Resolution (A)	29.54 - 1.55	EDS
% Data completeness	94.4 (29.54-1.55)	Depositor
(in resolution range)	94.4 (29.54-1.55)	EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	0.03	Depositor
$< I/\sigma(I) > 1$	5.46 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
P. P.	0.143 , 0.189	Depositor
$R, R_{free}$	0.144 , 0.189	DCC
$R_{free}$ test set	4962 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	8.8	Xtriage
Anisotropy	1.457	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.32, 59.1	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5991	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, F9F, PLP, CL, CS, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.40	0/2080	0.70	1/2824 (0.0%)	
2	В	0.45	0/3150	0.73	0/4250	
All	All	0.43	0/5230	0.72	1/7074 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	212	PHE	N-CA-CB	5.03	119.66	110.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	296[A]	GLU	Mainchain
2	В	296[B]	GLU	Mainchain

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2041	0	2021	29	0
2	В	3092	0	3056	28	0
3	A	22	0	9	2	0
4	A	12	0	18	7	0
4	В	4	0	6	1	0
5	A	3	0	0	2	0
5	В	5	0	0	0	0
6	A	4	0	0	1	0
6	В	5	0	0	0	0
7	В	15	0	6	1	0
8	В	12	0	18	4	0
9	A	308	0	0	9	0
9	В	468	0	0	15	0
All	All	5991	0	5134	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance } (\text{\AA}) \end{array}$	Clash overlap (Å)	
4:A:304:EDO:H21	9:A:544:HOH:O	1.33	1.29	
4:A:304:EDO:H22	9:A:534[B]:HOH:O	1.48	1.12	
4:A:304:EDO:C1	9:A:534[A]:HOH:O	1.99	1.09	
4:A:304:EDO:H11	9:A:534[A]:HOH:O	1.54	1.03	
2:B:296[B]:GLU:HG2	9:B:516:HOH:O	1.60	1.00	
1:A:190:ALA:HB1	1:A:192:PRO:HD2	1.57	0.86	
1:A:190:ALA:HB3	1:A:193:LEU:HD11	1.59	0.83	
1:A:192:PRO:HD2	1:A:193:LEU:HG	1.57	0.83	
1:A:178[B]:SER:O	1:A:190:ALA:HB2	1.81	0.81	
6:A:309:CS:CS	9:A:568:HOH:O	2.28	0.80	
1:A:190:ALA:CB	1:A:193:LEU:HD11	2.12	0.80	
2:B:217[B]:LEU:O	2:B:217[B]:LEU:HD23	1.88	0.73	
1:A:190:ALA:C	1:A:192:PRO:HD3	2.10	0.72	
2:B:217[A]:LEU:HG	9:B:710:HOH:O	1.89	0.71	
2:B:217[B]:LEU:C	2:B:217[B]:LEU:CD2	2.58	0.71	
1:A:190:ALA:CA	1:A:192:PRO:HD3	2.21	0.71	
2:B:165[B]:THR:CG2	9:B:622:HOH:O	2.41	0.69	
2:B:165[B]:THR:HG22	9:B:622:HOH:O	1.93	0.68	

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance} \; ({ m \AA})$	$-$ overlap $(\AA)$
1:A:190:ALA:HB1	1:A:192:PRO:CD	2.26	0.66
8:B:406:DMS:H23	9:B:913:HOH:O	1.96	0.65
1:A:178[A]:SER:O	1:A:190:ALA:HB2	1.97	0.64
1:A:5:GLU:OE1	5:A:306:CL:CL	2.55	0.61
2:B:217[B]:LEU:O	2:B:217[B]:LEU:CD2	2.48	0.61
8:B:406:DMS:C2	9:B:913:HOH:O	2.49	0.61
1:A:133:VAL:HG12	1:A:152:PHE:CD2	2.36	0.60
2:B:216:ILE:HG23	9:B:710:HOH:O	2.02	0.60
1:A:112:ASP:OD1	1:A:146:HIS:HE1	1.85	0.59
1:A:188:ARG:HH22	1:A:216:SER:HB3	1.68	0.58
4:A:304:EDO:H12	9:A:534[A]:HOH:O	1.85	0.57
2:B:217[B]:LEU:HD23	2:B:217[B]:LEU:C	2.23	0.57
1:A:133:VAL:HG12	1:A:152:PHE:HD2	1.70	0.55
2:B:300:ILE:HD11	2:B:390:ILE:CD1	2.38	0.53
1:A:190:ALA:HB1	1:A:193:LEU:CG	2.38	0.53
2:B:165[B]:THR:HG22	2:B:166:LEU:N	2.19	0.53
2:B:217[B]:LEU:C	2:B:217[B]:LEU:HD22	2.28	0.53
1:A:190:ALA:HA	1:A:192:PRO:HD3	1.90	0.52
2:B:216:ILE:HG21	2:B:224:PRO:HD3	1.89	0.52
2:B:268:GLY:O	2:B:270:PRO:HD2	2.09	0.52
1:A:190:ALA:HB1	1:A:193:LEU:HG	1.91	0.52
1:A:175:TYR:CE2	3:A:301:F9F:H5	2.45	0.52
2:B:87:LYS:NZ	7:B:401:PLP:O3	2.43	0.52
2:B:44[A]:GLN:HG3	9:B:688:HOH:O	2.09	0.51
2:B:296[B]:GLU:CG	9:B:516:HOH:O	2.37	0.51
2:B:223:LEU:HD23	9:B:710:HOH:O	2.10	0.51
1:A:178[B]:SER:HB3	1:A:190:ALA:H	1.77	0.50
1:A:195:HIS:HB3	9:A:424:HOH:O	2.12	0.50
2:B:115:HIS:CE1	2:B:189:GLY:HA2	2.47	0.49
2:B:216:ILE:CG2	9:B:710:HOH:O	2.61	0.47
1:A:190:ALA:CB	1:A:193:LEU:CD1	2.91	0.47
1:A:178[A]:SER:HB2	1:A:190:ALA:H	1.81	0.46
1:A:15:ARG:O	1:A:268:ALA:HB2	2.16	0.45
4:A:304:EDO:O1	5:A:306:CL:CL	2.53	0.45
1:A:22:PHE:HA	1:A:49:GLU:O	2.17	0.43
1:A:156:PRO:HA	1:A:196:LEU:HD11	2.00	0.43
2:B:360[A]:LYS:HG3	9:B:674:HOH:O	2.17	0.43
1:A:153:ILE:HD13	1:A:175:TYR:HB3	1.99	0.43
2:B:193:GLY:HA2	2:B:280:PHE:O	2.19	0.43
2:B:303:GLY:HA3	2:B:350:GLU:OE2	2.18	0.43
3:A:301:F9F:H3	9:A:521:HOH:O	2.19	0.43

Continued on next page...



I 'omtamalod	trom	mmonia	maaa
Continued	11 0116	DICUIUUS	Daue
	.,	10	1

Atom-1	Atom-2	$egin{array}{c}  ext{Interatomic} \  ext{distance } ( ext{Å}) \end{array}$	Clash overlap (Å)
1:A:100:LEU:C	1:A:100:LEU:HD13	2.39	0.42
1:A:188:ARG:HH22	1:A:216:SER:CB	2.32	0.42
8:B:407:DMS:H12	9:B:878:HOH:O	2.20	0.41
2:B:195:HIS:ND1	2:B:196:PRO:HA	2.36	0.41
4:A:304:EDO:C2	9:A:534[B]:HOH:O	2.29	0.41
2:B:65:ILE:HA	4:B:404:EDO:H12	2.03	0.41
1:A:156:PRO:HA	1:A:196:LEU:CD1	2.51	0.41
2:B:302:ALA:HB2	9:B:877:HOH:O	2.21	0.41
8:B:405:DMS:H11	9:B:583:HOH:O	2.21	0.40
2:B:70:ARG:HD3	2:B:367[A]:GLU:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	A	270/268 (101%)	267 (99%)	2 (1%)	1 (0%)	34	14
2	В	$405/397 \; (102\%)$	397 (98%)	8 (2%)	0	100	100
All	All	675/665 (102%)	664 (98%)	10 (2%)	1 (0%)	47	26

#### All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	PHE

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	A	209/208 (100%)	209 (100%)	0	100	100
2	В	321/311 (103%)	319 (99%)	2 (1%)	86	73
All	All	530/519 (102%)	528 (100%)	2 (0%)	91	82

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	В	207	MET
2	В	297	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	HIS
2	В	63	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 17 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	В	404	-	3,3,3	0.09	0	2,2,2	0.79	0
4	EDO	A	302	-	3,3,3	0.13	0	2,2,2	0.23	0
8	DMS	В	406	-	3,3,3	0.20	0	3,3,3	0.09	0
4	EDO	A	303	-	3,3,3	0.12	0	2,2,2	0.30	0
4	EDO	A	304	-	3,3,3	0.14	0	2,2,2	0.74	0
7	PLP	В	401	2	15,15,16	0.92	1 (6%)	20,22,23	1.11	1 (5%)
8	DMS	В	407	-	3,3,3	0.22	0	3,3,3	0.34	0
8	DMS	В	405	-	3,3,3	0.48	0	3,3,3	0.30	0
3	F9F	A	301	_	22,22,22	0.45	0	32,33,33	0.86	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	В	404	-	-	1/1/1/1	-
4	EDO	A	302	-	-	1/1/1/1	-
4	EDO	A	303	-	-	0/1/1/1	-
4	EDO	A	304	-	-	1/1/1/1	-
7	PLP	В	401	2	-	0/6/6/8	0/1/1/1
3	F9F	A	301	-	-	6/20/20/20	0/1/1/1

#### All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$Ideal(\AA)$
7	В	401	PLP	C4A-C4	-2.49	1.46	1.51

#### All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
7	В	401	PLP	O4P-C5A-C5	2.53	114.17	109.35
3	A	301	F9F	C14-N13-S12	-2.00	113.95	120.41

There are no chirality outliers.

All (9) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	A	302	EDO	O1-C1-C2-O2
4	В	404	EDO	O1-C1-C2-O2
3	A	301	F9F	C3-C4-S12-O21
4	A	304	EDO	O1-C1-C2-O2
3	A	301	F9F	C5-C4-S12-O21
3	A	301	F9F	N13-C14-C15-O16
3	A	301	F9F	C6-C1-O7-C8
3	A	301	F9F	C2-C1-O7-C8
3	A	301	F9F	C3-C4-S12-N13


There are no ring outliers.

7 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	404	EDO	1	0
8	В	406	DMS	2	0
4	A	304	EDO	7	0
7	В	401	PLP	1	0
8	В	407	DMS	1	0
8	В	405	DMS	1	0
3	A	301	F9F	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





# 5.7 Other polymers (i)

There are no such residues in this entry.

Torsions

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



Rings

# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	267/268 (99%)	-0.06	9 (3%) 45 52	10, 15, 29, 41	0
2	В	395/397~(99%)	-0.13	10 (2%) 57 64	7, 11, 24, 59	0
All	All	662/665 (99%)	-0.10	19 (2%) 51 59	7, 13, 28, 59	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	GLY	4.5
2	В	395	GLY	4.1
1	A	185	ALA	3.8
2	В	385	PHE	3.5
1	A	268	ALA	3.5
2	В	139	VAL	3.3
1	A	190	ALA	3.1
1	A	192	PRO	3.0
2	В	159	VAL	2.9
1	A	194	HIS	2.6
1	A	195	HIS	2.5
1	A	188	ARG	2.5
2	В	112	ALA	2.4
1	A	186	GLU	2.3
2	В	141	ARG	2.2
2	В	157	ILE	2.0
2	В	393	ALA	2.0
2	В	140	GLU	2.0
2	В	136	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-}factors}({f \AA}^2)$	Q<0.9
5	CL	В	408	1/1	0.82	0.10	64,64,64,64	0
8	DMS	В	405	4/4	0.88	0.15	30,31,37,44	0
4	EDO	В	404	4/4	0.89	0.14	20,25,29,33	0
8	DMS	В	406	4/4	0.93	0.29	39,52,54,59	0
4	EDO	A	303	4/4	0.94	0.22	29,38,40,41	0
4	EDO	A	304	4/4	0.95	0.17	24,27,28,29	0
5	CL	В	410	1/1	0.95	0.06	66,66,66,66	0
4	EDO	A	302	4/4	0.96	0.09	30,32,33,36	0
3	F9F	A	301	22/22	0.96	0.09	13,15,18,19	0
8	DMS	В	407	4/4	0.97	0.14	25,31,32,36	0
7	PLP	В	401	15/16	0.98	0.10	8,10,14,15	0
5	CL	A	305	1/1	0.98	0.04	36,36,36,36	0
5	CL	A	307	1/1	0.98	0.05	38,38,38,38	0
5	CL	В	411	1/1	0.98	0.16	56,56,56,56	0
6	CS	A	310[B]	1/1	0.99	0.21	35,35,35,35	1
6	CS	В	403[A]	1/1	0.99	0.10	26,26,26,26	1
6	CS	В	403[B]	1/1	0.99	0.10	24,24,24,24	1
6	CS	В	403[C]	1/1	0.99	0.10	40,40,40,40	1
5	CL	A	306	1/1	0.99	0.05	37,37,37,37	0
5	CL	В	409	1/1	0.99	0.08	39,39,39,39	0
6	CS	A	309	1/1	0.99	0.02	31,31,31,31	1
6	CS	A	310[A]	1/1	0.99	0.21	34,34,34,34	1
6	CS	A	308	1/1	1.00	0.03	11,11,11,11	1
5	CL	В	412	1/1	1.00	0.03	19,19,19,19	0
6	CS	В	402	1/1	1.00	0.01	14,14,14,14	0
6	CS	В	413	1/1	1.00	0.03	21,21,21,21	1


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




## Electron density around F9F A 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around PLP B 401:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












## Electron density around CS A 310 (B):












## Electron density around CS B 403 (A):












## Electron density around CS B 403 (B):












## Electron density around CS B 403 (C):

 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












## Electron density around CS A 309:

 $2mF_o$ -DF<sub>c</sub> (at 0.7 rmsd) in gray  $mF_o$ -DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)












## Electron density around CS A 310 (A):












## Electron density around CS A 308:












# Electron density around CS B 402:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











# 6.5 Other polymers (i)

There are no such residues in this entry.

