



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2023 – 06:32 PM JST

PDB ID : 6LWO  
Title : Crystal structure of human NEIL1(R242, Y244H) bound to duplex DNA containing 2'-fluoro-2'-deoxy-5,6-dihydrouridine  
Authors : Liu, M.H.; Zhang, J.; Zhu, C.X.; Zhang, X.X.; Gao, Y.Q.; Yi, C.Q.  
Deposited on : 2020-02-07  
Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	 3% (Poor fit), 82% (0-3 outliers), 6% (1 outlier), 10% (2+ outliers or not modelled)
1	D	295	 % (Poor fit), 81% (0-3 outliers), 5% (1 outlier), 12% (2+ outliers or not modelled)
1	G	295	 9% (Poor fit), 69% (0-3 outliers), 5% (1 outlier), 24% (2+ outliers or not modelled)
2	B	13	 15% (Poor fit), 85% (0-3 outliers), 8% (1 outlier), 8% (2+ outliers or not modelled)
2	E	13	 15% (Poor fit), 92% (0-3 outliers), 8% (1 outlier or 2+ outliers)
2	H	13	 23% (Poor fit), 85% (0-3 outliers), 8% (1 outlier), 8% (2+ outliers or not modelled)

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	13	
3	F	13	
3	I	13	

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endonuclease 8-like 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	265	Total 2111	C 1347	N 387	O 367	S 10	0	0	0
1	D	260	Total 2045	C 1305	N 374	O 356	S 10	0	0	0
1	G	223	Total 1574	C 984	N 297	O 286	S 7	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	242	ARG	LYS	variant	UNP Q96FI4
A	244	HIS	TYR	engineered mutation	UNP Q96FI4
D	242	ARG	LYS	variant	UNP Q96FI4
D	244	HIS	TYR	engineered mutation	UNP Q96FI4
G	242	ARG	LYS	variant	UNP Q96FI4
G	244	HIS	TYR	engineered mutation	UNP Q96FI4


- Molecule 2 is a DNA chain called DNA (5'-D(\*CP\*GP\*TP\*CP\*CP\*AP\*(FDU)P\*GP\*TP\*CP\*TP\*AP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	F	N	O				P
2	B	13	Total 258	C 124	F 1	N 43	O 78	P 12	0	0	0
2	E	13	Total 258	C 124	F 1	N 43	O 78	P 12	0	0	0
2	H	13	Total 258	C 124	F 1	N 43	O 78	P 12	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(\*TP\*AP\*GP\*AP\*CP\*CP\*TP\*GP\*GP\*AP\*CP\*GP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	13	Total 267	C 127	N 53	O 75	P 12	0	0	0
3	F	13	Total 267	C 127	N 53	O 75	P 12	0	0	0
3	I	13	Total 267	C 127	N 53	O 75	P 12	0	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	D	1	Total 6	C 3	O 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	133	Total 133 O 133	0	0
5	B	16	Total 16 O 16	0	0
5	C	9	Total 9 O 9	0	0
5	D	91	Total 91 O 91	0	0
5	E	12	Total 12 O 12	0	0

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
5	F	11	Total O 11 11	0	0
5	G	3	Total O 3 3	0	0
5	I	2	Total O 2 2	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Endonuclease 8-like 1




- Molecule 1: Endonuclease 8-like 1




- Molecule 1: Endonuclease 8-like 1





- Molecule 2: DNA (5'-D(\*CP\*GP\*TP\*CP\*CP\*AP\*(FDU)P\*GP\*TP\*CP\*TP\*AP\*C)-3')








- Molecule 2: DNA (5'-D(\*CP\*GP\*TP\*CP\*CP\*AP\*(FDU)P\*GP\*TP\*CP\*TP\*AP\*C)-3')





- Molecule 2: DNA (5'-D(\*CP\*GP\*TP\*CP\*CP\*AP\*(FDU)P\*GP\*TP\*CP\*TP\*AP\*C)-3')





- Molecule 3: DNA (5'-D(\*TP\*AP\*GP\*AP\*CP\*CP\*TP\*GP\*GP\*AP\*CP\*GP\*G)-3')



- Molecule 3: DNA (5'-D(\*TP\*AP\*GP\*AP\*CP\*CP\*TP\*GP\*GP\*AP\*CP\*GP\*G)-3')



- Molecule 3: DNA (5'-D(\*TP\*AP\*GP\*AP\*CP\*CP\*TP\*GP\*GP\*AP\*CP\*GP\*G)-3')





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.93Å 109.45Å 170.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.15 – 2.51 36.12 – 2.51	Depositor EDS
% Data completeness (in resolution range)	98.2 (36.15-2.51) 98.3 (36.12-2.51)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.203 , 0.243 0.204 , 0.244	Depositor DCC
$R_{free}$ test set	2466 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.3	Xtrriage
Anisotropy	0.391	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 57.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7588	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EW3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	2/2169 (0.1%)	0.87	2/2933 (0.1%)
1	D	0.71	0/2099	0.88	2/2837 (0.1%)
1	G	0.70	0/1605	0.81	2/2144 (0.1%)
2	B	0.51	0/264	0.84	0/402
2	E	0.47	0/264	0.82	0/402
2	H	0.41	0/264	0.73	0/402
3	C	0.53	0/300	0.85	1/462 (0.2%)
3	F	0.50	0/300	0.82	0/462
3	I	0.43	0/300	0.88	0/462
All	All	0.68	2/7565 (0.0%)	0.85	7/10506 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	92	GLU	CD-OE2	7.62	1.34	1.25
1	A	92	GLU	CD-OE1	-5.90	1.19	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	105	PRO	C-N-CD	-13.55	90.79	120.60
1	A	133	ARG	NE-CZ-NH2	-6.14	117.23	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	284	ASP	CB-CA-C	6.00	122.40	110.40
1	D	133	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	A	133	ARG	NE-CZ-NH1	5.33	122.97	120.30
3	C	7	DT	O5'-P-OP2	-5.22	101.00	105.70
1	G	185	ARG	CG-CD-NE	-5.07	101.15	111.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	105	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2111	0	2085	11	0
1	D	2045	0	2006	15	0
1	G	1574	0	1361	6	0
2	B	258	0	137	3	0
2	E	258	0	137	1	0
2	H	258	0	137	3	0
3	C	267	0	147	2	0
3	F	267	0	147	1	0
3	I	267	0	147	1	0
4	D	6	0	8	0	0
5	A	133	0	0	3	0
5	B	16	0	0	0	0
5	C	9	0	0	1	0
5	D	91	0	0	0	0
5	E	12	0	0	0	0
5	F	11	0	0	0	0
5	G	3	0	0	0	0
5	I	2	0	0	0	0
All	All	7588	0	6312	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:104:ALA:HB1	1:D:105:PRO:HD2	1.44	0.98
1:D:105:PRO:CB	1:D:106:PRO:HD2	1.91	0.98
1:D:105:PRO:HB3	1:D:106:PRO:HD2	1.55	0.87
1:A:41:GLU:O	1:A:70:GLN:NE2	2.16	0.79
1:G:76:VAL:HG23	1:G:127:GLY:HA2	1.65	0.78
1:G:2:PRO:N	2:H:7:EW3:F2	2.11	0.74
1:D:104:ALA:CB	1:D:105:PRO:HD2	2.16	0.72
3:C:3:DG:N7	5:C:101:HOH:O	2.30	0.65
1:D:104:ALA:HB1	1:D:105:PRO:CD	2.27	0.62
1:A:2:PRO:N	2:B:7:EW3:F2	2.23	0.61
1:D:2:PRO:N	2:E:7:EW3:F2	2.26	0.59
1:A:126:GLY:HA3	5:A:396:HOH:O	2.02	0.57
1:D:130:GLN:HE21	1:D:133:ARG:HD2	1.70	0.57
2:B:13:DC:H2''	2:H:1:DC:C6	2.39	0.57
3:C:1:DT:H2'	3:C:2:DA:C8	2.44	0.53
1:A:242:ARG:HB3	5:A:382:HOH:O	2.09	0.52
1:G:123:TRP:C	1:G:124:ASP:OD1	2.49	0.52
1:A:201:LEU:HD12	1:A:201:LEU:C	2.32	0.50
3:F:1:DT:H2'	3:F:2:DA:C8	2.48	0.48
1:D:155:LYS:H	1:D:155:LYS:HD2	1.79	0.48
3:I:1:DT:H2'	3:I:2:DA:C8	2.49	0.47
1:A:130:GLN:HE21	1:A:133:ARG:HD2	1.79	0.47
1:D:130:GLN:NE2	1:D:133:ARG:HD2	2.28	0.47
1:D:155:LYS:H	1:D:155:LYS:CD	2.27	0.47
1:A:130:GLN:NE2	1:A:133:ARG:HD2	2.32	0.45
1:G:142:GLN:O	1:G:142:GLN:NE2	2.51	0.44
1:G:76:VAL:HG21	1:G:128:LYS:H	1.82	0.44
1:A:106:PRO:CD	1:A:106:PRO:O	2.64	0.44
1:A:130:GLN:HB3	1:A:133:ARG:CD	2.49	0.43
1:D:201:LEU:CD1	1:D:201:LEU:N	2.82	0.42
1:A:90:ARG:NH1	5:A:305:HOH:O	2.47	0.42
1:D:155:LYS:HD2	1:D:155:LYS:N	2.35	0.42
1:A:155:LYS:HD3	1:A:155:LYS:H	1.85	0.42
1:D:130:GLN:NE2	1:D:133:ARG:HH11	2.18	0.42
1:G:130:GLN:HE21	1:G:133:ARG:HE	1.67	0.41
2:B:13:DC:C2'	2:H:1:DC:C6	3.03	0.41
1:D:146:GLU:OE1	1:D:146:GLU:HA	2.19	0.41
1:D:107:GLY:O	1:D:108:PRO:C	2.59	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/295 (88%)	248 (96%)	10 (4%)	1 (0%)	34	54
1	D	254/295 (86%)	240 (94%)	12 (5%)	2 (1%)	19	35
1	G	207/295 (70%)	198 (96%)	9 (4%)	0	100	100
All	All	720/885 (81%)	686 (95%)	31 (4%)	3 (0%)	34	54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	106	PRO
1	A	106	PRO
1	D	105	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/248 (89%)	211 (96%)	10 (4%)	27	51
1	D	212/248 (86%)	201 (95%)	11 (5%)	23	44
1	G	138/248 (56%)	129 (94%)	9 (6%)	17	33
All	All	571/744 (77%)	541 (95%)	30 (5%)	22	43

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	ARG
1	A	155	LYS
1	A	177	TYR
1	A	202	GLN
1	A	222	ASN
1	A	238	GLN
1	A	251	GLU
1	A	252	ASP
1	A	257	ARG
1	A	274	ARG
1	D	11	SER
1	D	95	ARG
1	D	100	ARG
1	D	155	LYS
1	D	177	TYR
1	D	201	LEU
1	D	202	GLN
1	D	222	ASN
1	D	238	GLN
1	D	257	ARG
1	D	265	MET
1	G	31	SER
1	G	84	SER
1	G	99	LEU
1	G	142	GLN
1	G	177	TYR
1	G	201	LEU
1	G	238	GLN
1	G	265	MET
1	G	284	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	130	GLN
1	A	139	GLN
1	A	222	ASN
1	D	69	GLN
1	D	130	GLN
1	D	202	GLN
1	D	222	ASN
1	G	130	GLN

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	G	142	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	EW3	E	7	2	18,21,22	1.20	2 (11%)	22,30,33	1.74	4 (18%)
2	EW3	B	7	2	18,21,22	1.35	2 (11%)	22,30,33	1.48	2 (9%)
2	EW3	H	7	2	18,21,22	1.01	0	22,30,33	1.66	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EW3	E	7	2	-	1/7/38/39	0/2/2/2
2	EW3	B	7	2	-	1/7/38/39	0/2/2/2
2	EW3	H	7	2	-	2/7/38/39	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	7	EW3	F2-C2'	-2.77	1.34	1.40
2	B	7	EW3	C2-N3	-2.61	1.33	1.38
2	E	7	EW3	F2-C2'	-2.51	1.34	1.40
2	E	7	EW3	C4-N3	-2.41	1.33	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	7	EW3	C4-N3-C2	-4.97	121.67	125.79
2	H	7	EW3	C4-N3-C2	-4.86	121.76	125.79
2	B	7	EW3	C4-N3-C2	-3.97	122.50	125.79
2	B	7	EW3	C5-C6-N1	-2.91	102.04	111.61
2	E	7	EW3	C5-C4-N3	2.75	119.74	116.65
2	H	7	EW3	C5-C6-N1	-2.59	103.07	111.61
2	E	7	EW3	C5-C6-N1	-2.58	103.11	111.61
2	H	7	EW3	C5-C4-N3	2.56	119.53	116.65
2	E	7	EW3	F2-C2'-C1'	2.19	113.64	109.08

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	7	EW3	C2'-C1'-N1-C2
2	E	7	EW3	O4'-C4'-C5'-O5'
2	B	7	EW3	O4'-C4'-C5'-O5'
2	H	7	EW3	O4'-C4'-C5'-O5'

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	7	EW3	1	0
2	B	7	EW3	1	0
2	H	7	EW3	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	D	301	-	5,5,5	0.18	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	301	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	301	GOL	O1-C1-C2-C3
4	D	301	GOL	O1-C1-C2-O2
4	D	301	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	265/295 (89%)	0.21	8 (3%) 50 53	34, 58, 97, 134	0
1	D	260/295 (88%)	0.40	2 (0%) 86 87	37, 68, 114, 126	0
1	G	223/295 (75%)	0.83	26 (11%) 4 4	82, 109, 136, 148	0
2	B	12/13 (92%)	0.76	2 (16%) 1 1	60, 97, 137, 145	0
2	E	12/13 (92%)	0.66	2 (16%) 1 1	56, 88, 108, 134	0
2	H	12/13 (92%)	1.11	3 (25%) 0 0	97, 115, 132, 138	0
3	C	13/13 (100%)	0.31	2 (15%) 2 1	70, 94, 122, 134	0
3	F	13/13 (100%)	0.66	3 (23%) 0 0	55, 83, 126, 139	0
3	I	13/13 (100%)	1.41	5 (38%) 0 0	85, 117, 134, 137	0
All	All	823/963 (85%)	0.49	53 (6%) 19 20	34, 80, 130, 148	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	60	LEU	5.4
2	E	1	DC	4.8
1	G	281	PHE	4.3
3	I	12	DG	4.0
1	G	269	SER	3.9
1	A	251	GLU	3.8
1	G	280	TRP	3.7
1	A	243	GLY	3.7
1	G	86	GLN	3.6
1	G	268	MET	3.6
3	I	13	DG	3.6
2	H	13	DC	3.6
1	G	141	TYR	3.5
1	A	244	HIS	3.4
2	B	13	DC	3.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
3	I	11	DC	3.1
1	G	125	LEU	3.0
1	G	267	GLY	3.0
1	D	41	GLU	2.9
3	F	11	DC	2.9
2	H	2	DG	2.9
3	I	1	DT	2.9
1	G	92	GLU	2.9
3	F	12	DG	2.8
1	G	89	PRO	2.8
1	G	91	GLU	2.7
1	G	76	VAL	2.6
1	A	250	GLU	2.5
1	G	164	ALA	2.5
1	A	274	ARG	2.5
1	D	184	TYR	2.5
1	G	101	PHE	2.5
3	F	13	DG	2.5
3	C	1	DT	2.4
2	E	13	DC	2.4
1	A	242	ARG	2.4
3	I	2	DA	2.4
2	H	1	DC	2.4
3	C	13	DG	2.3
1	G	266	PRO	2.3
1	G	100	ARG	2.3
1	G	42	SER	2.3
1	G	286	GLY	2.3
2	B	1	DC	2.3
1	A	252	ASP	2.3
1	G	190	PRO	2.3
1	G	283	GLY	2.3
1	G	79	PHE	2.2
1	G	270	SER	2.2
1	G	102	TYR	2.2
1	G	127	GLY	2.1
1	G	77	PHE	2.1
1	A	2	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EW3	H	7	20/21	0.93	0.16	98,108,120,123	0
2	EW3	E	7	20/21	0.96	0.15	65,77,99,108	0
2	EW3	B	7	20/21	0.97	0.18	63,76,100,106	0

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	D	301	6/6	0.69	0.34	61,78,80,82	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.