



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2023 – 06:33 PM JST

PDB ID : 6LZY  
Title : Structure of Phytolacca americana UGT3 with 18-crown-6  
Authors : Maharjan, R.; Fukuda, Y.; Nakayama, T.; Nakayama, T.; Hamada, H.; Ozaki, S.; Inoue, T.  
Deposited on : 2020-02-19  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	
1	B	505	

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7434 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	454	3642	2349	624	652	17	0	0	0
1	B	453	3636	2346	623	650	17	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP B5MGN9
A	-18	ASN	-	expression tag	UNP B5MGN9
A	-17	HIS	-	expression tag	UNP B5MGN9
A	-16	LYS	-	expression tag	UNP B5MGN9
A	-15	VAL	-	expression tag	UNP B5MGN9
A	-14	HIS	-	expression tag	UNP B5MGN9
A	-13	HIS	-	expression tag	UNP B5MGN9
A	-12	HIS	-	expression tag	UNP B5MGN9
A	-11	HIS	-	expression tag	UNP B5MGN9
A	-10	HIS	-	expression tag	UNP B5MGN9
A	-9	HIS	-	expression tag	UNP B5MGN9
A	-8	LEU	-	expression tag	UNP B5MGN9
A	-7	GLN	-	expression tag	UNP B5MGN9
A	-6	GLU	-	expression tag	UNP B5MGN9
A	-5	ASN	-	expression tag	UNP B5MGN9
A	-4	LEU	-	expression tag	UNP B5MGN9
A	-3	TYR	-	expression tag	UNP B5MGN9
A	-2	PHE	-	expression tag	UNP B5MGN9
A	-1	GLN	-	expression tag	UNP B5MGN9
A	0	GLY	-	expression tag	UNP B5MGN9
B	-19	MET	-	initiating methionine	UNP B5MGN9
B	-18	ASN	-	expression tag	UNP B5MGN9
B	-17	HIS	-	expression tag	UNP B5MGN9
B	-16	LYS	-	expression tag	UNP B5MGN9
B	-15	VAL	-	expression tag	UNP B5MGN9

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP B5MGN9
B	-13	HIS	-	expression tag	UNP B5MGN9
B	-12	HIS	-	expression tag	UNP B5MGN9
B	-11	HIS	-	expression tag	UNP B5MGN9
B	-10	HIS	-	expression tag	UNP B5MGN9
B	-9	HIS	-	expression tag	UNP B5MGN9
B	-8	LEU	-	expression tag	UNP B5MGN9
B	-7	GLN	-	expression tag	UNP B5MGN9
B	-6	GLU	-	expression tag	UNP B5MGN9
B	-5	ASN	-	expression tag	UNP B5MGN9
B	-4	LEU	-	expression tag	UNP B5MGN9
B	-3	TYR	-	expression tag	UNP B5MGN9
B	-2	PHE	-	expression tag	UNP B5MGN9
B	-1	GLN	-	expression tag	UNP B5MGN9
B	0	GLY	-	expression tag	UNP B5MGN9

- Molecule 2 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Br 2 2	0	0
2	B	2	Total Br 2 2	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0

- Molecule 5 is 1,4,7,10,13,16-HEXAOXACYCLOOCTADECANE (three-letter code: O4B) (formula: C<sub>12</sub>H<sub>24</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	C O	0	0
			18	12 6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	43	Total	O	0	0
			43	43		
6	B	38	Total	O	0	0
			38	38		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycosyltransferase



- Molecule 1: Glycosyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.73Å 103.86Å 111.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.22 – 2.40 49.17 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.22-2.40) 99.8 (49.17-2.40)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.241 , 0.279 0.247 , 0.281	Depositor DCC
$R_{free}$ test set	2119 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.3	Xtrriage
Anisotropy	0.083	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 35.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7434	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, BR, EDO, O4B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	1/3731 (0.0%)	0.81	1/5041 (0.0%)
1	B	0.69	0/3725	0.81	2/5033 (0.0%)
All	All	0.70	1/7456 (0.0%)	0.81	3/10074 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	368	VAL	C-O	5.26	1.33	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	177	PRO	N-CA-C	7.45	131.47	112.10
1	B	177	PRO	N-CA-C	7.15	130.68	112.10
1	B	371	CYS	CB-CA-C	-5.19	100.03	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3642	0	3651	22	0
1	B	3636	0	3646	23	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	16	0	24	1	0
3	B	36	0	54	6	0
4	A	1	0	0	0	0
5	B	18	0	24	2	0
6	A	43	0	0	0	0
6	B	38	0	0	2	0
All	All	7434	0	7399	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:ALA:HB1	3:B:511:EDO:H22	1.46	0.97
1:B:116:THR:HG23	1:B:118:PRO:HD3	1.63	0.81
1:B:29:ARG:HG2	1:B:65:ILE:HD11	1.74	0.69
1:B:19:GLY:HA3	3:B:511:EDO:H12	1.80	0.62
1:B:446:GLU:OE2	1:B:449:ARG:NH1	2.33	0.61
1:A:167:VAL:HG13	1:A:172:GLU:HB3	1.83	0.60
1:A:238:GLU:HA	5:B:501:O4B:HAI1	1.84	0.60
1:B:30:LEU:HD11	3:B:510:EDO:H22	1.84	0.59
1:B:182:ASP:OD2	1:B:184:LYS:NZ	2.39	0.56
1:A:90:LEU:HD23	1:A:90:LEU:H	1.72	0.55
1:B:156:GLU:OE1	1:B:156:GLU:HA	2.08	0.53
1:B:18:HIS:HA	1:B:21:MET:HE3	1.91	0.53
1:A:89:ALA:O	1:A:95:ILE:HG13	2.10	0.51
1:A:18:HIS:HA	1:A:21:MET:HE3	1.93	0.51
1:B:19:GLY:HA3	3:B:511:EDO:C1	2.41	0.48
1:B:371:CYS:SG	1:B:388:THR:HB	2.53	0.48
1:B:112:TYR:O	1:B:116:THR:HG22	2.13	0.48
1:A:370:HIS:HA	1:A:389:TRP:O	2.14	0.48
1:B:251:LEU:HD21	1:B:381:THR:HG21	1.95	0.48
1:A:463:VAL:HG12	3:A:606:EDO:H12	1.96	0.47
1:B:370:HIS:HA	1:B:389:TRP:O	2.14	0.47
1:A:147:THR:OG1	1:A:394:GLU:HG3	2.16	0.46
1:B:296:GLN:HA	6:B:627:HOH:O	2.15	0.45
1:A:58:LYS:O	1:A:59:LYS:HG2	2.16	0.45
1:B:394:GLU:OE1	1:B:394:GLU:N	2.43	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:ALA:N	1:A:299:PRO:HD2	2.32	0.45
1:B:155:LEU:HD13	3:B:504:EDO:H21	1.99	0.45
1:B:298:ALA:HB3	1:B:299:PRO:HD3	1.98	0.44
1:B:87:GLU:HB2	6:B:604:HOH:O	2.17	0.43
1:A:240:GLY:HA3	5:B:501:O4B:HAH1	2.00	0.43
1:A:91:GLY:O	1:A:95:ILE:HD12	2.19	0.43
1:A:156:GLU:HA	1:A:156:GLU:OE2	2.18	0.43
1:A:334:LEU:HD23	1:A:334:LEU:HA	1.89	0.43
1:B:147:THR:OG1	1:B:394:GLU:HG3	2.19	0.43
1:A:22:ILE:HB	1:A:23:PRO:HD3	2.01	0.42
1:A:294:ALA:O	1:A:320:ARG:HD3	2.19	0.42
1:A:162:GLU:N	1:A:163:PRO:HD3	2.35	0.42
1:A:285:VAL:HB	1:A:364:ILE:HD13	2.00	0.42
1:A:313:GLN:HG2	1:A:440:MET:HG3	2.00	0.42
1:A:242:ARG:HH21	1:A:479:GLU:CD	2.21	0.42
1:A:119:ASN:O	1:A:140:PRO:HD2	2.20	0.41
1:B:463:VAL:HG22	3:B:510:EDO:H12	2.03	0.41
1:B:252:CYS:SG	1:B:253:ASN:N	2.94	0.41
1:A:251:LEU:HD21	1:A:381:THR:HG21	2.04	0.40
1:B:285:VAL:HB	1:B:364:ILE:HD13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/505 (89%)	436 (97%)	12 (3%)	0	100	100
1	B	447/505 (88%)	435 (97%)	12 (3%)	0	100	100
All	All	895/1010 (89%)	871 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/434 (90%)	377 (97%)	12 (3%)	40	60
1	B	388/434 (89%)	370 (95%)	18 (5%)	27	43
All	All	777/868 (90%)	747 (96%)	30 (4%)	32	50

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	MET
1	A	58	LYS
1	A	60	ASN
1	A	88	GLN
1	A	141	ARG
1	A	177	PRO
1	A	207	LYS
1	A	245	ASN
1	A	392	PHE
1	A	421	THR
1	A	461	LYS
1	A	472	ASP
1	B	16	MET
1	B	60	ASN
1	B	76	ASP
1	B	88	GLN
1	B	141	ARG
1	B	166	ASN
1	B	177	PRO
1	B	185	MET
1	B	193	ASP
1	B	245	ASN
1	B	295	HIS
1	B	320	ARG
1	B	343	GLU
1	B	347	LEU
1	B	392	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	420	GLU
1	B	421	THR
1	B	472	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	ASN
1	A	296	GLN
1	B	7	GLN
1	B	295	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 5 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	B	505	-	3,3,3	0.22	0	2,2,2	0.42	0
3	EDO	B	507	-	3,3,3	0.15	0	2,2,2	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	B	510	-	3,3,3	0.50	0	2,2,2	0.70	0
3	EDO	B	511	-	3,3,3	0.13	0	2,2,2	0.42	0
3	EDO	B	509	-	3,3,3	0.10	0	2,2,2	0.17	0
5	O4B	B	501	4	18,18,18	0.25	0	18,18,18	0.21	0
3	EDO	A	606	-	3,3,3	0.29	0	2,2,2	0.40	0
3	EDO	A	603	-	3,3,3	0.19	0	2,2,2	0.32	0
3	EDO	A	604	-	3,3,3	0.39	0	2,2,2	0.54	0
3	EDO	B	503	-	3,3,3	0.18	0	2,2,2	0.41	0
3	EDO	B	506	-	3,3,3	0.09	0	2,2,2	0.25	0
3	EDO	B	504	-	3,3,3	0.33	0	2,2,2	0.54	0
3	EDO	A	605	-	3,3,3	0.37	0	2,2,2	0.38	0
3	EDO	B	508	-	3,3,3	0.05	0	2,2,2	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	505	-	-	1/1/1/1	-
3	EDO	B	507	-	-	1/1/1/1	-
3	EDO	B	510	-	-	1/1/1/1	-
3	EDO	B	511	-	-	1/1/1/1	-
3	EDO	B	509	-	-	0/1/1/1	-
5	O4B	B	501	4	-	6/18/18/18	0/1/1/1
3	EDO	A	606	-	-	1/1/1/1	-
3	EDO	A	603	-	-	1/1/1/1	-
3	EDO	A	604	-	-	1/1/1/1	-
3	EDO	B	503	-	-	1/1/1/1	-
3	EDO	B	506	-	-	1/1/1/1	-
3	EDO	B	504	-	-	0/1/1/1	-
3	EDO	A	605	-	-	1/1/1/1	-
3	EDO	B	508	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	603	EDO	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	A	604	EDO	O1-C1-C2-O2
3	B	503	EDO	O1-C1-C2-O2
3	B	508	EDO	O1-C1-C2-O2
3	B	507	EDO	O1-C1-C2-O2
3	B	511	EDO	O1-C1-C2-O2
5	B	501	O4B	OAM-CAC-CAD-OAO
5	B	501	O4B	OAQ-CAK-CAL-OAR
3	B	505	EDO	O1-C1-C2-O2
5	B	501	O4B	OAN-CAE-CAF-OAP
5	B	501	O4B	CAA-CAB-OAN-CAE
3	B	510	EDO	O1-C1-C2-O2
3	A	605	EDO	O1-C1-C2-O2
3	A	606	EDO	O1-C1-C2-O2
3	B	506	EDO	O1-C1-C2-O2
5	B	501	O4B	CAI-CAJ-OAR-CAL
5	B	501	O4B	CAL-CAK-OAQ-CAH

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	510	EDO	2	0
3	B	511	EDO	3	0
5	B	501	O4B	2	0
3	A	606	EDO	1	0
3	B	504	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	454/505 (89%)	0.47	34 (7%) <b>14</b> <b>13</b>	21, 40, 88, 133	0
1	B	453/505 (89%)	0.72	47 (10%) <b>6</b> <b>6</b>	21, 45, 104, 146	0
All	All	907/1010 (89%)	0.60	81 (8%) <b>9</b> <b>9</b>	21, 42, 98, 146	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	293	THR	9.1
1	B	57	GLY	9.0
1	B	294	ALA	8.4
1	B	291	GLY	8.2
1	A	197	HIS	6.6
1	B	60	ASN	6.3
1	B	75	GLN	6.0
1	B	197	HIS	6.0
1	A	57	GLY	5.7
1	A	195	TRP	5.6
1	A	92	SER	5.5
1	A	60	ASN	5.4
1	A	253	ASN	5.2
1	A	484	HIS	5.2
1	A	61	GLY	4.8
1	A	200	ALA	4.8
1	A	55	GLU	4.8
1	B	93	SER	4.8
1	B	352	TRP	4.6
1	B	61	GLY	4.5
1	B	200	ALA	4.5
1	B	193	ASP	4.4
1	B	354	PRO	4.3
1	B	199	LYS	4.2

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	193	ASP	4.2
1	A	196	LYS	4.2
1	A	59	LYS	4.1
1	A	295	HIS	3.9
1	B	82	GLY	3.9
1	A	199	LYS	3.8
1	A	273	CYS	3.8
1	A	255	SER	3.8
1	B	292	SER	3.7
1	A	198	GLU	3.6
1	B	6	GLN	3.5
1	B	81	GLU	3.5
1	B	59	LYS	3.3
1	A	64	THR	3.2
1	A	292	SER	3.2
1	A	6	GLN	3.2
1	B	305	ALA	3.1
1	A	347	LEU	3.1
1	B	420	GLU	3.0
1	B	92	SER	3.0
1	A	56	MET	3.0
1	B	62	SER	3.0
1	B	56	MET	2.9
1	B	78	GLY	2.9
1	B	77	VAL	2.9
1	A	166	ASN	2.9
1	B	344	GLY	2.8
1	B	295	HIS	2.8
1	B	55	GLU	2.7
1	B	51	THR	2.7
1	A	58	LYS	2.5
1	A	62	SER	2.5
1	A	192	GLU	2.5
1	B	90	LEU	2.5
1	B	253	ASN	2.5
1	B	58	LYS	2.5
1	B	97	LYS	2.4
1	A	187	ARG	2.4
1	B	353	ALA	2.4
1	A	93	SER	2.4
1	B	484	HIS	2.3
1	A	340	GLN	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	350	ARG	2.3
1	B	74	ALA	2.3
1	B	198	GLU	2.3
1	B	351	GLY	2.3
1	B	83	CYS	2.3
1	B	339	GLU	2.2
1	A	63	PRO	2.2
1	A	274	LEU	2.2
1	A	331	ASP	2.2
1	B	333	TRP	2.1
1	B	53	ALA	2.1
1	B	290	PHE	2.0
1	B	332	SER	2.0
1	B	64	THR	2.0
1	A	420	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	603	4/4	0.65	0.36	69,74,78,78	0
3	EDO	A	606	4/4	0.66	0.31	60,68,68,69	0
3	EDO	A	605	4/4	0.69	0.34	48,56,57,64	0
3	EDO	A	604	4/4	0.74	0.27	46,51,52,54	0
3	EDO	B	507	4/4	0.74	0.26	66,70,71,72	0
3	EDO	B	505	4/4	0.79	0.27	61,63,65,68	0
3	EDO	B	504	4/4	0.79	0.24	51,52,56,58	0
2	BR	B	512	1/1	0.81	0.10	106,106,106,106	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	B	510	4/4	0.82	0.33	59,65,69,73	0
3	EDO	B	509	4/4	0.87	0.13	53,57,58,65	0
3	EDO	B	508	4/4	0.88	0.20	72,72,73,76	0
3	EDO	B	511	4/4	0.88	0.17	57,59,59,63	0
4	NA	A	607	1/1	0.88	0.13	34,34,34,34	0
3	EDO	B	506	4/4	0.89	0.24	68,73,75,78	0
3	EDO	B	503	4/4	0.92	0.23	38,49,49,56	0
5	O4B	B	501	18/18	0.92	0.20	29,31,32,33	0
2	BR	A	602	1/1	0.98	0.06	45,45,45,45	0
2	BR	A	601	1/1	0.99	0.08	34,34,34,34	0
2	BR	B	502	1/1	0.99	0.12	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.