



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 11:40 PM EDT

PDB ID : 7LZR
Title : Crystal structure of the BCL6 BTB domain in complex with OICR-10256
Authors : Kuntz, D.A.; Prive, G.G.
Deposited on : 2021-03-10
Resolution : 1.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1385 (1.36-1.32)
Clashscore	141614	1417 (1.36-1.32)
Ramachandran outliers	138981	1397 (1.36-1.32)
Sidechain outliers	138945	1397 (1.36-1.32)
RSRZ outliers	127900	1369 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	125	 2% 94% 5%
1	B	125	 3% 95%
1	C	125	 2% 90% 7%
1	D	125	 2% 93% 5%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8525 atoms, of which 3975 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called B-cell lymphoma 6 protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	123	1995	634	996	176	181	8	0	3	0
1	B	123	2022	643	1010	179	182	8	0	5	0
1	C	122	1977	632	981	173	183	8	0	4	0
1	D	122	2000	638	988	181	185	8	0	4	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	8	GLN	CYS	engineered mutation	UNP P41182
A	67	ARG	CYS	engineered mutation	UNP P41182
A	84	ASN	CYS	engineered mutation	UNP P41182
B	8	GLN	CYS	engineered mutation	UNP P41182
B	67	ARG	CYS	engineered mutation	UNP P41182
B	84	ASN	CYS	engineered mutation	UNP P41182
C	8	GLN	CYS	engineered mutation	UNP P41182
C	67	ARG	CYS	engineered mutation	UNP P41182
C	84	ASN	CYS	engineered mutation	UNP P41182
D	8	GLN	CYS	engineered mutation	UNP P41182
D	67	ARG	CYS	engineered mutation	UNP P41182
D	84	ASN	CYS	engineered mutation	UNP P41182

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	1
			12	6	6		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	2	Total	Cl	0	0
			2	2		
4	C	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		

- Molecule 5 is N-[5-chloro-2-(morpholin-4-yl)pyridin-4-yl]-2-[5-(3-cyano-4-hydroxy-5-methylphenyl)-3-methyl-4-oxo-3,4-dihydro-7H-pyrrolo[2,3-d]pyrimidin-7-yl]acetamide (three-letter code: YJJ) (formula: C₂₆H₂₄ClN₇O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	A	1	Total	C	Cl	N	O	0	0
			38	26	1	7	4		
5	B	1	Total	C	Cl	N	O	0	0
			38	26	1	7	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	C	1	Total	C	Cl	N	O	0	0
			38	26	1	7	4		
5	D	1	Total	C	Cl	N	O	0	0
			38	26	1	7	4		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	80	Total	O	0	5
			81	81		
6	B	75	Total	O	0	0
			75	75		
6	C	73	Total	O	0	2
			75	75		
6	D	91	Total	O	0	3
			94	94		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: B-cell lymphoma 6 protein




- Molecule 1: B-cell lymphoma 6 protein



- Molecule 1: B-cell lymphoma 6 protein



- Molecule 1: B-cell lymphoma 6 protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.36Å 73.06Å 64.42Å 90.00° 109.50° 90.00°	Depositor
Resolution (Å)	31.30 – 1.34 60.73 – 1.34	Depositor EDS
% Data completeness (in resolution range)	99.5 (31.30-1.34) 99.5 (60.73-1.34)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.34Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.135 , 0.159 0.135 , 0.158	Depositor DCC
R_{free} test set	5349 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtrriage
Anisotropy	0.039	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	8525	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.78 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4764e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL, SO4, YJJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	0/1023	0.85	0/1380
1	B	0.76	0/1039	0.82	0/1400
1	C	0.76	0/1020	0.79	0/1377
1	D	0.76	0/1030	0.82	0/1388
All	All	0.76	0/4112	0.82	0/5545

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	999	996	1025	3	0
1	B	1012	1010	1042	3	0
1	C	996	981	1015	8	0
1	D	1012	988	1028	5	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	10	0	0	1	0
3	A	18	0	24	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	6	0	8	0	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	38	0	0	0	0
5	B	38	0	0	0	0
5	C	38	0	0	0	0
5	D	38	0	0	0	0
6	A	81	0	0	0	0
6	B	75	0	0	0	0
6	C	75	0	0	1	0
6	D	94	0	0	3	0
All	All	4550	3975	4142	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:24:ARG:O	1:B:28:ARG:HD3	1.86	0.75
1:D:110:MET:HG2	6:D:387:HOH:O	2.06	0.55
1:C:11:PHE:HB3	6:C:357:HOH:O	2.06	0.54
1:C:28:ARG:HD2	6:D:345:HOH:O	2.08	0.53
1:D:40:ARG:HG3	6:D:381:HOH:O	2.11	0.51
1:C:8:GLN:NE2	1:C:8:GLN:H	2.09	0.50
1:D:124:PHE:O	1:D:128:SER:HB2	2.12	0.49
1:A:58:TYR:O	1:A:62:THR:HG23	2.13	0.48
1:B:27[B]:SER:OG	1:B:28:ARG:NH2	2.36	0.48
1:C:35[B]:VAL:CG2	1:C:44:ARG:NH1	2.78	0.46
1:C:58:TYR:O	1:C:62:THR:HG23	2.16	0.46
1:C:116:HIS:HE1	2:D:202:SO4:O2	2.00	0.45
1:C:11:PHE:CD2	1:D:116:HIS:CD2	3.07	0.43
1:D:123:LYS:HD2	1:D:123:LYS:N	2.35	0.41
1:A:88:ASP:CG	3:A:202:GOL:H12	2.40	0.41
1:A:86:LEU:HD21	1:A:117:VAL:HG11	2.03	0.41
1:B:28:ARG:CZ	1:C:88[A]:ASP:OD2	2.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/125 (99%)	122 (98%)	2 (2%)	0	100	100
1	B	126/125 (101%)	123 (98%)	3 (2%)	0	100	100
1	C	124/125 (99%)	122 (98%)	2 (2%)	0	100	100
1	D	124/125 (99%)	121 (98%)	3 (2%)	0	100	100
All	All	498/500 (100%)	488 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/114 (101%)	114 (99%)	1 (1%)	78	52
1	B	116/114 (102%)	115 (99%)	1 (1%)	78	52
1	C	115/114 (101%)	113 (98%)	2 (2%)	60	27
1	D	116/114 (102%)	116 (100%)	0	100	100
All	All	462/456 (101%)	458 (99%)	4 (1%)	78	52

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	102	ILE
1	B	84	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	8	GLN
1	C	98	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	8	GLN
1	D	116	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 5 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	YJJ	D	201	-	39,42,42	2.13	11 (28%)	43,61,61	1.37	6 (13%)
2	SO4	A	201	-	4,4,4	0.11	0	6,6,6	0.51	0
2	SO4	D	203	-	4,4,4	0.15	0	6,6,6	0.66	0
3	GOL	A	205[A]	-	5,5,5	0.35	0	5,5,5	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	202	-	5,5,5	0.13	0	5,5,5	1.05	0
2	SO4	D	202	-	4,4,4	0.42	0	6,6,6	0.69	0
5	YJJ	C	203	-	39,42,42	2.19	12 (30%)	43,61,61	1.76	13 (30%)
5	YJJ	B	201	-	39,42,42	2.20	12 (30%)	43,61,61	1.58	10 (23%)
2	SO4	C	201	-	4,4,4	0.11	0	6,6,6	0.34	0
3	GOL	C	202	-	5,5,5	0.64	0	5,5,5	0.51	0
2	SO4	B	202	-	4,4,4	0.29	0	6,6,6	0.25	0
5	YJJ	A	204	-	39,42,42	3.13	15 (38%)	43,61,61	1.62	7 (16%)
3	GOL	A	205[B]	-	5,5,5	0.32	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	YJJ	D	201	-	-	0/18/26/26	0/5/5/5
3	GOL	A	205[B]	-	-	0/4/4/4	-
3	GOL	A	205[A]	-	-	3/4/4/4	-
3	GOL	A	202	-	-	2/4/4/4	-
5	YJJ	C	203	-	-	0/18/26/26	0/5/5/5
5	YJJ	B	201	-	-	1/18/26/26	0/5/5/5
3	GOL	C	202	-	-	0/4/4/4	-
5	YJJ	A	204	-	-	1/18/26/26	0/5/5/5

All (50) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	204	YJJ	C03-N04	11.51	1.43	1.29
5	D	201	YJJ	C03-N04	6.98	1.37	1.29
5	A	204	YJJ	C12-N11	-6.81	1.40	1.47
5	B	201	YJJ	C03-N04	6.66	1.37	1.29
5	C	203	YJJ	C03-N04	6.62	1.37	1.29
5	A	204	YJJ	C35-C37	5.58	1.52	1.44
5	B	201	YJJ	C12-N11	-5.57	1.41	1.47
5	D	201	YJJ	C12-N11	-5.41	1.41	1.47
5	C	203	YJJ	C12-N11	-4.86	1.42	1.47
5	A	204	YJJ	C10-N11	-4.41	1.32	1.38
5	A	204	YJJ	C09-C29	-4.33	1.41	1.49
5	A	204	YJJ	C18-N23	4.24	1.46	1.37
5	B	201	YJJ	C28-N23	-3.91	1.40	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	204	YJJ	C28-N23	-3.72	1.40	1.46
5	A	204	YJJ	O14-C13	-3.67	1.15	1.23
5	C	203	YJJ	C35-C37	3.66	1.49	1.44
5	A	204	YJJ	C24-N23	-3.52	1.41	1.46
5	C	203	YJJ	C28-N23	-3.48	1.41	1.46
5	B	201	YJJ	C35-C37	3.45	1.49	1.44
5	C	203	YJJ	C09-C29	-3.42	1.43	1.49
5	D	201	YJJ	C09-C29	-3.35	1.43	1.49
5	A	204	YJJ	O34-C33	3.33	1.44	1.37
5	C	203	YJJ	C36-C35	-3.22	1.35	1.39
5	D	201	YJJ	C35-C37	3.22	1.49	1.44
5	A	204	YJJ	C06-C05	-3.12	1.36	1.40
5	C	203	YJJ	C18-N23	3.11	1.43	1.37
5	A	204	YJJ	C09-C06	3.08	1.45	1.41
5	B	201	YJJ	C18-N23	3.01	1.43	1.37
5	D	201	YJJ	C18-N23	2.98	1.43	1.37
5	D	201	YJJ	C10-C09	2.93	1.42	1.38
5	C	203	YJJ	O34-C33	2.93	1.43	1.37
5	A	204	YJJ	C36-C35	-2.90	1.35	1.39
5	B	201	YJJ	C06-C05	-2.88	1.37	1.40
5	D	201	YJJ	O14-C13	-2.83	1.17	1.23
5	C	203	YJJ	C09-C06	2.65	1.44	1.41
5	B	201	YJJ	C10-N11	-2.59	1.35	1.38
5	B	201	YJJ	O14-C13	-2.59	1.18	1.23
5	D	201	YJJ	C17-C16	-2.48	1.35	1.39
5	A	204	YJJ	C01-N02	-2.45	1.41	1.46
5	D	201	YJJ	C09-C06	2.42	1.44	1.41
5	B	201	YJJ	C13-N15	2.41	1.41	1.35
5	B	201	YJJ	O34-C33	2.39	1.42	1.37
5	A	204	YJJ	C03-N02	2.37	1.40	1.35
5	B	201	YJJ	C09-C06	2.36	1.44	1.41
5	C	203	YJJ	C24-N23	-2.31	1.43	1.46
5	D	201	YJJ	C36-C35	-2.25	1.36	1.39
5	B	201	YJJ	C09-C29	-2.24	1.45	1.49
5	C	203	YJJ	O14-C13	-2.22	1.18	1.23
5	D	201	YJJ	C13-N15	2.14	1.40	1.35
5	C	203	YJJ	C18-N19	-2.10	1.29	1.34

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	204	YJJ	N02-C03-N04	-5.29	119.86	126.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	201	YJJ	N02-C03-N04	-4.77	120.46	126.02
5	D	201	YJJ	N02-C03-N04	-3.96	121.40	126.02
5	C	203	YJJ	C36-C35-C33	3.96	123.23	119.00
5	C	203	YJJ	C17-C18-N23	-3.69	117.90	122.29
5	D	201	YJJ	C36-C35-C33	3.56	122.80	119.00
5	B	201	YJJ	C28-N23-C24	3.41	119.05	111.52
5	A	204	YJJ	C20-C21-C16	-3.35	118.11	120.06
5	C	203	YJJ	C30-C31-C33	3.21	120.70	118.08
5	C	203	YJJ	C20-C21-C16	-3.13	118.23	120.06
5	B	201	YJJ	C20-C21-C16	-3.09	118.26	120.06
5	C	203	YJJ	C35-C33-C31	-2.97	117.53	120.57
5	A	204	YJJ	C28-N23-C24	2.83	117.76	111.52
5	A	204	YJJ	C16-C21-CL22	2.82	122.75	119.54
5	B	201	YJJ	C03-N02-C07	2.80	125.80	122.11
5	C	203	YJJ	C28-N23-C24	2.79	117.67	111.52
5	A	204	YJJ	C27-C28-N23	2.74	115.07	110.02
5	C	203	YJJ	N02-C03-N04	-2.69	122.89	126.02
5	C	203	YJJ	C16-C21-CL22	2.67	122.58	119.54
5	D	201	YJJ	C29-C36-C35	-2.66	119.14	121.53
5	B	201	YJJ	C16-C21-CL22	2.62	122.52	119.54
5	C	203	YJJ	C24-N23-C18	-2.59	114.37	120.39
5	D	201	YJJ	C35-C33-C31	-2.49	118.02	120.57
5	C	203	YJJ	C29-C30-C31	-2.43	119.45	122.33
5	B	201	YJJ	C24-N23-C18	-2.39	114.83	120.39
5	B	201	YJJ	O26-C27-C28	-2.37	106.57	111.80
5	C	203	YJJ	C36-C35-C37	-2.36	116.18	119.74
5	B	201	YJJ	C17-C18-N23	-2.36	119.49	122.29
5	C	203	YJJ	C03-N02-C07	2.34	125.19	122.11
5	A	204	YJJ	C20-N19-C18	2.20	121.31	117.30
5	B	201	YJJ	C36-C35-C33	2.19	121.33	119.00
5	D	201	YJJ	O14-C13-C12	2.15	123.53	120.04
5	A	204	YJJ	C21-C20-N19	-2.04	121.09	122.84
5	D	201	YJJ	C24-N23-C18	-2.04	115.65	120.39
5	C	203	YJJ	C29-C36-C35	-2.01	119.72	121.53
5	B	201	YJJ	C35-C37-N38	-2.00	174.58	177.88

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	205[A]	GOL	O1-C1-C2-C3
3	A	205[A]	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
3	A	205[A]	GOL	O1-C1-C2-O2
3	A	202	GOL	O1-C1-C2-C3
3	A	202	GOL	O1-C1-C2-O2
5	A	204	YJJ	C33-C35-C37-N38
5	B	201	YJJ	N11-C12-C13-N15


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	202	GOL	1	0
2	D	202	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	123/125 (98%)	-0.29	2 (1%) 72 76	13, 21, 38, 62	0
1	B	123/125 (98%)	-0.18	4 (3%) 46 52	13, 20, 41, 53	0
1	C	122/125 (97%)	-0.21	2 (1%) 72 76	13, 22, 44, 58	0
1	D	122/125 (97%)	-0.20	2 (1%) 72 76	13, 18, 38, 83	0
All	All	490/500 (98%)	-0.22	10 (2%) 65 70	13, 20, 42, 83	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	128	SER	6.9
1	B	7	SER	6.0
1	C	8	GLN	3.8
1	D	127	ALA	3.7
1	A	7	SER	3.6
1	B	40	ARG	3.0
1	A	8	GLN	3.0
1	C	39	SER	2.8
1	B	18	VAL	2.5
1	B	8	GLN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	202	6/6	0.62	0.25	37,39,43,46	6
2	SO4	C	201	5/5	0.74	0.23	37,38,44,48	5
3	GOL	C	202	6/6	0.74	0.14	23,29,32,32	0
3	GOL	A	205[B]	6/6	0.84	0.15	26,29,31,33	6
3	GOL	A	205[A]	6/6	0.84	0.15	25,29,30,33	6
2	SO4	A	201	5/5	0.89	0.15	48,50,52,53	5
2	SO4	D	203	5/5	0.90	0.12	43,49,51,55	5
4	CL	B	204	1/1	0.90	0.08	50,50,50,50	0
4	CL	D	204	1/1	0.90	0.07	63,63,63,63	0
2	SO4	B	202	5/5	0.92	0.12	44,45,47,54	5
2	SO4	D	202	5/5	0.94	0.12	31,33,36,38	0
4	CL	B	203	1/1	0.95	0.03	63,63,63,63	0
4	CL	A	203	1/1	0.96	0.05	59,59,59,59	0
5	YJJ	A	204	38/38	0.98	0.08	14,16,42,43	0
5	YJJ	B	201	38/38	0.98	0.07	13,15,26,28	0
5	YJJ	C	203	38/38	0.98	0.07	13,15,24,26	0
4	CL	C	204	1/1	0.99	0.04	38,38,38,38	0
5	YJJ	D	201	38/38	0.99	0.07	12,14,16,17	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around YJJ A 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YJJ B 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YJJ C 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around YJJ D 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.