



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2023 – 08:41 PM JST

PDB ID : 6M64
Title : Crystal structure of SMAD2 in complex with CBP
Authors : Miyazono, K.; Ito, T.; Wada, H.; Tanokura, M.
Deposited on : 2020-03-13
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	 3% 91% 6% .
1	C	208	 % 87% 6% 7%
1	E	208	 2% 87% 6% 7%
2	B	28	 68% 32%
2	D	28	 7% 61% 11% 29%
2	F	28	 14% 82% 14% .

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 5863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mothers against decapentaplegic homolog 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	203	1622	1024	283	301	14	0	2	0
1	C	194	1548	981	271	284	12	0	1	0
1	E	193	1534	972	269	281	12	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	260	GLY	-	expression tag	UNP Q15796
A	261	PRO	-	expression tag	UNP Q15796
A	465	GLU	-	expression tag	UNP Q15796
A	466	MET	-	expression tag	UNP Q15796
A	467	GLU	-	expression tag	UNP Q15796
C	260	GLY	-	expression tag	UNP Q15796
C	261	PRO	-	expression tag	UNP Q15796
C	465	GLU	-	expression tag	UNP Q15796
C	466	MET	-	expression tag	UNP Q15796
C	467	GLU	-	expression tag	UNP Q15796
E	260	GLY	-	expression tag	UNP Q15796
E	261	PRO	-	expression tag	UNP Q15796
E	465	GLU	-	expression tag	UNP Q15796
E	466	MET	-	expression tag	UNP Q15796
E	467	GLU	-	expression tag	UNP Q15796

- Molecule 2 is a protein called CBP.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	19	140	86	28	26	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	20	Total	C	N	O	0	0	0
			149	91	30	28			
2	F	27	Total	C	N	O	0	0	0
			216	131	40	45			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	202	Total	O	0	0
			202	202		
3	B	24	Total	O	0	0
			24	24		
3	C	192	Total	O	0	0
			192	192		
3	D	23	Total	O	0	0
			23	23		
3	E	178	Total	O	0	0
			178	178		
3	F	35	Total	O	0	0
			35	35		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mothers against decapentaplegic homolog 2




- Molecule 1: Mothers against decapentaplegic homolog 2




- Molecule 1: Mothers against decapentaplegic homolog 2




- Molecule 2: CBP



- Molecule 2: CBP



- Molecule 2: CBP

Chain F:  14% 82% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.37Å 70.80Å 165.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.67 – 1.45 36.67 – 1.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (36.67-1.45) 100.0 (36.67-1.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 1.45Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.169 , 0.189 0.169 , 0.189	Depositor DCC
R_{free} test set	7239 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.5	Xtrriage
Anisotropy	0.374	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.025 for k,h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5863	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/1667	0.84	1/2265 (0.0%)
1	C	0.63	0/1589	0.83	5/2161 (0.2%)
1	E	0.59	0/1574	0.78	1/2140 (0.0%)
2	B	0.64	0/142	0.64	0/193
2	D	0.63	0/151	0.72	0/205
2	F	0.43	0/220	0.80	1/297 (0.3%)
All	All	0.64	0/5343	0.81	8/7261 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	285	ARG	NE-CZ-NH2	6.90	123.75	120.30
1	C	285	ARG	NE-CZ-NH1	-6.66	116.97	120.30
1	E	440	LEU	CA-CB-CG	6.12	129.38	115.30
1	A	285	ARG	NE-CZ-NH2	5.84	123.22	120.30
1	C	456	MET	C-N-CA	-5.34	111.08	122.30
1	C	321	ARG	NE-CZ-NH1	5.31	122.96	120.30
2	F	1975	ASP	CB-CG-OD1	5.31	123.08	118.30
1	C	321	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1622	0	1567	8	0
1	C	1548	0	1494	6	0
1	E	1534	0	1481	9	0
2	B	140	0	141	0	0
2	D	149	0	149	2	0
2	F	216	0	201	1	0
3	A	202	0	0	0	0
3	B	24	0	0	0	0
3	C	192	0	0	0	0
3	D	23	0	0	0	0
3	E	178	0	0	1	0
3	F	35	0	0	0	0
All	All	5863	0	5033	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:357:VAL:HG22	1:E:416:MET:HG2	1.81	0.63
1:A:323:ALA:O	1:A:327:MET:HG2	2.03	0.59
1:E:458:SER:O	1:E:458:SER:OG	2.24	0.55
1:A:336:VAL:HG21	1:A:416:MET:HE1	1.91	0.53
1:C:330:ARG:NH1	1:E:403:GLU:OE1	2.23	0.52
2:F:1957:GLU:OE2	2:F:1960:ARG:NH2	2.41	0.50
1:C:330:ARG:NH2	1:E:403:GLU:OE2	2.32	0.49
1:C:357:VAL:HG22	1:C:416:MET:HG2	1.95	0.47
1:A:280:TYR:CZ	1:A:285:ARG:HD3	2.51	0.46
1:C:388:GLN:OE1	2:D:1967:GLN:HG2	2.17	0.45
2:D:1964:ARG:O	2:D:1968:GLN:HG2	2.16	0.45
1:C:280:TYR:CZ	1:C:285:ARG:HD3	2.51	0.45
1:E:413:THR:HG22	1:E:441:HIS:CD2	2.53	0.44
1:E:439:GLU:HG3	3:E:520:HOH:O	2.17	0.44
1:A:328:THR:HG21	1:A:435:PRO:HG2	1.98	0.43
1:A:467:GLU:HG3	1:E:420:LYS:NZ	2.34	0.43
1:E:280:TYR:CZ	1:E:285:ARG:HD3	2.54	0.42
1:A:467:GLU:HG3	1:E:420:LYS:HZ2	1.84	0.42
1:A:285:ARG:HH11	1:A:285:ARG:HD2	1.58	0.42
1:A:365:ARG:HD2	1:A:366:TYR:CZ	2.56	0.40
1:C:339:TYR:CZ	1:C:346:PHE:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/208 (98%)	198 (98%)	5 (2%)	0	100	100
1	C	193/208 (93%)	189 (98%)	3 (2%)	1 (0%)	29	9
1	E	191/208 (92%)	187 (98%)	4 (2%)	0	100	100
2	B	17/28 (61%)	17 (100%)	0	0	100	100
2	D	18/28 (64%)	18 (100%)	0	0	100	100
2	F	25/28 (89%)	25 (100%)	0	0	100	100
All	All	647/708 (91%)	634 (98%)	12 (2%)	1 (0%)	47	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	424	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/181 (99%)	176 (98%)	3 (2%)	60	28
1	C	169/181 (93%)	167 (99%)	2 (1%)	71	43
1	E	167/181 (92%)	165 (99%)	2 (1%)	71	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	13/22 (59%)	13 (100%)	0	100	100
2	D	14/22 (64%)	14 (100%)	0	100	100
2	F	22/22 (100%)	21 (96%)	1 (4%)	27	3
All	All	564/609 (93%)	556 (99%)	8 (1%)	67	37

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	266	VAL
1	A	425	GLU
1	A	464	SER
1	C	425	GLU
1	C	427	ARG
1	E	267	THR
1	E	297	LEU
2	F	1963	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/208 (97%)	-0.05	7 (3%) 45 48	16, 23, 45, 75	0
1	C	194/208 (93%)	-0.11	3 (1%) 73 74	17, 25, 44, 66	0
1	E	193/208 (92%)	-0.02	5 (2%) 56 58	17, 29, 49, 61	0
2	B	19/28 (67%)	-0.16	0 100 100	18, 22, 46, 59	0
2	D	20/28 (71%)	0.11	2 (10%) 7 8	20, 27, 55, 67	0
2	F	27/28 (96%)	0.61	4 (14%) 2 3	29, 40, 57, 58	0
All	All	656/708 (92%)	-0.03	21 (3%) 47 50	16, 26, 50, 75	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	1951	PRO	5.0
1	A	461	VAL	5.0
2	D	1968	GLN	3.8
1	A	467	GLU	3.4
1	A	460	SER	3.4
1	E	457	GLY	3.4
1	E	266	VAL	3.3
1	A	462	ARG	3.1
2	F	1952	PRO	2.9
1	A	466	MET	2.7
1	A	266	VAL	2.7
2	F	1953	PRO	2.6
1	C	341	ILE	2.5
1	A	425	GLU	2.2
1	C	425	GLU	2.2
2	F	1956	VAL	2.2
1	E	267	THR	2.2
1	E	339	TYR	2.2
1	C	459	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	366	TYR	2.1
2	D	1967	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.