



## Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 08:43 pm BST

PDB ID : 4M8S  
Title : Crystal structure of 3-ketoacyl -(acyl carrier protein) reductase (FabG) from *Neisseria meningitidis*  
Authors : Nanson, J.D.; Forwood, J.K.  
Deposited on : 2013-08-13  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	 90% 9%
1	B	248	 91% 8%
1	C	248	 90% 10%
1	D	248	 87% 8%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative 3-oxoacyl-[acyl-carrier protein] reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	247	1813	1122	325	355	11	0	0	0
1	B	246	1807	1119	324	353	11	11	0	0
1	C	247	1813	1122	325	355	11	0	0	0
1	D	237	1734	1075	312	337	10	0	0	0



- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	174	Total 174	O 174	0	0
2	B	202	Total 202	O 202	0	0
2	C	155	Total 155	O 155	0	0
2	D	146	Total 146	O 146	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative 3-oxoacyl-[acyl-carrier protein] reductase





- Molecule 1: Putative 3-oxoacyl-[acyl-carrier protein] reductase



- Molecule 1: Putative 3-oxoacyl-[acyl-carrier protein] reductase



- Molecule 1: Putative 3-oxoacyl-[acyl-carrier protein] reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.89Å 111.27Å 117.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.54 - 2.00 34.54 - 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.54-2.00) 100.0 (34.54-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.58 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.173 , 0.199 0.180 , 0.206	Depositor DCC
$R_{free}$ test set	3020 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.4	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 44.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.022 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7844	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/1830	0.66	0/2468
1	B	0.43	0/1824	0.66	0/2460
1	C	0.41	0/1830	0.66	0/2468
1	D	0.42	0/1749	0.65	0/2356
All	All	0.42	0/7233	0.66	0/9752

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1813	0	1841	21	0
1	B	1807	0	1836	15	0
1	C	1813	0	1841	25	0
1	D	1734	0	1765	17	0
2	A	174	0	0	1	0
2	B	202	0	0	1	0
2	C	155	0	0	1	0
2	D	146	0	0	0	0
All	All	7844	0	7283	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:141:THR:O	1:C:184:ALA:HA	1.68	0.94
1:C:200:THR:O	1:C:203:THR:HG22	1.76	0.85
1:D:141:THR:O	1:D:184:ALA:HA	1.78	0.83
1:D:242:ASN:HB2	1:D:245:MET:HE2	1.59	0.81
1:A:141:THR:O	1:A:141:THR:HG23	1.80	0.81
1:B:141:THR:O	1:B:185:PRO:HD2	1.81	0.79
1:B:139:ASN:HB3	1:B:162:LEU:HD21	1.71	0.73
1:A:133:ARG:HG2	1:A:176:ARG:HH21	1.54	0.72
1:A:100:MET:HB3	1:C:126:LEU:CD2	2.20	0.70
1:C:242:ASN:HB2	1:C:245:MET:HG2	1.75	0.68
1:B:242:ASN:HB2	1:B:245:MET:HG2	1.76	0.67
1:C:187:PHE:CD2	1:C:204:PHE:HB3	2.29	0.67
1:A:100:MET:HB3	1:C:126:LEU:HD22	1.78	0.65
1:B:200:THR:O	1:B:203:THR:HG22	1.97	0.64
1:B:141:THR:O	1:B:185:PRO:CD	2.49	0.60
1:A:60:ARG:NH2	1:A:76:ASP:OD2	2.31	0.59
1:A:34:LYS:HE2	1:A:81:PHE:CZ	2.39	0.58
1:D:66:GLU:HG3	1:D:69:THR:HG21	1.87	0.57
1:A:100:MET:HB3	1:C:126:LEU:HD21	1.87	0.57
1:A:141:THR:O	1:A:141:THR:CG2	2.52	0.56
1:B:141:THR:HB	1:B:162:LEU:HD22	1.88	0.56
1:B:123:LYS:NZ	1:D:102:MET:O	2.36	0.55
1:C:63:ASN:HB2	2:C:401:HOH:O	2.08	0.54
1:C:187:PHE:CZ	1:C:204:PHE:CD1	2.95	0.54
1:C:126:LEU:HD21	1:C:130:MET:SD	2.49	0.53
1:C:187:PHE:CE2	1:C:204:PHE:HB3	2.44	0.52
1:A:34:LYS:HE2	1:A:81:PHE:CE1	2.43	0.52
1:A:118:VAL:HG11	1:A:162:LEU:HD11	1.92	0.52
1:D:242:ASN:HB2	1:D:245:MET:CE	2.34	0.52
1:C:200:THR:HA	1:C:203:THR:HG22	1.92	0.52
1:C:10:ILE:CD1	1:C:81:PHE:HB3	2.39	0.52
1:D:143:VAL:HG11	1:D:245:MET:HE1	1.91	0.52
1:D:118:VAL:HG11	1:D:162:LEU:HD11	1.92	0.51
1:D:242:ASN:CB	1:D:245:MET:HE2	2.34	0.51
1:A:102:MET:O	1:C:123:LYS:NZ	2.40	0.51
1:B:97:ASN:OD1	1:B:101:ARG:HD2	2.10	0.51
1:C:118:VAL:HG11	1:C:162:LEU:HD11	1.94	0.50
1:B:139:ASN:CB	1:B:162:LEU:HD21	2.42	0.49
1:C:10:ILE:HD13	1:C:81:PHE:HB3	1.95	0.49
1:A:126:LEU:HD21	1:A:130:MET:SD	2.53	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:GLU:N	1:A:42:GLU:OE1	2.36	0.49
1:B:65:ALA:HA	1:B:112:GLN:NE2	2.27	0.49
1:A:65:ALA:HA	1:A:112:GLN:HE21	1.78	0.48
1:C:199:GLU:O	1:C:203:THR:HB	2.14	0.47
1:C:14:THR:O	1:C:90:ASN:HB3	2.15	0.46
1:C:187:PHE:CZ	1:C:204:PHE:CG	3.03	0.46
1:B:131:LYS:HE3	2:B:490:HOH:O	2.15	0.46
1:D:141:THR:O	1:D:141:THR:HG23	2.15	0.45
1:B:14:THR:O	1:B:90:ASN:HB3	2.16	0.45
1:C:126:LEU:CD2	1:C:130:MET:SD	3.04	0.45
1:A:242:ASN:HB2	1:A:245:MET:HG3	1.98	0.45
1:D:14:THR:O	1:D:90:ASN:HB3	2.16	0.45
1:A:14:THR:O	1:A:90:ASN:HB3	2.16	0.45
1:C:10:ILE:HD13	1:C:81:PHE:CB	2.46	0.45
1:B:198:GLU:O	1:B:202:GLN:HG2	2.17	0.45
1:A:133:ARG:CG	1:A:176:ARG:HH21	2.28	0.45
1:C:200:THR:C	1:C:203:THR:HG22	2.36	0.45
1:C:141:THR:O	1:C:141:THR:HG23	2.16	0.44
1:D:64:SER:O	1:D:120:ARG:NH1	2.50	0.44
1:A:70:VAL:HG21	2:A:323:HOH:O	2.17	0.44
1:D:16:ALA:O	1:D:51:ARG:NH2	2.51	0.43
1:D:34:LYS:HE2	1:D:81:PHE:CZ	2.54	0.43
1:A:126:LEU:CD2	1:A:130:MET:SD	3.07	0.43
1:C:64:SER:O	1:C:120:ARG:NH1	2.52	0.43
1:C:34:LYS:HE2	1:C:81:PHE:CZ	2.54	0.42
1:A:141:THR:O	1:A:184:ALA:HA	2.20	0.42
1:B:102:MET:O	1:D:123:LYS:NZ	2.51	0.42
1:A:79:LYS:HB3	1:A:79:LYS:HE2	1.93	0.41
1:D:66:GLU:HG3	1:D:69:THR:CG2	2.50	0.41
1:D:186:GLY:HA3	1:D:245:MET:HE2	2.03	0.41
1:B:126:LEU:HD23	1:D:100:MET:HG2	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/248 (99%)	241 (98%)	4 (2%)	0	100	100
1	B	244/248 (98%)	241 (99%)	3 (1%)	0	100	100
1	C	245/248 (99%)	242 (99%)	3 (1%)	0	100	100
1	D	233/248 (94%)	231 (99%)	2 (1%)	0	100	100
All	All	967/992 (98%)	955 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/186 (100%)	185 (100%)	0	100	100
1	B	184/186 (99%)	183 (100%)	1 (0%)	88	92
1	C	185/186 (100%)	183 (99%)	2 (1%)	73	78
1	D	176/186 (95%)	176 (100%)	0	100	100
All	All	730/744 (98%)	727 (100%)	3 (0%)	91	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	189	ASP
1	C	42	GLU
1	C	202	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	247/248 (99%)	-0.09	3 (1%) 79 78	9, 20, 42, 60	0
1	B	245/248 (98%)	-0.01	14 (5%) 23 23	10, 17, 52, 71	0
1	C	247/248 (99%)	-0.02	10 (4%) 38 37	10, 20, 46, 74	0
1	D	237/248 (95%)	0.07	13 (5%) 25 24	10, 22, 48, 61	0
All	All	976/992 (98%)	-0.01	40 (4%) 37 36	9, 20, 46, 74	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	203	THR	5.8
1	D	204	PHE	5.1
1	B	192	MET	4.9
1	B	195	ALA	4.1
1	B	3	THR	3.9
1	D	190	THR	3.4
1	B	64	SER	3.4
1	B	204	PHE	3.2
1	C	187	PHE	3.0
1	C	197	PRO	3.0
1	B	202	GLN	2.9
1	C	2	SER	2.9
1	D	202	GLN	2.8
1	D	82	GLY	2.8
1	D	64	SER	2.8
1	C	100	MET	2.7
1	B	200	THR	2.7
1	C	192	MET	2.7
1	C	195	ALA	2.6
1	D	213	PHE	2.6
1	D	203	THR	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	100	MET	2.5
1	B	198	GLU	2.5
1	A	65	ALA	2.5
1	D	201	ARG	2.4
1	D	81	PHE	2.3
1	C	196	LEU	2.3
1	A	133	ARG	2.3
1	B	211	GLY	2.3
1	B	196	LEU	2.2
1	C	200	THR	2.2
1	D	8	GLY	2.2
1	C	204	PHE	2.2
1	B	190	THR	2.2
1	B	141	THR	2.1
1	B	197	PRO	2.1
1	A	68	GLU	2.1
1	D	80	THR	2.1
1	C	199	GLU	2.0
1	D	76	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.