



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 02:45 AM EDT

PDB ID : 6M92  
Title : Monophosphorylated pSer33 b-Catenin peptide, b-TrCP/Skp1, NRX-2663 ternary complex  
Authors : Simonetta, K.R.; Clifton, M.C.; Walter, R.L.; Ranieri, G.M.; Carter, J.J.; Lee, S.J.  
Deposited on : 2018-08-22  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	 81% 12% 7%
2	B	144	 10% 85% 8% 8%
3	C	33	 3% 27% 70%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called F-box/WD repeat-containing protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3171	1998	567	584	22	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	138	SER	-	expression tag	UNP Q9Y297

- Molecule 2 is a protein called S-phase kinase-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	133	985	629	159	192	5	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ASP	deletion	UNP P63208
B	?	-	ASP	deletion	UNP P63208
B	?	-	GLU	deletion	UNP P63208
B	?	-	GLY	deletion	UNP P63208
B	?	-	ASP	deletion	UNP P63208
B	?	-	ASP	deletion	UNP P63208
B	?	-	PRO	deletion	UNP P63208
B	?	-	PRO	deletion	UNP P63208
B	?	-	GLU	deletion	UNP P63208
B	?	-	ASP	deletion	UNP P63208
B	?	-	ASP	deletion	UNP P63208
B	?	-	GLU	deletion	UNP P63208
B	?	-	ASN	deletion	UNP P63208
B	?	-	LYS	deletion	UNP P63208
B	?	-	GLU	deletion	UNP P63208

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	LYS	deletion	UNP P63208
B	?	-	ARG	deletion	UNP P63208
B	?	-	THR	deletion	UNP P63208


- Molecule 3 is a protein called Catenin beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	10	68	38	12	17	1	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	16	CYS	-	expression tag	UNP P35222

- Molecule 4 is 3-{[2-oxo-4-phenoxy-6-(trifluoromethyl)-1,2-dihydropyridine-3-carbonyl]amino}benzoic acid (three-letter code: J8V) (formula: C<sub>20</sub>H<sub>13</sub>F<sub>3</sub>N<sub>2</sub>O<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
4	A	1	30	20	3	2	5	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	33	Total	O	0	0
			33	33		
6	C	4	Total	O	0	0
			4	4		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.59Å 82.59Å 111.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.30 – 2.35 41.30 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.30-2.35) 92.3 (41.30-2.35)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.72 (at 2.34Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, $R_{free}$	0.189 , 0.221 0.189 , 0.221	Depositor DCC
$R_{free}$ test set	1330 reflections (3.77%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.3	Xtrriage
Anisotropy	0.217	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 58.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.014 for -h,-k,l 0.039 for h,-h-k,-l 0.026 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4296	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: J8V, SEP, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3234	0.61	0/4388
2	B	0.32	0/1000	0.53	0/1368
3	C	0.21	0/57	0.42	0/74
All	All	0.35	0/4291	0.59	0/5830

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3171	0	3108	36	0
2	B	985	0	927	12	0
3	C	68	0	54	1	0
4	A	30	0	0	0	0
5	A	5	0	0	0	0
6	A	33	0	0	3	0
6	C	4	0	0	0	0
All	All	4296	0	4089	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.



All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:20:ILE:HD12	2:B:21:ALA:N	1.84	0.92
1:A:470:GLU:OE2	1:A:491:LYS:NZ	2.15	0.79
1:A:476:ILE:HD12	1:A:485:SER:HB3	1.66	0.77
2:B:20:ILE:HD12	2:B:21:ALA:H	1.48	0.77
1:A:196:MET:HA	1:A:199:THR:HG22	1.65	0.77
2:B:16:VAL:HG13	2:B:20:ILE:HD11	1.67	0.77
2:B:16:VAL:CG1	2:B:20:ILE:HD11	2.18	0.74
1:A:363:CYS:HB2	1:A:393:VAL:HG13	1.78	0.66
2:B:16:VAL:CG1	2:B:20:ILE:CD1	2.77	0.63
2:B:16:VAL:HG13	2:B:20:ILE:CD1	2.29	0.63
1:A:200:ASP:OD2	1:A:202:LEU:HB3	1.98	0.62
1:A:363:CYS:HB2	1:A:393:VAL:CG1	2.29	0.62
1:A:512:LEU:O	1:A:513:ARG:HD2	2.00	0.61
1:A:140:LEU:HD23	1:A:141:GLN:HG3	1.83	0.60
1:A:395:VAL:HG21	1:A:435:CYS:HA	1.82	0.60
1:A:140:LEU:CD2	1:A:141:GLN:HG3	2.34	0.58
1:A:452:ILE:HB	1:A:466:LEU:HB2	1.87	0.55
1:A:305:GLY:HA3	1:A:339:MET:HE2	1.88	0.55
1:A:251:ARG:O	1:A:513:ARG:HD3	2.08	0.54
1:A:296:THR:HG23	1:A:298:GLU:H	1.73	0.53
2:B:53:ILE:O	2:B:57:THR:HG23	2.09	0.53
1:A:270:VAL:HB	1:A:535:SER:HB2	1.91	0.52
1:A:476:ILE:CD1	1:A:485:SER:HB3	2.38	0.52
1:A:386:LEU:HD21	1:A:420:CYS:HB3	1.92	0.51
1:A:153:LEU:HD13	1:A:156:ILE:HD12	1.94	0.49
1:A:444:VAL:HG12	1:A:454:LEU:HG	1.94	0.49
1:A:517:GLU:HB2	1:A:544:TRP:CH2	2.48	0.48
1:A:477:ARG:NH2	6:A:701:HOH:O	2.25	0.48
1:A:483:ILE:HB	1:A:495:TRP:HB2	1.96	0.48
1:A:259:ARG:NH1	6:A:703:HOH:O	2.37	0.47
1:A:515:LEU:HB3	1:A:544:TRP:CZ3	2.50	0.46
2:B:49:LEU:HA	2:B:52:VAL:HG12	1.98	0.46
1:A:426:LEU:HD22	1:A:460:GLY:HA2	1.97	0.45
2:B:16:VAL:HG12	2:B:20:ILE:HD11	1.96	0.45
1:A:477:ARG:NE	6:A:701:HOH:O	2.43	0.44
1:A:478:PHE:CD1	1:A:478:PHE:C	2.91	0.44
1:A:437:GLN:OE1	1:A:477:ARG:HA	2.19	0.43
1:A:285:ARG:NH2	3:C:33:SEP:O1P	2.44	0.43
1:A:491:LYS:HG2	1:A:516:VAL:HG22	2.00	0.42
2:B:4:ILE:HD12	2:B:4:ILE:HA	1.79	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:MET:HA	1:A:199:THR:CG2	2.42	0.42
1:A:477:ARG:HA	1:A:477:ARG:HD2	1.83	0.42
1:A:161:LEU:HD13	1:A:183:THR:HG22	2.02	0.42
1:A:476:ILE:HD12	1:A:485:SER:CB	2.44	0.42
1:A:145:ILE:HB	1:A:179:TRP:CE2	2.56	0.41
1:A:293:ASP:HB3	1:A:296:THR:HG22	2.01	0.41
2:B:59:HIS:NE2	2:B:71:ASP:OD2	2.47	0.41
2:B:124:LYS:HB3	2:B:124:LYS:HE2	1.89	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/432 (92%)	375 (94%)	21 (5%)	1 (0%)	41	47
2	B	129/144 (90%)	128 (99%)	1 (1%)	0	100	100
3	C	7/33 (21%)	6 (86%)	1 (14%)	0	100	100
All	All	533/609 (88%)	509 (96%)	23 (4%)	1 (0%)	47	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	345	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/382 (90%)	343 (100%)	1 (0%)	92	96
2	B	102/132 (77%)	102 (100%)	0	100	100
3	C	5/25 (20%)	5 (100%)	0	100	100
All	All	451/539 (84%)	450 (100%)	1 (0%)	93	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	301	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SEP	C	33	3	8,9,10	1.55	1 (12%)	8,12,14	1.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SEP	C	33	3	-	0/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	33	SEP	P-O1P	3.34	1.61	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	33	SEP	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	J8V	A	601	-	32,32,32	1.38	6 (18%)	40,46,46	2.78	6 (15%)
5	PO4	A	602	-	4,4,4	2.09	3 (75%)	6,6,6	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	J8V	A	601	-	-	10/22/22/22	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	601	J8V	C06-N07	3.27	1.48	1.41
4	A	601	J8V	C02-C09	2.74	1.55	1.49
4	A	601	J8V	C08-N07	2.73	1.41	1.35
5	A	602	PO4	P-O2	-2.43	1.47	1.54
4	A	601	J8V	C13-C18	2.42	1.49	1.44
5	A	602	PO4	P-O4	-2.41	1.47	1.54
4	A	601	J8V	C13-C08	2.34	1.52	1.48
5	A	602	PO4	P-O3	-2.14	1.48	1.54
4	A	601	J8V	C20-C16	2.01	1.55	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	601	J8V	C14-C13-C08	15.31	129.59	118.35
4	A	601	J8V	O12-C08-C13	-4.02	115.38	121.53
4	A	601	J8V	C13-C08-N07	3.42	121.75	115.96
4	A	601	J8V	C02-C01-C06	3.16	124.71	120.44
4	A	601	J8V	O19-C18-N17	-3.09	114.19	120.12
4	A	601	J8V	C05-C06-C01	-2.62	116.55	119.65

There are no chirality outliers.


All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	601	J8V	O12-C08-C13-C18
4	A	601	J8V	C03-C02-C09-O10
4	A	601	J8V	C03-C02-C09-O11
4	A	601	J8V	C01-C02-C09-O10
4	A	601	J8V	C01-C02-C09-O11
4	A	601	J8V	C15-C16-C20-F21
4	A	601	J8V	C15-C16-C20-F22
4	A	601	J8V	O12-C08-C13-C14
4	A	601	J8V	N07-C08-C13-C18
4	A	601	J8V	C15-C16-C20-F23

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	401/432 (92%)	0.12	6 (1%) 73 81	55, 78, 104, 127	0
2	B	133/144 (92%)	0.45	14 (10%) 6 9	77, 103, 141, 165	0
3	C	9/33 (27%)	1.07	1 (11%) 5 8	56, 77, 120, 127	0
All	All	543/609 (89%)	0.22	21 (3%) 39 52	55, 83, 121, 165	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	20	ILE	4.8
2	B	18	VAL	3.9
2	B	21	ALA	3.9
2	B	55	TRP	3.2
2	B	56	CYS	3.2
2	B	4	ILE	2.8
2	B	127	PHE	2.7
1	A	446	GLY	2.6
3	C	39	ALA	2.5
2	B	68	PRO	2.5
1	A	436	LEU	2.5
2	B	54	GLN	2.5
2	B	2	PRO	2.4
2	B	75	LEU	2.3
2	B	67	ILE	2.3
1	A	234	ALA	2.3
2	B	69	VAL	2.2
1	A	476	ILE	2.2
1	A	444	VAL	2.2
1	A	547	LEU	2.1
2	B	70	TRP	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SEP	C	33	10/11	0.99	0.17	57,60,70,72	0

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	J8V	A	601	30/30	0.93	0.20	45,64,80,98	30
5	PO4	A	602	5/5	0.97	0.14	114,114,123,124	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.