



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 01:00 pm BST

PDB ID : 5MCV  
Title : New Insights into the Role of DNA Shape on Its Recognition by p53 Proteins  
(complex p53DBD-LWC1)  
Authors : Golovenko, D.; Rozenberg, H.; Shakked, Z.  
Deposited on : 2016-11-10  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	 88% 11%
1	B	200	 88% 12%
2	C	21	 67% 29% 5%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6915 atoms, of which 2569 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	200	2856	1024	1204	307	300	21	0	19	0
1	B	200	2849	1028	1190	307	303	21	0	19	0

- Molecule 2 is a DNA chain called DNA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
2	C	20	585	192	175	78	120	20	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	253	Total O 253 253	0	12
6	B	257	Total O 257 257	0	12
6	C	65	Total O 65 65	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cellular tumor antigen p53

Chain A:  88% 11%




- Molecule 1: Cellular tumor antigen p53

Chain B:  88% 12%



- Molecule 2: DNA

Chain C:  67% 29% 5%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	138.08Å 49.53Å 68.05Å 90.00° 92.91° 90.00°	Depositor
Resolution (Å)	34.48 – 1.60 34.47 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (34.48-1.60) 73.9 (34.47-1.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.79 (at 1.60Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.154 , 0.188 0.157 , 0.192	Depositor DCC
$R_{free}$ test set	3018 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.9	Xtrriage
Anisotropy	0.441	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 59.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.051 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6915	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 97.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7523e-10. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ACT, EDO, 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/1732	0.55	0/2340
1	B	0.35	0/1740	0.56	0/2350
2	C	0.72	0/365	0.84	0/556
All	All	0.40	0/3837	0.59	0/5246

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1652	1204	1631	20	0
1	B	1659	1190	1645	23	0
2	C	410	175	223	4	1
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	20	0	30	2	0
4	B	20	0	30	4	0
5	A	4	0	3	1	0
5	B	4	0	3	1	0
6	A	253	0	0	6	4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	257	0	0	5	3
6	C	65	0	0	3	2
All	All	4346	2569	3565	48	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:133[B]:MET:HE3	4:B:302:EDO:H22	1.58	0.84
1:B:221:GLU:O	1:B:230[B]:THR:HG21	1.83	0.78
1:A:133[B]:MET:HE3	4:A:302:EDO:H22	1.66	0.77
1:A:221:GLU:O	1:A:230[B]:THR:HG21	1.85	0.76
1:A:167:GLN:NE2	6:A:401[B]:HOH:O	2.19	0.74
1:B:167:GLN:NE2	6:B:401[A]:HOH:O	2.20	0.73
2:C:15:DI:OP2	6:C:101:HOH:O	2.15	0.64
1:A:230[B]:THR:HG22	6:A:587:HOH:O	1.98	0.61
1:A:202[A]:ARG:HE	1:B:94:SER:N	2.01	0.58
1:B:203[A]:VAL:HG12	6:B:451:HOH:O	2.04	0.57
1:B:230[B]:THR:HG22	6:B:578:HOH:O	2.04	0.57
5:B:307:ACT:H2	6:B:558:HOH:O	2.04	0.57
1:B:188:LEU:HD23	1:B:203[A]:VAL:CG1	2.36	0.55
2:C:5:DI:OP2	6:C:102:HOH:O	2.18	0.55
5:A:307:ACT:H2	6:A:557:HOH:O	2.07	0.54
1:A:203[A]:VAL:HG12	6:A:477:HOH:O	2.07	0.53
1:B:248[B]:ARG:HD2	6:B:404:HOH:O	2.07	0.53
1:B:133[B]:MET:HE3	4:B:302:EDO:C2	2.38	0.51
1:A:188:LEU:HD23	1:A:203[B]:VAL:CG2	2.42	0.49
1:B:110:ARG:HG2	1:B:146:TRP:HB2	1.94	0.49
1:A:248[B]:ARG:HD2	6:A:403:HOH:O	2.11	0.49
1:B:137:LEU:HD23	1:B:138:ALA:N	2.28	0.48
1:A:110:ARG:HG2	1:A:146:TRP:HB2	1.94	0.48
1:B:200:ASN:HB3	1:B:203[B]:VAL:CG2	2.45	0.47
1:B:98:PRO:HG2	1:B:160[B]:MET:SD	2.55	0.47
1:B:136:GLN:HB2	1:B:139:LYS:HG3	1.97	0.47
1:B:178:HIS:O	1:B:181:ARG:HG3	2.15	0.47
1:A:178:HIS:O	1:A:181:ARG:HG3	2.15	0.46
1:A:98:PRO:HG2	1:A:160[B]:MET:SD	2.56	0.46
1:B:200:ASN:HB3	1:B:203[B]:VAL:HG23	1.99	0.45
1:A:142:PRO:HD2	4:A:302:EDO:H21	1.98	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:11:DG:OP2	6:C:103:HOH:O	2.21	0.45
1:B:142:PRO:HD2	4:B:302:EDO:H21	1.99	0.44
1:B:146:TRP:CD2	4:B:305:EDO:H22	2.52	0.44
1:B:188:LEU:HD23	1:B:203[A]:VAL:HG13	2.00	0.43
1:A:136:GLN:HB2	1:A:139:LYS:HG3	2.00	0.43
2:C:10:DC:H2'	2:C:11:DG:C8	2.53	0.43
1:A:136:GLN:OE1	1:A:139:LYS:NZ	2.51	0.43
1:B:166[B]:SER:HA	1:B:169[B]:MET:HG3	2.00	0.42
1:B:175:ARG:HD3	1:B:191:PRO:O	2.18	0.42
1:B:136:GLN:OE1	1:B:139:LYS:NZ	2.53	0.41
1:A:247:ASN:O	1:A:248[B]:ARG:HB2	2.21	0.41
1:A:175:ARG:HD3	1:A:191:PRO:O	2.20	0.41
1:B:181:ARG:HD2	1:B:181:ARG:C	2.42	0.41
1:A:181:ARG:C	1:A:181:ARG:HD2	2.42	0.40
1:A:200:ASN:HB3	1:A:203[B]:VAL:CG1	2.52	0.40
1:A:192:GLN:HB3	6:A:516[B]:HOH:O	2.22	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1:DG:P	2:C:20:DC:O3'[1_556]	1.60	0.60
6:B:592:HOH:O	6:C:140:HOH:O[2_557]	2.13	0.07
6:A:472:HOH:O	6:B:435:HOH:O[2_557]	2.15	0.05
6:A:591:HOH:O	6:C:151:HOH:O[2_557]	2.15	0.05
6:A:458:HOH:O	6:B:499:HOH:O[2_557]	2.16	0.04
6:A:509:HOH:O	6:A:633:HOH:O[4_557]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/200 (108%)	214 (100%)	1 (0%)	0	100	100
1	B	216/200 (108%)	216 (100%)	0	0	100	100
All	All	431/400 (108%)	430 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/179 (106%)	186 (98%)	3 (2%)	62	41
1	B	191/179 (107%)	188 (98%)	3 (2%)	62	41
All	All	380/358 (106%)	374 (98%)	6 (2%)	60	41

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ARG
1	A	181	ARG
1	A	207	ASP
1	B	110	ARG
1	B	181	ARG
1	B	290	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	5CM	C	6	2	15,21,22	1.82	3 (20%)	19,30,33	2.26	6 (31%)
2	5CM	C	16	2	15,21,22	1.84	4 (26%)	19,30,33	1.67	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5CM	C	6	2	-	1/4/21/22	0/2/2/2
2	5CM	C	16	2	-	1/4/21/22	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	16	5CM	C4-N4	5.67	1.48	1.34
2	C	6	5CM	C4-N4	5.07	1.46	1.34
2	C	6	5CM	C2-N3	-3.76	1.30	1.38
2	C	16	5CM	C2-N3	-2.49	1.33	1.38
2	C	16	5CM	O5'-C5'	-2.24	1.39	1.44
2	C	6	5CM	C5A-C5	2.19	1.55	1.51
2	C	16	5CM	C5A-C5	2.15	1.55	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	5CM	C2-N3-C4	5.49	122.64	116.02
2	C	16	5CM	C5-C6-N1	-5.06	116.75	122.19
2	C	6	5CM	C5-C6-N1	-4.06	117.82	122.19
2	C	6	5CM	N4-C4-N3	3.57	122.08	117.03

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	16	5CM	C2-N3-C4	3.53	120.27	116.02
2	C	6	5CM	C5-C4-N3	-3.00	116.51	121.26
2	C	6	5CM	C5A-C5-C4	-2.77	118.92	121.72
2	C	6	5CM	O4'-C1'-C2'	-2.65	101.25	106.25

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	6	5CM	O4'-C1'-N1-C6
2	C	16	5CM	O4'-C1'-N1-C6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ACT	B	307	-	1,3,3	1.47	0	0,3,3	0.00	-
4	EDO	A	305	-	3,3,3	0.46	0	2,2,2	0.60	0
4	EDO	B	303	-	3,3,3	0.45	0	2,2,2	0.38	0
5	ACT	A	307	-	1,3,3	1.22	0	0,3,3	0.00	-
4	EDO	A	306	-	3,3,3	0.43	0	2,2,2	0.43	0
4	EDO	B	306	-	3,3,3	0.41	0	2,2,2	0.53	0
4	EDO	A	302	-	3,3,3	0.46	0	2,2,2	0.26	0
4	EDO	B	302	-	3,3,3	0.50	0	2,2,2	0.28	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	303	-	3,3,3	0.47	0	2,2,2	0.31	0
4	EDO	B	305	-	3,3,3	0.44	0	2,2,2	0.42	0
4	EDO	B	304	-	3,3,3	0.47	0	2,2,2	0.32	0
4	EDO	A	304	-	3,3,3	0.44	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	305	-	-	0/1/1/1	-
4	EDO	B	303	-	-	0/1/1/1	-
4	EDO	A	306	-	-	0/1/1/1	-
4	EDO	B	306	-	-	0/1/1/1	-
4	EDO	A	302	-	-	0/1/1/1	-
4	EDO	B	302	-	-	0/1/1/1	-
4	EDO	A	303	-	-	0/1/1/1	-
4	EDO	B	305	-	-	0/1/1/1	-
4	EDO	B	304	-	-	1/1/1/1	-
4	EDO	A	304	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	304	EDO	O1-C1-C2-O2
4	A	304	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	307	ACT	1	0
5	A	307	ACT	1	0
4	A	302	EDO	2	0
4	B	302	EDO	3	0
4	B	305	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	200/200 (100%)	-0.65	0 100 100	8, 15, 40, 71	3 (1%)
1	B	200/200 (100%)	-0.63	0 100 100	8, 15, 40, 75	3 (1%)
2	C	16/21 (76%)	-0.18	0 100 100	17, 30, 38, 38	0
All	All	416/421 (98%)	-0.63	0 100 100	8, 15, 40, 75	6 (1%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	5CM	C	6	20/21	0.97	0.07	11,16,23,23	0
2	5CM	C	16	20/21	0.97	0.07	11,16,21,26	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ACT	A	307	4/4	0.50	0.24	31,32,32,32	4
5	ACT	B	307	4/4	0.57	0.23	34,34,34,35	4
4	EDO	B	306	4/4	0.83	0.18	24,24,25,26	4
4	EDO	A	305	4/4	0.86	0.16	23,24,25,26	4
4	EDO	B	305	4/4	0.90	0.13	43,43,44,44	0
4	EDO	A	306	4/4	0.91	0.13	45,45,45,45	0
4	EDO	A	302	4/4	0.93	0.10	28,29,31,32	0
4	EDO	A	304	4/4	0.93	0.15	27,28,28,28	4
4	EDO	B	304	4/4	0.94	0.14	28,28,29,29	4
4	EDO	B	302	4/4	0.95	0.10	28,30,31,32	0
4	EDO	A	303	4/4	0.98	0.08	10,11,12,12	4
4	EDO	B	303	4/4	0.98	0.10	11,13,13,13	4
3	ZN	B	301	1/1	1.00	0.05	12,12,12,12	0
3	ZN	A	301	1/1	1.00	0.05	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.