



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 7, 2020 – 05:16 am BST

PDB ID : 6MDC
Title : Non-receptor Protein Tyrosine Phosphatase SHP2 in Complex with Allosteric Inhibitor Pyrazolo-pyrimidinone 1 SHP389
Authors : Fodor, M.; Stams, T.
Deposited on : 2018-09-04
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	
1	B	526	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	483	Total 3924	C 2477	N 693	O 735	S 19	0	2	0
1	B	470	Total 3816	C 2409	N 677	O 711	S 19	0	1	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q06124
B	0	SER	-	expression tag	UNP Q06124

- Molecule 2 is 6-[(3S,4S)-4-amino-3-methyl-2-oxa-8-azaspiro[4.5]decan-8-yl]-3-[3-chloro-2-(cyclopropylamino)pyridin-4-yl]-5-methyl-2,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one (three-letter code: JEA) (formula: C₂₃H₂₉ClN₈O₂) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	Cl	N	O	0	0
			34	23	1	8	2		
2	B	1	Total	C	Cl	N	O	0	0
			34	23	1	8	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O P	0	0
			5	4 1		
3	A	1	Total	O P	0	0
			5	4 1		
3	A	1	Total	O P	0	0
			5	4 1		
3	B	1	Total	O P	0	0
			5	4 1		
3	B	1	Total	O P	0	0
			5	4 1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	300	Total	O	0	0
			300	300		
4	B	287	Total	O	0	0
			287	287		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.49Å 213.77Å 56.30Å 90.00° 97.02° 90.00°	Depositor
Resolution (Å)	21.38 – 2.14 21.38 – 2.14	Depositor EDS
% Data completeness (in resolution range)	98.0 (21.38-2.14) 98.0 (21.38-2.14)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.13Å)	Xtrriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.194 , 0.239 0.205 , 0.251	Depositor DCC
R_{free} test set	2894 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtrriage
Anisotropy	0.255	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8420	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, JEA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/4011	0.66	0/5408
1	B	0.51	0/3896	0.66	0/5252
All	All	0.51	0/7907	0.66	0/10660

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3924	0	3869	20	0
1	B	3816	0	3768	8	0
2	A	34	0	0	1	0
2	B	34	0	0	1	0
3	A	15	0	0	0	0
3	B	10	0	0	0	0
4	A	300	0	0	1	0
4	B	287	0	0	0	0
All	All	8420	0	7637	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:GLU:HB2	1:A:301:VAL:HG11	1.52	0.88
1:A:232:GLU:HA	1:A:235:LYS:HE2	1.83	0.61
1:A:272:GLU:HB2	1:A:301:VAL:CG1	2.31	0.57
1:A:65:LEU:HD23	1:A:68:GLY:HA3	1.86	0.56
1:A:107:PRO:HG3	1:A:190:LEU:HD12	1.91	0.53
1:B:330:THR:HG23	1:B:458:HIS:HB3	1.90	0.53
1:A:272:GLU:CB	1:A:301:VAL:HG11	2.31	0.52
1:B:309:ILE:HD13	1:B:328:ILE:HG12	1.92	0.51
1:A:309:ILE:HD13	1:A:328:ILE:HG12	1.95	0.49
1:A:132:HIS:HD1	1:A:154:GLY:H	1.60	0.48
1:B:271:GLN:HA	1:B:274:LYS:HE3	1.97	0.47
1:A:222:ASN:O	1:A:225:GLU:HG2	2.16	0.46
1:B:117:LEU:O	1:B:138:ARG:HD2	2.15	0.46
1:A:408:GLN:HB3	1:A:411:THR:HG23	1.98	0.46
2:A:601:JEA:C14	2:A:601:JEA:C5	2.93	0.45
1:B:12:THR:HG22	1:B:35:LYS:HE3	1.98	0.45
1:A:388:VAL:HG21	1:A:402:LYS:HE2	1.98	0.45
1:A:272:GLU:HG2	1:A:272:GLU:H	1.57	0.45
1:A:54:ILE:HD11	1:A:96:ILE:HD13	1.98	0.44
1:B:222:ASN:O	1:B:225:GLU:HG2	2.17	0.44
1:A:56:ILE:HG12	1:A:65:LEU:HD12	2.00	0.43
1:A:125:LEU:HB3	1:A:216:LEU:HD21	2.00	0.43
1:A:498:ARG:HD3	4:A:982:HOH:O	2.19	0.43
1:B:418:TYR:HB3	1:B:438:PHE:CE1	2.54	0.41
1:A:126:LEU:HD23	1:A:216:LEU:HD13	2.02	0.41
2:B:601:JEA:C14	2:B:601:JEA:C5	2.98	0.41
1:A:290:VAL:HG11	1:A:344:MET:HG3	2.01	0.41
1:A:292:LEU:HB2	1:A:303:ASP:HA	2.01	0.41
1:A:377:LEU:HD11	1:A:384:ARG:HG2	2.02	0.41
1:B:284:PRO:HG3	1:B:306:ASN:HA	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/526 (90%)	459 (97%)	12 (2%)	2 (0%)	34	29
1	B	457/526 (87%)	448 (98%)	8 (2%)	1 (0%)	47	45
All	All	930/1052 (88%)	907 (98%)	20 (2%)	3 (0%)	41	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	GLY
1	A	505	VAL
1	B	505	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	432/468 (92%)	420 (97%)	12 (3%)	43	42
1	B	420/468 (90%)	402 (96%)	18 (4%)	29	25
All	All	852/936 (91%)	822 (96%)	30 (4%)	36	33

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	152	ARG
1	A	166	LYS
1	A	173	ARG
1	A	199	LYS
1	A	221	ILE
1	A	313	GLU
1	A	325	LYS
1	A	335	GLN
1	A	402	LYS
1	A	413	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	485	ASP
1	A	502	SER
1	B	37	ASN
1	B	47	ARG
1	B	69	GLU
1	B	96	ILE
1	B	152	ARG
1	B	166	LYS
1	B	199	LYS
1	B	206	LEU
1	B	211	GLN
1	B	275	ASN
1	B	280	LYS
1	B	303	ASP
1	B	372	PRO
1	B	378	LYS
1	B	413	ARG
1	B	477	ASP
1	B	484	VAL
1	B	525	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	281	ASN
1	B	211	GLN
1	B	245	GLN
1	B	256	GLN
1	B	275	ASN
1	B	335	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	B	602	-	4,4,4	2.44	1 (25%)	6,6,6	0.54	0
3	PO4	A	604	-	4,4,4	2.49	1 (25%)	6,6,6	0.54	0
2	JEA	B	601	-	34,39,39	1.79	9 (26%)	32,60,60	1.19	2 (6%)
2	JEA	A	601	-	34,39,39	1.73	8 (23%)	32,60,60	1.27	2 (6%)
3	PO4	B	603	-	4,4,4	1.93	0	6,6,6	0.56	0
3	PO4	A	603	-	4,4,4	2.48	1 (25%)	6,6,6	0.46	0
3	PO4	A	602	-	4,4,4	1.98	1 (25%)	6,6,6	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JEA	B	601	-	-	2/10/42/42	0/6/6/6
2	JEA	A	601	-	-	2/10/42/42	0/6/6/6

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	JEA	C3-C2	5.06	1.50	1.41
2	A	601	JEA	C3-C2	4.19	1.49	1.41
3	A	604	PO4	P-O1	4.14	1.60	1.50
3	A	603	PO4	P-O1	4.09	1.60	1.50
3	B	602	PO4	P-O1	4.00	1.60	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	JEA	C18-N8	3.89	1.41	1.35
2	B	601	JEA	C18-N8	3.58	1.40	1.35
2	A	601	JEA	O1-C11	-3.15	1.40	1.44
2	B	601	JEA	C16-C17	3.06	1.42	1.39
2	B	601	JEA	C6-C7	3.01	1.58	1.53
2	A	601	JEA	C1-N6	-2.99	1.29	1.34
2	B	601	JEA	C8-C7	2.91	1.58	1.53
2	A	601	JEA	C8-C7	2.87	1.58	1.53
2	A	601	JEA	C16-C17	2.72	1.42	1.39
2	A	601	JEA	C6-C7	2.67	1.57	1.53
2	B	601	JEA	C10-C7	2.62	1.56	1.53
2	B	601	JEA	C16-C15	-2.60	1.46	1.49
2	B	601	JEA	C1-N6	-2.31	1.31	1.34
2	B	601	JEA	O1-C11	-2.22	1.41	1.44
3	A	602	PO4	P-O1	2.09	1.55	1.50
2	A	601	JEA	C16-C15	-2.00	1.46	1.49

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	JEA	C6-C7-C10	-4.15	107.76	111.41
2	B	601	JEA	C6-C7-C10	-3.68	108.17	111.41
2	A	601	JEA	C16-C17-CL1	2.89	122.18	119.66
2	B	601	JEA	C16-C17-C18	-2.40	118.00	121.78

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	601	JEA	N2-C4-N3-C5
2	A	601	JEA	N2-C4-N3-C5
2	B	601	JEA	N2-C4-N3-C9
2	A	601	JEA	N2-C4-N3-C9


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	JEA	1	0
2	A	601	JEA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	483/526 (91%)	0.71	58 (12%) 4 5	16, 36, 62, 85	0
1	B	470/526 (89%)	0.65	45 (9%) 8 10	18, 35, 58, 73	0
All	All	953/1052 (90%)	0.68	103 (10%) 5 7	16, 36, 61, 85	0

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	205	THR	7.5
1	A	323	PRO	5.9
1	A	236	LEU	5.9
1	A	296	ASP	5.2
1	B	301	VAL	5.2
1	A	525	LEU	4.8
1	B	409	GLY	4.5
1	A	66	TYR	4.4
1	A	37	ASN	4.1
1	B	205	THR	4.0
1	A	207	GLY	3.9
1	B	177	LEU	3.8
1	A	171	MET	3.8
1	A	38	PRO	3.8
1	A	300	PRO	3.7
1	A	48	ASN	3.7
1	B	426	HIS	3.7
1	A	451	ASP	3.6
1	A	116	HIS	3.6
1	A	481	GLU	3.6
1	A	298	ASN	3.6
1	B	116	HIS	3.5
1	B	120	LYS	3.5
1	B	450	MET	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	523	GLU	3.4
1	B	144	PRO	3.4
1	B	362	ARG	3.4
1	B	524	THR	3.2
1	A	120	LYS	3.2
1	A	313	GLU	3.2
1	A	409	GLY	3.2
1	B	365	SER	3.1
1	A	68	GLY	3.1
1	B	481	GLU	3.1
1	A	325	LYS	3.0
1	B	451	ASP	3.0
1	B	149	LEU	3.0
1	B	525	LEU	2.9
1	A	151	VAL	2.9
1	A	523	GLU	2.9
1	B	137	VAL	2.9
1	A	322	LYS	2.9
1	B	171	MET	2.9
1	A	135	PHE	2.8
1	A	127	THR	2.8
1	A	182	GLY	2.8
1	B	37	ASN	2.8
1	B	151	VAL	2.8
1	B	38	PRO	2.8
1	A	248	TRP	2.8
1	A	314	PHE	2.8
1	B	410	ASN	2.8
1	A	353	ILE	2.7
1	A	137	VAL	2.7
1	A	305	ILE	2.7
1	A	67	GLY	2.7
1	A	177	LEU	2.7
1	B	74	LEU	2.7
1	A	297	PRO	2.6
1	A	83	GLU	2.6
1	B	47	ARG	2.6
1	B	143	HIS	2.6
1	A	117	LEU	2.6
1	A	118	SER	2.6
1	A	299	GLU	2.6
1	A	410	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	155	ASP	2.5
1	B	425	ASP	2.5
1	A	84	HIS	2.5
1	B	293	HIS	2.5
1	B	364	LYS	2.5
1	B	363	GLY	2.5
1	A	235	LYS	2.5
1	B	311	MET	2.5
1	A	115	GLY	2.4
1	B	353	ILE	2.4
1	B	142	SER	2.4
1	A	450	MET	2.4
1	B	207	GLY	2.4
1	B	128	GLU	2.3
1	B	235	LYS	2.3
1	B	68	GLY	2.3
1	A	294	ASP	2.3
1	A	136	LEU	2.3
1	B	490	VAL	2.3
1	A	394	HIS	2.3
1	B	48	ASN	2.3
1	A	123	GLU	2.3
1	A	35	LYS	2.2
1	B	188	ASP	2.2
1	A	524	THR	2.2
1	A	231	ARG	2.2
1	B	195	GLU	2.1
1	B	275	ASN	2.1
1	A	203	VAL	2.1
1	B	283	LEU	2.1
1	A	188	ASP	2.1
1	B	66	TYR	2.1
1	A	324	LYS	2.0
1	A	74	LEU	2.0
1	A	149	LEU	2.0
1	B	136	LEU	2.0
1	A	301	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	A	602	5/5	0.77	0.22	120,120,121,121	0
3	PO4	A	604	5/5	0.83	0.17	80,80,80,82	0
3	PO4	A	603	5/5	0.89	0.15	86,86,87,87	0
2	JEA	B	601	34/34	0.91	0.14	17,28,42,44	0
3	PO4	B	603	5/5	0.91	0.14	70,72,75,75	0
2	JEA	A	601	34/34	0.93	0.14	18,25,41,42	0
3	PO4	B	602	5/5	0.97	0.08	43,43,45,48	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around JEA B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around JEA A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.