

# Full wwPDB X-ray Structure Validation Report (i)

#### Dec 9, 2024 – 02:05 PM EST

PDB ID : 9MED

Title : Crystal Structure Pyrophosphate-fructose 6-phosphate 1-phosphotransferase

1 (Pfk1) from Trichomonas vaginalis (AMP/beta-D-Glucose-6-phosphate

complex)

Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural

Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2024-12-06

Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$ 

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.21

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.004 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.40




## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
$R_{free}$	164625	4623 (2.64-2.60)
Clashscore	180529	5071 (2.64-2.60)
Ramachandran outliers	177936	5006 (2.64-2.60)
Sidechain outliers	177891	5006 (2.64-2.60)
RSRZ outliers	164620	4622 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	434	% 	7% •
1	В	434	91%	6% • •
1	С	434	90%	7% •
1	D	434	91%	6% •



## 2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 13064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Pyrophosphate--fructose 6-phosphate 1-phosphotransferase 1.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	Λ	417	Total	С	N	О	S	0	0	0
1	A	417	3191	2028	556	591	16	U	U	U
1	В	423	Total	С	N	О	S	0	0	0
1	Ъ	420	3248	2062	566	603	17	0	0	
1	С	423	Total	С	N	О	S	0	0	0
1		420	3244	2060	566	601	17	U		
1	D	417	Total	С	N	О	S	0	0	0
1	ע	417	3180	2019	553	591	17	U		U

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP O61068
A	-6	ALA	-	expression tag	UNP O61068
A	-5	HIS	-	expression tag	UNP O61068
A	-4	HIS	-	expression tag	UNP O61068
A	-3	HIS	-	expression tag	UNP O61068
A	-2	HIS	-	expression tag	UNP O61068
A	-1	HIS	-	expression tag	UNP O61068
A	0	HIS	-	expression tag	UNP O61068
В	-7	MET	-	initiating methionine	UNP O61068
В	-6	ALA	-	expression tag	UNP O61068
В	-5	HIS	-	expression tag	UNP O61068
В	-4	HIS	-	expression tag	UNP O61068
В	-3	HIS	-	expression tag	UNP O61068
В	-2	HIS	-	expression tag	UNP O61068
В	-1	HIS	-	expression tag	UNP O61068
В	0	HIS	-	expression tag	UNP O61068
С	-7	MET	-	initiating methionine	UNP O61068
С	-6	ALA	-	expression tag	UNP O61068
С	-5	HIS	-	expression tag	UNP O61068
С	-4	HIS	-	expression tag	UNP O61068
С	-3	HIS	-	expression tag	UNP O61068



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	HIS	-	expression tag	UNP O61068
С	-1	HIS	-	expression tag	UNP O61068
С	0	HIS	-	expression tag	UNP O61068
D	-7	MET	_	initiating methionine	UNP O61068
D	-6	ALA	-	expression tag	UNP O61068
D	-5	HIS	_	expression tag	UNP O61068
D	-4	HIS	-	expression tag	UNP O61068
D	-3	HIS	-	expression tag	UNP O61068
D	-2	HIS	_	expression tag	UNP O61068
D	-1	HIS	_	expression tag	UNP O61068
D	0	HIS	-	expression tag	UNP O61068

 $\bullet$  Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula:  $\mathrm{O_4P}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	В	1	Total O P 5 4 1	0	0
2	В	1	Total O P 5 4 1	0	0
2	В	1	Total O P 5 4 1	0	0
2	С	1	Total O P 5 4 1	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	1	Total O P 5 4 1	0	0
2	С	1	Total O P 10 8 2	0	1
2	D	1	Total O P 5 4 1	0	0

• Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula:  $C_{10}H_{14}N_5O_7P$ ) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
3	Λ	1	Total	С	N	О	Р	0	0	
9	Λ	1	23	10	5	7	1	U		
3	В	1	Total	С	N	О	Р	0	0	
9	Б	1	23	10	5	7	1		0	
3	С	1	Total	С	N	О	Р	0	0	
3		1	23	10	5	7	1	U		
9	D	1	Total	С	N	О	Р	0	0	
3	D	1	23	10	5	7	1	U	U	

• Molecule 4 is PYROPHOSPHATE (three-letter code: PPV) (formula:  $H_4O_7P_2$ ) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	А	1	Total	О	Р	0	0
_ T	11	1	9	7	2		

• Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0

• Molecule 6 is 6-O-phosphono-beta-D-glucopyranose (three-letter code: BG6) (formula:  $C_6H_{13}O_9P$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	Total C O P 16 6 9 1	0	0
6	С	1	Total C O P 16 6 9 1	0	0

 $\bullet$  Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	С	1	Total Cl 1 1	0	0

#### • Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	2	Total O 2 2	0	0
8	В	4	Total O 4 4	0	0
8	С	5	Total O 5 5	0	0
8	D	5	Total O 5 5	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyrophosphate--fructose 6-phosphate 1-phosphotransferase 1









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	86.88Å 86.88Å 475.98Å	Donogitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.57 - 2.62	Depositor
rtesolution (A)	48.57 - 2.62	EDS
% Data completeness	100.0 (48.57-2.62)	Depositor
(in resolution range)	100.0 (48.57-2.62)	EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.83 (at 2.61Å)	Xtriage
Refinement program	PHENIX dev_5508	Depositor
P. P.	0.192 , 0.234	Depositor
$R, R_{free}$	0.197 , $0.234$	DCC
$R_{free}$ test set	2768 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	67.6	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.32, 53.4	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.45, < L^2>=0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13064	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PPV, MG, AMP, BG6, CL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.26	0/3250	0.52	0/4396
1	В	0.25	0/3308	0.51	0/4472
1	С	0.26	0/3304	0.53	0/4467
1	D	0.25	0/3236	0.52	0/4377
All	All	0.25	0/13098	0.52	0/17712

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	3191	0	3231	21	0
1	В	3248	0	3299	16	0
1	С	3244	0	3295	15	0
1	D	3180	0	3216	8	0
2	A	10	0	0	0	0
2	В	15	0	0	0	0
2	С	20	0	0	0	0
2	D	5	0	0	0	0
3	A	23	0	12	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	23	0	12	0	0
3	С	23	0	12	0	0
3	D	23	0	12	0	0
4	A	9	0	0	0	0
5	A	1	0	0	0	0
6	В	16	0	11	2	0
6	С	16	0	10	2	0
7	С	1	0	0	0	0
8	A	2	0	0	0	0
8	В	4	0	0	0	0
8	С	5	0	0	0	0
8	D	5	0	0	0	0
All	All	13064	0	13110	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:HIS:HE2	6:C:503:BG6:HC61	1.50	0.77
1:A:87:ARG:NH1	1:A:365:ARG:NH2	2.36	0.73
1:A:204:HIS:NE2	6:C:503:BG6:HC61	2.09	0.66
1:D:342:LEU:HD12	1:D:358:LEU:HD12	1.79	0.64
1:B:346:GLU:OE1	1:B:351:LYS:HE2	1.99	0.63
1:A:355:PHE:CE2	1:A:359:LEU:HD21	2.34	0.61
1:D:114:ASP:OD1	1:D:115:ASP:N	2.33	0.61
1:C:342:LEU:HD12	1:C:358:LEU:HD12	1.83	0.60
1:B:204:HIS:HE2	6:B:502:BG6:HC61	1.68	0.58
1:C:275:ILE:HD12	1:C:275:ILE:H	1.68	0.57
1:D:363:THR:HG22	1:D:365:ARG:HD3	1.86	0.57
1:C:214:GLN:HE22	1:C:423:VAL:HG13	1.69	0.57
1:C:90:PRO:HA	1:C:93:GLN:HB2	1.86	0.56
1:A:112:GLY:HA2	1:A:139:LYS:HE3	1.86	0.56
1:C:124:ALA:HB2	1:C:134:VAL:HG21	1.88	0.55
1:C:270:ASP:HB3	1:C:276:LEU:HD21	1.90	0.54
1:D:20:LEU:HD21	1:D:139:LYS:HD3	1.90	0.53
1:A:52:ASP:OD2	1:A:55:VAL:HG23	2.07	0.53
1:A:121:VAL:HG11	1:A:355:PHE:CG	2.43	0.53
1:A:136:SER:HB3	1:A:342:LEU:CD2	2.39	0.52
1:A:87:ARG:HH12	1:A:365:ARG:NH2	2.05	0.51



 $Continued\ from\ previous\ page...$ 

Continued from prediction		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:139:LYS:O	1:A:140:THR:HB	2.10	0.51	
1:B:270:ASP:HB3	1:B:276:LEU:HD21	1.91	0.51	
1:B:342:LEU:HD12	1:B:358:LEU:HD12	1.93	0.50	
1:A:121:VAL:HG11	1:A:355:PHE:CD2	2.46	0.50	
1:C:56:VAL:O	1:C:60:THR:HG23	2.12	0.49	
1:C:179:ARG:HG3	1:C:180:TRP:N	2.26	0.49	
1:A:270:ASP:HB3	1:A:276:LEU:HD21	1.94	0.48	
1:B:136:SER:HB3	1:B:342:LEU:CD2	2.44	0.48	
1:C:141:ILE:HD11	1:C:159:ALA:HB2	1.96	0.47	
1:B:56:VAL:O	1:B:60:THR:HG23	2.14	0.47	
1:A:147:LEU:HD11	1:A:153:THR:HG22	1.97	0.47	
1:C:52:ASP:OD2	1:C:55:VAL:HG23	2.15	0.47	
1:B:20:LEU:HD21	1:B:139:LYS:HD3	1.98	0.46	
1:B:179:ARG:HG3	1:B:180:TRP:N	2.31	0.46	
1:B:12:CYS:SG	1:B:77:THR:HG21	2.56	0.45	
1:A:275:ILE:HD12	1:A:275:ILE:H	1.80	0.45	
1:D:124:ALA:HB2	1:D:134:VAL:HG21	1.97	0.45	
1:A:179:ARG:HG3	1:A:180:TRP:N	2.31	0.45	
1:A:110:THR:O	1:A:136:SER:HA	2.17	0.45	
1:B:346:GLU:OE1	1:B:351:LYS:CE	2.64	0.45	
1:B:214:GLN:CG	1:B:216:GLU:O	2.65	0.45	
1:B:204:HIS:NE2	6:B:502:BG6:HC61	2.31	0.44	
1:D:174:SER:O	1:D:178:PRO:HA	2.17	0.44	
1:A:317:ALA:HB2	1:C:161:SER:HA	2.01	0.43	
1:C:179:ARG:HB2	1:C:300:ARG:HG3	2.01	0.43	
1:D:138:PRO:HD2	1:D:343:ILE:O	2.19	0.43	
1:B:382:ARG:NH1	1:B:387:ARG:CZ	2.82	0.42	
1:A:186:MET:HB3	1:A:186:MET:HE2	1.87	0.42	
1:B:138:PRO:HB2	1:B:145:LEU:HD11	2.02	0.42	
1:B:52:ASP:OD2	1:B:55:VAL:HG23	2.20	0.42	
1:A:342:LEU:HD12	1:A:358:LEU:CD1	2.49	0.42	
1:D:217:ILE:HD12	1:D:252:LYS:HB3	2.02	0.42	
1:C:138:PRO:HD2	1:C:343:ILE:O	2.20	0.42	
1:A:174:SER:O	1:A:178:PRO:HA	2.20	0.41	
1:B:138:PRO:HB2	1:B:145:LEU:CD1	2.50	0.41	
1:C:11:LEU:C	1:C:11:LEU:HD12	2.40	0.41	
1:C:135:ILE:CG2	1:C:343:ILE:HG12	2.51	0.41	
1:A:180:TRP:HA	1:A:241:GLY:O	2.20	0.41	

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	413/434 (95%)	403 (98%)	8 (2%)	2 (0%)	25	45
1	В	421/434 (97%)	414 (98%)	6 (1%)	1 (0%)	44	65
1	С	421/434 (97%)	412 (98%)	8 (2%)	1 (0%)	44	65
1	D	413/434 (95%)	404 (98%)	8 (2%)	1 (0%)	44	65
All	All	1668/1736 (96%)	1633 (98%)	30 (2%)	5 (0%)	37	57

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	140	THR
1	D	307	GLY
1	A	307	GLY
1	В	307	GLY
1	С	307	GLY

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	337/355~(95%)	331 (98%)	6 (2%)	54 76
1	В	345/355 (97%)	338 (98%)	7 (2%)	50 73
1	С	344/355 (97%)	337 (98%)	7 (2%)	50 73
1	D	335/355 (94%)	328 (98%)	7 (2%)	48 72
All	All	1361/1420 (96%)	1334 (98%)	27 (2%)	50 73



All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	THR
1	A	179	ARG
1	A A	218	GLU
1	A	322	TYR
1	Α	324	ARG
1	A	382	ARG
1	В	69	SER
1	В	127	MET
1	В	179	ARG
1	В	186	MET
1	В	270	ASP
1	В	322	TYR
1	В	382	ARG
1	С	77	THR
1	С	179	ARG
1	C C C C C D	272	HIS
1	С	300	ARG
1	С	322	TYR
1	С	345	ARG
1	С	382	ARG
1		127	MET
1	D	141	ILE
1	D	179	ARG
1	D	300	ARG
1	D	322	TYR
1	D	376	GLN
1	D	382	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	ths	Bond angles		
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	504	-	4,4,4	1.59	1 (25%)	6,6,6	0.48	0
2	PO4	С	506[A]	-	4,4,4	1.54	1 (25%)	6,6,6	0.44	0
2	PO4	С	505	-	4,4,4	1.59	1 (25%)	6,6,6	0.49	0
3	AMP	В	501	-	21,25,25	0.78	1 (4%)	23,38,38	1.33	2 (8%)
2	PO4	С	506[B]	-	4,4,4	1.58	1 (25%)	6,6,6	0.49	0
6	BG6	С	503	-	16,16,16	1.28	1 (6%)	23,24,24	1.48	4 (17%)
2	PO4	В	505	-	4,4,4	1.60	1 (25%)	6,6,6	0.50	0
3	AMP	A	502	-	21,25,25	0.81	0	23,38,38	1.37	2 (8%)
2	PO4	В	504	-	4,4,4	1.57	1 (25%)	6,6,6	0.44	0
3	AMP	С	502	-	21,25,25	0.79	0	23,38,38	1.35	2 (8%)
3	AMP	D	501	-	21,25,25	0.80	0	23,38,38	1.33	2 (8%)
2	PO4	С	504	-	4,4,4	1.50	1 (25%)	6,6,6	0.55	0
2	PO4	D	502	-	4,4,4	1.58	1 (25%)	6,6,6	0.51	0
4	PPV	A	503	5	6,8,8	0.62	0	12,13,13	0.78	0
6	BG6	В	502	-	16,16,16	1.31	1 (6%)	23,24,24	1.46	3 (13%)
2	PO4	A	501	-	4,4,4	1.58	1 (25%)	6,6,6	0.53	0
2	PO4	В	503	-	4,4,4	1.59	1 (25%)	6,6,6	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	В	501	-	-	0/6/26/26	0/3/3/3
6	BG6	С	503	-	-	5/6/26/26	0/1/1/1
3	AMP	С	502	-	-	0/6/26/26	0/3/3/3
3	AMP	A	502	-	-	2/6/26/26	0/3/3/3
3	AMP	D	501	-	-	2/6/26/26	0/3/3/3
4	PPV	A	503	5	-	2/6/6/6	-
6	BG6	В	502	-	-	5/6/26/26	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
6	В	502	BG6	O5-C1	3.58	1.51	1.42
6	С	503	BG6	O5-C1	3.53	1.51	1.42
2	В	503	PO4	P-O1	2.80	1.57	1.50
2	В	505	PO4	P-O1	2.77	1.57	1.50
2	С	506[B]	PO4	P-O1	2.77	1.57	1.50
2	С	505	PO4	P-O1	2.76	1.57	1.50
2	A	504	PO4	P-O1	2.75	1.57	1.50
2	A	501	PO4	P-O1	2.74	1.57	1.50
2	D	502	PO4	P-O1	2.74	1.57	1.50
2	В	504	PO4	P-O1	2.73	1.57	1.50
2	С	506[A]	PO4	P-O1	2.72	1.56	1.50
2	С	504	PO4	P-O1	2.57	1.56	1.50
3	В	501	AMP	C2-N3	2.05	1.35	1.32

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
6	В	502	BG6	O5-C1-C2	4.60	118.38	110.30
6	С	503	BG6	O5-C1-C2	4.41	118.06	110.30
3	С	502	AMP	N3-C2-N1	-4.36	122.75	128.67
3	D	501	AMP	N3-C2-N1	-4.30	122.83	128.67
3	В	501	AMP	N3-C2-N1	-4.29	122.85	128.67
3	A	502	AMP	N3-C2-N1	-4.27	122.87	128.67
6	С	503	BG6	C1-C2-C3	2.76	115.99	110.36
6	С	503	BG6	C3-C4-C5	-2.69	105.36	110.23
3	A	502	AMP	C4-C5-N7	-2.44	106.76	109.34
6	В	502	BG6	C3-C4-C5	-2.39	105.90	110.23
3	С	502	AMP	C4-C5-N7	-2.37	106.84	109.34
3	D	501	AMP	C4-C5-N7	-2.31	106.90	109.34
6	С	503	BG6	O5-C5-C6	2.16	110.97	106.69
3	В	501	AMP	C4-C5-N7	-2.14	107.08	109.34



Continued from previous page...

$\mathbf{Mol}$	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
6	В	502	BG6	C1-C2-C3	2.01	114.46	110.36

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	PPV	P1-OPP-P2-O32
6	В	502	BG6	O5-C5-C6-O6
6	В	502	BG6	C4-C5-C6-O6
6	В	502	BG6	C6-O6-P-O2P
6	В	502	BG6	C6-O6-P-O3P
6	С	503	BG6	O5-C5-C6-O6
6	С	503	BG6	C4-C5-C6-O6
6	С	503	BG6	C6-O6-P-O2P
6	С	503	BG6	C6-O6-P-O3P
3	A	502	AMP	O4'-C4'-C5'-O5'
3	D	501	AMP	O4'-C4'-C5'-O5'
3	A	502	AMP	C3'-C4'-C5'-O5'
3	D	501	AMP	C3'-C4'-C5'-O5'
6	С	503	BG6	C6-O6-P-O1P
4	A	503	PPV	P1-OPP-P2-O22
6	В	502	BG6	C6-O6-P-O1P


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	С	503	BG6	2	0
6	В	502	BG6	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	417/434 (96%)	-0.25	4 (0%) 79	76	44, 72, 135, 161	0
1	В	423/434 (97%)	-0.37	0 100 100	0	44, 77, 122, 174	0
1	С	423/434 (97%)	-0.40	1 (0%) 92	90	42, 69, 111, 160	0
1	D	417/434 (96%)	-0.31	3 (0%) 84	81	47, 87, 141, 173	0
All	All	1680/1736 (96%)	-0.33	8 (0%) 87	85	42, 76, 132, 174	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	363	THR	3.7
1	D	355	PHE	3.3
1	A	358	LEU	2.6
1	A	145	LEU	2.5
1	A	141	ILE	2.5
1	D	368	THR	2.2
1	A	132	ILE	2.2
1	С	362	ALA	2.1

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ m \AA}^2)$	Q<0.9
5	MG	A	505	1/1	0.64	0.17	108,108,108,108	0
2	PO4	С	505	5/5	0.68	0.13	131,137,146,165	0
2	PO4	D	502	5/5	0.74	0.07	85,98,112,114	0
4	PPV	A	503	9/9	0.75	0.10	111,123,131,139	0
3	AMP	С	502	23/23	0.79	0.10	75,90,105,114	0
2	PO4	В	505	5/5	0.81	0.09	97,105,126,128	0
3	AMP	D	501	23/23	0.83	0.09	80,98,108,114	0
2	PO4	A	504	5/5	0.88	0.08	75,77,93,101	0
2	PO4	A	501	5/5	0.89	0.07	79,91,116,119	0
3	AMP	В	501	23/23	0.89	0.09	74,82,93,99	0
2	PO4	В	504	5/5	0.90	0.11	82,82,88,99	0
2	PO4	С	506[A]	5/5	0.90	0.12	59,76,79,83	5
2	PO4	С	506[B]	5/5	0.90	0.12	66,75,82,83	5
6	BG6	В	502	16/16	0.91	0.10	74,82,89,91	0
3	AMP	A	502	23/23	0.92	0.07	51,69,79,91	0
2	PO4	С	504	5/5	0.93	0.09	61,67,71,73	0
2	PO4	В	503	5/5	0.93	0.09	69,70,84,85	0
6	BG6	С	503	16/16	0.93	0.10	60,71,79,87	0
7	CL	С	501	1/1	0.94	0.14	88,88,88,88	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




## Electron density around PPV A 503:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around AMP C 502:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












#### Electron density around AMP D 501:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around AMP B 501:

 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











# 







## 6.5 Other polymers (i)

There are no such residues in this entry.

