



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 30, 2023 – 12:01 PM EDT

PDB ID : 3MG4
Title : Structure of yeast 20S proteasome with Compound 1
Authors : Sintchak, M.D.
Deposited on : 2010-04-05
Resolution : 3.11 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1292 (3.14-3.10)
Clashscore	141614	1389 (3.14-3.10)
Ramachandran outliers	138981	1337 (3.14-3.10)
Sidechain outliers	138945	1337 (3.14-3.10)
RSRZ outliers	127900	1260 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	
1	O	250	
2	B	244	
2	P	244	
3	C	241	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	241	6% 74% 22% .
4	D	242	4% 83% 15% .
4	R	242	3% 82% 17% .
5	E	233	3% 80% 19% .
5	S	233	3% 76% 21% .
6	F	244	77% 20% .
6	T	244	% 81% 16% .
7	G	243	77% 21% .
7	U	243	75% 21% .
8	H	222	86% 13% .
8	V	222	80% 18% .
9	I	204	86% 13%
9	W	204	83% 15% .
10	J	198	% 81% 19% .
10	X	198	2% 84% 14% .
11	K	212	78% 19% .
11	Y	212	79% 18% .
12	L	222	77% 21% .
12	Z	222	77% 18% 5%
13	1	233	80% 17% .
13	M	233	83% 16% .
14	2	196	% 80% 19% .
14	N	196	80% 20% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
15	MG	F	242	-	-	-	X
15	MG	G	241	-	-	-	X

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 49672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome component Y7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome component Y13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	Total	C	N	O	S	0	0	0
			1905	1201	321	380	3			
2	P	244	Total	C	N	O	S	0	0	0
			1905	1201	321	380	3			

- Molecule 3 is a protein called Proteasome component PRE6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	241	Total	C	N	O	S	0	0	0
			1891	1181	331	375	4			
3	Q	241	Total	C	N	O	S	0	0	0
			1891	1181	331	375	4			

- Molecule 4 is a protein called Proteasome component PUP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	242	Total	C	N	O	S	0	0	0
			1862	1162	314	379	7			
4	R	242	Total	C	N	O	S	0	0	0
			1862	1162	314	379	7			

- Molecule 5 is a protein called Proteasome component PRE5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	233	Total 1795	C 1129	N 312	O 350	S 4	0	0	0
5	S	233	Total 1795	C 1129	N 312	O 350	S 4	0	0	0

- Molecule 6 is a protein called Proteasome component C1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	244	Total 1897	C 1205	N 330	O 358	S 4	0	0	0
6	T	244	Total 1897	C 1205	N 330	O 358	S 4	0	0	0

- Molecule 7 is a protein called Proteasome component C7-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	243	Total 1921	C 1221	N 322	O 370	S 8	0	0	0
7	U	243	Total 1921	C 1221	N 322	O 370	S 8	0	0	0

- Molecule 8 is a protein called Proteasome component PUP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	222	Total 1685	C 1061	N 293	O 324	S 7	0	0	0
8	V	222	Total 1685	C 1061	N 293	O 324	S 7	0	0	0

- Molecule 9 is a protein called Proteasome component PUP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	204	Total 1581	C 1010	N 258	O 305	S 8	0	0	0
9	W	204	Total 1581	C 1010	N 258	O 305	S 8	0	0	0

- Molecule 10 is a protein called Proteasome component C11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	J	198	Total 1585	C 1005	N 269	O 305	S 6	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

- Molecule 11 is a protein called Proteasome component PRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome component C5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome component PRE4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	1	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome component PRE3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	2	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	D	1	Total	Mg	0	0
			1	1		
15	F	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
15	G	2	Total Mg 2 2	0	0
15	H	1	Total Mg 1 1	0	0
15	I	2	Total Mg 2 2	0	0
15	K	1	Total Mg 1 1	0	0
15	L	1	Total Mg 1 1	0	0
15	N	1	Total Mg 1 1	0	0

- Molecule 16 is (2S)-2-amino-N-[(1S)-1-((1S)-1-[(4-methylbenzyl)carbamoyl]-3-phenylpropyl)carbamoyl]-3-phenylpropyl]-4-phenylbutanamide (three-letter code: LXT) (formula: C₃₈H₄₄N₄O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	K	1	Total C N O 45 38 4 3	0	0
16	Y	1	Total C N O 45 38 4 3	0	0

- Molecule 17 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
17	K	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
17	Y	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome component Y7




- Molecule 1: Proteasome component Y7




- Molecule 2: Proteasome component Y13





- Molecule 2: Proteasome component Y13









- Molecule 3: Proteasome component PRE6





- Molecule 3: Proteasome component PRE6




- Molecule 4: Proteasome component PUP2




- Molecule 4: Proteasome component PUP2




- Molecule 5: Proteasome component PRE5




- Molecule 5: Proteasome component PRE5




- Molecule 6: Proteasome component C1



- Molecule 6: Proteasome component C1





- Molecule 7: Proteasome component C7-alpha





- Molecule 7: Proteasome component C7-alpha

Chain U: 75% 21%





- Molecule 8: Proteasome component PUP1

Chain H: 86% 13%



- Molecule 8: Proteasome component PUP1

Chain V: 80% 18%



- Molecule 9: Proteasome component PUP3

Chain I: 86% 13%




- Molecule 9: Proteasome component PUP3


Chain W: 83% 15%




- Molecule 10: Proteasome component C11




- Molecule 10: Proteasome component C11




- Molecule 11: Proteasome component PRE2




- Molecule 11: Proteasome component PRE2




- Molecule 12: Proteasome component C5




- Molecule 12: Proteasome component C5




- Molecule 13: Proteasome component PRE4




- Molecule 13: Proteasome component PRE4



- Molecule 14: Proteasome component PRE3



- Molecule 14: Proteasome component PRE3



M175	V176	L178	P187D
V177	T179	D187E	D187E
A180	A180	E187F	E187F
		L187J	L187J

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	134.56Å 300.79Å 144.71Å 90.00° 112.73° 90.00°	Depositor
Resolution (Å)	50.00 – 3.11 47.61 – 3.11	Depositor EDS
% Data completeness (in resolution range)	93.9 (50.00-3.11) 93.9 (47.61-3.11)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 3.12Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.203 , 0.250 0.200 , 0.244	Depositor DCC
R_{free} test set	3565 reflections (2.01%)	wwPDB-VP
Wilson B-factor (Å ²)	67.9	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 19.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	49672	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MES, LXT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/1952	0.57	0/2642
1	O	0.44	0/1952	0.59	0/2642
2	B	0.42	0/1935	0.58	0/2618
2	P	0.43	0/1935	0.61	0/2618
3	C	0.45	2/1920 (0.1%)	0.59	0/2598
3	Q	0.45	0/1920	0.60	1/2598 (0.0%)
4	D	0.46	1/1887 (0.1%)	0.60	0/2541
4	R	0.44	0/1887	0.59	0/2541
5	E	0.40	0/1823	0.57	0/2463
5	S	0.42	0/1823	0.57	0/2463
6	F	0.41	0/1937	0.55	0/2614
6	T	0.44	0/1937	0.58	0/2614
7	G	0.43	0/1959	0.57	0/2652
7	U	0.45	0/1959	0.60	0/2652
8	H	0.43	0/1716	0.58	0/2326
8	V	0.42	0/1716	0.59	0/2326
9	I	0.42	0/1611	0.59	0/2174
9	W	0.44	0/1611	0.60	0/2174
10	J	0.42	0/1613	0.58	0/2173
10	X	0.42	0/1613	0.58	0/2173
11	K	0.43	0/1681	0.58	1/2274 (0.0%)
11	Y	0.43	0/1681	0.60	1/2274 (0.0%)
12	L	0.44	0/1795	0.57	0/2420
12	Z	0.47	0/1795	0.60	0/2420
13	1	0.44	0/1855	0.62	1/2514 (0.0%)
13	M	0.44	0/1855	0.61	0/2514
14	2	0.43	0/1541	0.59	0/2087
14	N	0.41	0/1541	0.55	0/2087
All	All	0.43	3/50450 (0.0%)	0.59	4/68192 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if

the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	O	0	1
4	D	0	1
12	Z	0	1
All	All	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	123(G)	GLU	C-O	7.39	1.37	1.23
3	C	57	LYS	CE-NZ	6.06	1.64	1.49
3	C	203	THR	C-O	5.08	1.32	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	Y	4	LEU	CA-CB-CG	5.38	127.68	115.30
11	K	4	LEU	CA-CB-CG	5.25	127.38	115.30
3	Q	56	LEU	CA-CB-CG	5.08	126.98	115.30
13	1	62	LEU	CA-CB-CG	5.07	126.97	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	D	123(G)	GLU	Mainchain
1	O	217(N)	THR	Peptide
12	Z	145	TYR	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1926	24	0
1	O	1915	0	1926	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1905	0	1901	32	0
2	P	1905	0	1901	36	0
3	C	1891	0	1900	30	0
3	Q	1891	0	1900	40	0
4	D	1862	0	1836	23	0
4	R	1862	0	1836	21	0
5	E	1795	0	1797	20	0
5	S	1795	0	1797	28	0
6	F	1897	0	1886	28	0
6	T	1897	0	1886	33	0
7	G	1921	0	1909	31	0
7	U	1921	0	1910	44	0
8	H	1685	0	1688	16	0
8	V	1685	0	1688	24	0
9	I	1581	0	1574	16	0
9	W	1581	0	1574	21	0
10	J	1585	0	1590	16	0
10	X	1585	0	1590	15	0
11	K	1644	0	1595	26	0
11	Y	1644	0	1595	29	0
12	L	1757	0	1711	31	0
12	Z	1757	0	1711	44	0
13	1	1824	0	1832	20	0
13	M	1824	0	1832	23	0
14	2	1512	0	1481	21	0
14	N	1512	0	1481	24	0
15	D	1	0	0	0	0
15	F	1	0	0	0	0
15	G	2	0	0	0	0
15	H	1	0	0	0	0
15	I	2	0	0	0	0
15	K	1	0	0	0	0
15	L	1	0	0	0	0
15	N	1	0	0	0	0
16	K	45	0	44	0	0
16	Y	45	0	44	4	0
17	K	12	0	13	0	0
17	Y	12	0	13	1	0
All	All	49672	0	49367	657	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 657 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
12:Z:144(O):LYS:HB3	12:Z:144(R):LYS:HZ3	1.01	1.13
5:E:97:ASN:HD21	12:L:61:ASN:HD21	1.06	0.96
12:Z:144(O):LYS:HB3	12:Z:144(R):LYS:NZ	1.81	0.95
3:C:163:GLN:HE21	3:C:164:THR:H	0.97	0.94
8:H:128:GLY:O	8:H:131:SER:HB2	1.69	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	233 (94%)	12 (5%)	3 (1%)	13	43
1	O	248/250 (99%)	230 (93%)	15 (6%)	3 (1%)	13	43
2	B	242/244 (99%)	218 (90%)	22 (9%)	2 (1%)	19	53
2	P	242/244 (99%)	217 (90%)	19 (8%)	6 (2%)	5	26
3	C	239/241 (99%)	232 (97%)	5 (2%)	2 (1%)	19	53
3	Q	239/241 (99%)	223 (93%)	11 (5%)	5 (2%)	7	29
4	D	240/242 (99%)	224 (93%)	13 (5%)	3 (1%)	12	41
4	R	240/242 (99%)	220 (92%)	19 (8%)	1 (0%)	34	68
5	E	231/233 (99%)	218 (94%)	10 (4%)	3 (1%)	12	41
5	S	231/233 (99%)	213 (92%)	13 (6%)	5 (2%)	6	28
6	F	242/244 (99%)	229 (95%)	12 (5%)	1 (0%)	34	68
6	T	242/244 (99%)	226 (93%)	14 (6%)	2 (1%)	19	53
7	G	241/243 (99%)	229 (95%)	11 (5%)	1 (0%)	34	68
7	U	241/243 (99%)	227 (94%)	13 (5%)	1 (0%)	34	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
8	H	220/222 (99%)	206 (94%)	13 (6%)	1 (0%)	29	63
8	V	220/222 (99%)	207 (94%)	11 (5%)	2 (1%)	17	51
9	I	202/204 (99%)	195 (96%)	6 (3%)	1 (0%)	29	63
9	W	202/204 (99%)	191 (95%)	10 (5%)	1 (0%)	29	63
10	J	196/198 (99%)	183 (93%)	10 (5%)	3 (2%)	10	38
10	X	196/198 (99%)	181 (92%)	10 (5%)	5 (3%)	5	25
11	K	210/212 (99%)	199 (95%)	9 (4%)	2 (1%)	15	48
11	Y	210/212 (99%)	199 (95%)	8 (4%)	3 (1%)	11	39
12	L	220/222 (99%)	205 (93%)	14 (6%)	1 (0%)	29	63
12	Z	220/222 (99%)	204 (93%)	14 (6%)	2 (1%)	17	51
13	1	231/233 (99%)	215 (93%)	15 (6%)	1 (0%)	34	68
13	M	231/233 (99%)	218 (94%)	12 (5%)	1 (0%)	34	68
14	2	194/196 (99%)	185 (95%)	8 (4%)	1 (0%)	29	63
14	N	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
All	All	6312/6368 (99%)	5916 (94%)	334 (5%)	62 (1%)	15	48

5 of 62 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	167	LYS
4	D	123(G)	GLU
5	E	5	ARG
5	E	203	ASP
5	E	217	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	198 (95%)	11 (5%)	22	53
1	O	209/209 (100%)	199 (95%)	10 (5%)	25	57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	203/203 (100%)	179 (88%)	24 (12%)	5	21
2	P	203/203 (100%)	186 (92%)	17 (8%)	11	37
3	C	213/213 (100%)	200 (94%)	13 (6%)	18	49
3	Q	213/213 (100%)	193 (91%)	20 (9%)	8	32
4	D	198/198 (100%)	185 (93%)	13 (7%)	16	46
4	R	198/198 (100%)	179 (90%)	19 (10%)	8	30
5	E	192/192 (100%)	176 (92%)	16 (8%)	11	37
5	S	192/192 (100%)	171 (89%)	21 (11%)	6	25
6	F	201/201 (100%)	180 (90%)	21 (10%)	7	26
6	T	201/201 (100%)	183 (91%)	18 (9%)	9	33
7	G	207/207 (100%)	188 (91%)	19 (9%)	9	32
7	U	207/207 (100%)	193 (93%)	14 (7%)	16	45
8	H	181/181 (100%)	175 (97%)	6 (3%)	38	68
8	V	181/181 (100%)	170 (94%)	11 (6%)	18	49
9	I	172/172 (100%)	166 (96%)	6 (4%)	36	67
9	W	172/172 (100%)	164 (95%)	8 (5%)	26	58
10	J	175/175 (100%)	163 (93%)	12 (7%)	15	44
10	X	175/175 (100%)	166 (95%)	9 (5%)	24	55
11	K	169/169 (100%)	158 (94%)	11 (6%)	17	47
11	Y	169/169 (100%)	159 (94%)	10 (6%)	19	50
12	L	185/185 (100%)	175 (95%)	10 (5%)	22	53
12	Z	185/185 (100%)	173 (94%)	12 (6%)	17	47
13	1	199/199 (100%)	184 (92%)	15 (8%)	13	41
13	M	199/199 (100%)	187 (94%)	12 (6%)	19	49
14	2	162/162 (100%)	154 (95%)	8 (5%)	25	57
14	N	162/162 (100%)	158 (98%)	4 (2%)	47	75
All	All	5332/5332 (100%)	4962 (93%)	370 (7%)	15	44

5 of 370 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	R	52	LYS
7	U	87	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	R	156	THR
5	S	208(B)	ASP
8	V	68	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 146 such sidechains are listed below:

Mol	Chain	Res	Type
8	V	172	ASN
14	2	69	GLN
10	X	85	GLN
12	Z	144(B)	ASN
10	J	85	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
16	LXT	Y	212	-	48,48,48	0.72	0	62,62,62	0.90	2 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
17	MES	K	214	-	12,12,12	2.16	1 (8%)	14,16,16	1.38	1 (7%)
17	MES	Y	213	-	12,12,12	2.01	1 (8%)	14,16,16	1.69	2 (14%)
16	LXT	K	213	-	48,48,48	0.72	0	62,62,62	0.85	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	LXT	Y	212	-	-	14/40/40/40	0/4/4/4
17	MES	K	214	-	-	3/6/14/14	0/1/1/1
17	MES	Y	213	-	-	5/6/14/14	0/1/1/1
16	LXT	K	213	-	-	13/40/40/40	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
17	K	214	MES	C8-S	-7.12	1.67	1.77
17	Y	213	MES	C8-S	-6.57	1.68	1.77

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
17	Y	213	MES	O3S-S-C8	3.98	112.20	105.77
17	K	214	MES	O3S-S-C8	3.63	111.64	105.77
16	Y	212	LXT	C13-C12-N11	-3.39	105.79	113.05
17	Y	213	MES	O1S-S-C8	3.23	110.81	106.92
16	K	213	LXT	C13-C12-N11	-2.99	106.64	113.05

There are no chirality outliers.

5 of 35 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
16	Y	212	LXT	N2-C34-C36-N38
16	Y	212	LXT	O35-C34-C36-N38
16	Y	212	LXT	C34-C36-C37-C41
16	Y	212	LXT	N38-C36-C37-C41
17	K	214	MES	C7-C8-S-O1S


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
16	Y	212	LXT	4	0
17	Y	213	MES	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.31	3 (1%) 79 63	42, 61, 89, 102	0
1	O	250/250 (100%)	-0.26	6 (2%) 59 37	43, 62, 89, 102	0
2	B	244/244 (100%)	-0.17	3 (1%) 79 63	47, 67, 100, 107	0
2	P	244/244 (100%)	-0.18	3 (1%) 79 63	47, 67, 100, 108	0
3	C	241/241 (100%)	-0.04	4 (1%) 70 50	49, 69, 118, 142	0
3	Q	241/241 (100%)	0.10	14 (5%) 23 10	49, 70, 118, 142	0
4	D	242/242 (100%)	-0.07	9 (3%) 41 21	50, 69, 104, 113	0
4	R	242/242 (100%)	-0.10	7 (2%) 51 29	50, 69, 103, 113	0
5	E	233/233 (100%)	-0.07	6 (2%) 56 33	53, 74, 104, 111	0
5	S	233/233 (100%)	0.03	6 (2%) 56 33	53, 75, 104, 111	0
6	F	244/244 (100%)	-0.17	1 (0%) 92 85	46, 66, 97, 109	0
6	T	244/244 (100%)	-0.12	2 (0%) 86 74	46, 66, 97, 109	0
7	G	243/243 (100%)	-0.24	1 (0%) 92 85	44, 61, 84, 105	0
7	U	243/243 (100%)	-0.29	1 (0%) 92 85	45, 61, 83, 105	0
8	H	222/222 (100%)	-0.39	0 100 100	44, 57, 76, 107	0
8	V	222/222 (100%)	-0.42	1 (0%) 91 82	45, 57, 76, 107	0
9	I	204/204 (100%)	-0.43	0 100 100	44, 57, 69, 77	0
9	W	204/204 (100%)	-0.38	0 100 100	43, 56, 69, 77	0
10	J	198/198 (100%)	-0.32	2 (1%) 82 69	43, 59, 72, 115	0
10	X	198/198 (100%)	-0.21	3 (1%) 73 56	43, 59, 73, 115	0
11	K	212/212 (100%)	-0.28	0 100 100	45, 60, 78, 89	0
11	Y	212/212 (100%)	-0.37	0 100 100	44, 60, 79, 89	0
12	L	222/222 (100%)	-0.37	1 (0%) 91 82	40, 57, 84, 90	0
12	Z	222/222 (100%)	-0.27	0 100 100	41, 57, 84, 90	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	1	233/233 (100%)	-0.32	1 (0%) 92 85	42, 55, 72, 82	0
13	M	233/233 (100%)	-0.42	1 (0%) 92 85	42, 55, 73, 83	0
14	2	196/196 (100%)	-0.38	1 (0%) 91 82	45, 53, 72, 79	0
14	N	196/196 (100%)	-0.38	0 100 100	45, 53, 72, 79	0
All	All	6368/6368 (100%)	-0.24	76 (1%) 79 63	40, 61, 97, 142	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	123(A)	GLY	7.3
4	D	123(C)	GLY	6.9
13	1	-8	THR	6.8
4	R	123(E)	SER	6.3
4	R	123(F)	GLY	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	MG	F	242	1/1	0.71	0.41	60,60,60,60	0
15	MG	G	241	1/1	0.72	0.83	85,85,85,85	0
16	LXT	Y	212	45/45	0.85	0.31	78,84,89,90	0
16	LXT	K	213	45/45	0.87	0.31	66,77,88,88	0
15	MG	I	196	1/1	0.90	0.18	57,57,57,57	0
15	MG	N	188	1/1	0.92	0.23	49,49,49,49	0
15	MG	H	224	1/1	0.92	0.12	56,56,56,56	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
15	MG	D	2	1/1	0.92	0.08	55,55,55,55	0
15	MG	K	212	1/1	0.93	0.17	56,56,56,56	0
17	MES	Y	213	12/12	0.93	0.23	90,91,92,92	0
15	MG	I	195	1/1	0.94	0.34	53,53,53,53	0
17	MES	K	214	12/12	0.96	0.17	84,86,87,87	0
15	MG	L	195	1/1	0.96	0.28	58,58,58,58	0
15	MG	G	1	1/1	0.98	0.06	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.