



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 06:32 PM EDT

PDB ID : 4MHX
Title : Crystal Structure of Sulfamidase
Authors : Sidhu, N.S.; Uson, I.; Schreiber, K.; Proepper, K.; Becker, S.; Gaertner, J.;
Kraetzner, R.; Steinfeld, R.; Sheldrick, G.M.
Deposited on : 2013-08-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	510	 2% 87% 7% • 5%
1	B	510	 2% 86% 7% • 5%
2	C	2	 100%
2	D	2	 100%
2	E	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-sulphoglucosamine sulphonydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	483	Total	C	N	O	P	S	0	0	0
			3792	2441	655	684	1	11			
1	B	482	Total	C	N	O	P	S	0	2	0
			3806	2445	654	695	1	11			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	FGP	CYS	modified residue	UNP P51688
A	503	ARG	-	expression tag	UNP P51688
A	504	SER	-	expression tag	UNP P51688
A	505	HIS	-	expression tag	UNP P51688
A	506	HIS	-	expression tag	UNP P51688
A	507	HIS	-	expression tag	UNP P51688
A	508	HIS	-	expression tag	UNP P51688
A	509	HIS	-	expression tag	UNP P51688
A	510	HIS	-	expression tag	UNP P51688
B	70	FGP	CYS	modified residue	UNP P51688
B	503	ARG	-	expression tag	UNP P51688
B	504	SER	-	expression tag	UNP P51688
B	505	HIS	-	expression tag	UNP P51688
B	506	HIS	-	expression tag	UNP P51688
B	507	HIS	-	expression tag	UNP P51688
B	508	HIS	-	expression tag	UNP P51688
B	509	HIS	-	expression tag	UNP P51688
B	510	HIS	-	expression tag	UNP P51688

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Mg	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	95	Total	O	0	0
			95	95		
6	B	110	Total	O	0	0
			110	110		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-sulphoglucosamine sulphohydrolase



- Molecule 1: N-sulphoglucosamine sulphohydrolase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.38Å 107.90Å 79.79Å 90.00° 104.07° 90.00°	Depositor
Resolution (Å)	44.30 – 2.00 44.26 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.30-2.00) 99.9 (44.26-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 2.00Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.192 , 0.230 0.197 , 0.230	Depositor DCC
R_{free} test set	3447 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtrriage
Anisotropy	0.388	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7975	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG, FGP, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/3902	0.76	4/5334 (0.1%)
1	B	0.66	0/3922	0.74	2/5361 (0.0%)
All	All	0.64	0/7824	0.75	6/10695 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	88	MET	CG-SD-CE	-13.29	78.93	100.20
1	A	154	ARG	NE-CZ-NH1	5.97	123.28	120.30
1	B	245	ARG	NE-CZ-NH2	-5.59	117.51	120.30
1	A	77	LEU	CA-CB-CG	5.36	127.63	115.30
1	B	428	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	A	441	ARG	NE-CZ-NH1	5.07	122.83	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	70	FGP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3792	0	3594	19	0
1	B	3806	0	3584	22	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
2	E	28	0	25	0	0
2	F	28	0	25	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	28	0	26	1	0
4	B	28	0	26	0	0
5	B	2	0	0	0	0
6	A	95	0	0	1	0
6	B	110	0	0	4	0
All	All	7975	0	7330	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:THR:HG22	6:B:809:HOH:O	1.95	0.67
1:A:61:ARG:HD2	1:A:310:GLU:OE2	1.99	0.63
1:B:385:ARG:HD3	1:B:447:GLU:OE2	2.00	0.61
1:B:43:SER:CB	6:B:765:HOH:O	2.50	0.60
1:B:61:ARG:HD2	1:B:310:GLU:OE2	2.03	0.58
1:B:357:LEU:HD12	1:B:357:LEU:O	2.04	0.57
1:B:179:ASP:OD2	1:B:245:ARG:HD2	2.05	0.57
1:B:366[B]:SER:OG	1:B:367:HIS:N	2.38	0.57
1:A:40:TYR:HA	4:A:602:NAG:H82	1.88	0.56
1:A:357:LEU:HD12	1:A:357:LEU:O	2.05	0.55
1:A:271:THR:HG21	1:A:315:LEU:HD11	1.89	0.54
1:B:156:LYS:O	1:B:159[A]:VAL:HG12	2.07	0.54
1:B:287:TRP:HB3	1:B:288:PRO:HD3	1.92	0.51
1:A:34:GLY:HA2	6:A:778:HOH:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:276:ILE:HD12	1:B:278:PHE:CD2	2.47	0.50
1:A:31:ASP:O	1:A:32:ASP:HB2	2.12	0.49
1:B:31:ASP:O	1:B:32:ASP:HB2	2.12	0.49
1:A:285:LEU:O	1:A:385:ARG:NH2	2.47	0.48
1:A:276:ILE:HD12	1:A:278:PHE:CD2	2.49	0.47
1:A:140:GLU:HB3	1:A:145:VAL:HG12	1.97	0.47
1:B:105:ARG:N	1:B:105:ARG:HD2	2.30	0.47
1:B:189:GLN:OE1	1:B:189:GLN:N	2.44	0.46
1:A:304:ARG:HG3	1:A:307:GLN:CD	2.36	0.46
1:B:69:SER:O	1:B:70:FGP:C	2.64	0.46
1:B:32:ASP:OD1	1:B:274:ASN:ND2	2.44	0.44
1:A:271:THR:HG22	1:A:272:SER:H	1.83	0.44
1:A:77:LEU:CD2	1:A:77:LEU:C	2.86	0.43
1:A:287:TRP:HB3	1:A:288:PRO:HD3	2.00	0.43
1:B:279:PRO:O	1:B:280:SER:HB2	2.19	0.42
1:B:385:ARG:HD2	1:B:441:ARG:HD2	2.02	0.42
1:A:77:LEU:C	1:A:77:LEU:HD23	2.38	0.42
1:B:55:ARG:HD2	6:B:776:HOH:O	2.18	0.42
1:B:426:ASP:HB2	6:B:784:HOH:O	2.20	0.41
1:A:199:ASN:OD1	1:A:199:ASN:C	2.59	0.41
1:B:64:PHE:HB2	1:B:361:VAL:HG11	2.02	0.41
1:A:140:GLU:HB3	1:A:145:VAL:CG1	2.50	0.41
1:A:380:GLN:OE1	1:A:385:ARG:NH1	2.53	0.41
1:A:317:ASP:C	1:A:320:PRO:HD2	2.41	0.41
1:B:196:LYS:O	1:B:204:MET:HB3	2.22	0.40
1:A:372:MET:HG2	1:A:397:PRO:HB2	2.03	0.40
1:B:287:TRP:N	1:B:288:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	480/510 (94%)	467 (97%)	12 (2%)	1 (0%)	47	44
1	B	481/510 (94%)	466 (97%)	14 (3%)	1 (0%)	47	44
All	All	961/1020 (94%)	933 (97%)	26 (3%)	2 (0%)	47	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	94	ASP
1	B	94	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/442 (88%)	379 (97%)	12 (3%)	40	40
1	B	394/442 (89%)	382 (97%)	12 (3%)	41	41
All	All	785/884 (89%)	761 (97%)	24 (3%)	40	40

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	LEU
1	A	94	ASP
1	A	95	VAL
1	A	109	LEU
1	A	235	ASP
1	A	271	THR
1	A	274	ASN
1	A	304	ARG
1	A	346	ARG
1	A	391	ASN
1	A	424	TYR
1	A	491	LEU
1	B	56	ARG
1	B	94	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	95	VAL
1	B	105	ARG
1	B	109	LEU
1	B	129	GLU
1	B	192	THR
1	B	274	ASN
1	B	332	SER
1	B	391	ASN
1	B	424	TYR
1	B	453	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	400	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	FGP	A	70	1,3	7,10,11	0.87	0	9,14,16	2.32	1 (11%)
1	FGP	B	70	1,3	7,10,11	1.14	0	9,14,16	2.68	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FGP	A	70	1,3	-	2/5/11/13	-
1	FGP	B	70	1,3	-	1/5/11/13	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	70	FGP	O-C-CA	-6.80	107.28	124.77
1	A	70	FGP	O-C-CA	-6.04	109.25	124.77
1	B	70	FGP	O3P-P-O2P	2.83	118.41	107.80
1	B	70	FGP	P-OG2-CB	-2.55	116.61	123.43

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	70	FGP	CA-CB-OG2-P
1	B	70	FGP	CA-CB-OG2-P
1	A	70	FGP	N-CA-CB-OG1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	70	FGP	1	0

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.71	0	17,19,21	1.22	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	2	2	14,14,15	0.58	0	17,19,21	1.57	4 (23%)
2	NAG	D	1	1,2	14,14,15	0.53	0	17,19,21	1.40	3 (17%)
2	NAG	D	2	2	14,14,15	0.59	0	17,19,21	1.46	4 (23%)
2	NAG	E	1	1,2	14,14,15	0.83	0	17,19,21	1.73	3 (17%)
2	NAG	E	2	2	14,14,15	0.65	0	17,19,21	0.97	1 (5%)
2	NAG	F	1	1,2	14,14,15	0.81	1 (7%)	17,19,21	1.65	3 (17%)
2	NAG	F	2	2	14,14,15	0.67	0	17,19,21	1.71	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	1	NAG	O5-C5	-2.31	1.38	1.43

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	NAG	C1-O5-C5	4.97	118.85	112.19
2	F	2	NAG	C3-C4-C5	4.36	118.13	110.23
2	C	2	NAG	O5-C1-C2	-3.71	105.55	111.29
2	F	1	NAG	C3-C4-C5	3.64	116.84	110.23
2	F	2	NAG	C4-C3-C2	3.62	116.33	111.02
2	F	1	NAG	O3-C3-C2	-3.55	102.03	109.40
2	D	1	NAG	C1-O5-C5	3.49	116.86	112.19
2	E	1	NAG	C1-C2-N2	-3.39	105.10	110.43
2	D	2	NAG	O4-C4-C5	3.07	116.89	109.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	NAG	C1-O5-C5	2.98	116.19	112.19
2	D	2	NAG	C2-N2-C7	2.80	126.65	122.90
2	D	1	NAG	C1-C2-N2	2.73	114.73	110.43
2	D	1	NAG	O5-C1-C2	-2.66	107.18	111.29
2	C	2	NAG	C2-N2-C7	-2.45	119.62	122.90
2	E	1	NAG	O7-C7-N2	2.42	126.26	121.98
2	C	1	NAG	O7-C7-C8	-2.40	117.78	122.05
2	E	2	NAG	O7-C7-C8	-2.37	117.83	122.05
2	D	2	NAG	C3-C4-C5	-2.33	106.00	110.23
2	C	1	NAG	C1-C2-N2	2.15	113.83	110.43
2	C	2	NAG	C4-C3-C2	-2.04	108.03	111.02
2	D	2	NAG	O7-C7-N2	2.04	125.58	121.98
2	F	2	NAG	O5-C1-C2	-2.03	108.15	111.29
2	C	2	NAG	C1-O5-C5	2.02	114.90	112.19

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	NAG	C1-C2-N2-C7
2	D	2	NAG	O5-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	F	2	NAG	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	607	1	14,14,15	0.82	1 (7%)	17,19,21	1.61	5 (29%)
4	NAG	B	607	1	14,14,15	0.56	0	17,19,21	1.74	3 (17%)
4	NAG	A	602	1	14,14,15	0.64	0	17,19,21	2.11	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	602	1	14,14,15	0.66	0	17,19,21	1.40	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	607	1	-	2/6/23/26	0/1/1/1
4	NAG	B	607	1	-	1/6/23/26	0/1/1/1
4	NAG	A	602	1	-	2/6/23/26	0/1/1/1
4	NAG	B	602	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	607	NAG	C1-C2	2.33	1.55	1.52

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	607	NAG	C1-O5-C5	5.21	119.16	112.19
4	A	602	NAG	C2-N2-C7	3.99	128.25	122.90
4	B	602	NAG	C1-C2-N2	-3.71	104.59	110.43
4	A	602	NAG	C1-C2-N2	-3.56	104.82	110.43
4	A	602	NAG	C1-O5-C5	3.49	116.87	112.19
4	A	602	NAG	O7-C7-C8	-3.38	116.03	122.05
4	A	602	NAG	O7-C7-N2	3.17	127.59	121.98
4	A	607	NAG	O3-C3-C2	3.00	115.63	109.40
4	B	607	NAG	C1-C2-N2	2.94	115.06	110.43
4	A	607	NAG	C1-O5-C5	2.67	115.76	112.19
4	B	602	NAG	C2-N2-C7	-2.48	119.58	122.90
4	A	607	NAG	O7-C7-C8	-2.48	117.65	122.05
4	A	602	NAG	O5-C1-C2	2.28	114.82	111.29
4	A	607	NAG	O4-C4-C5	2.20	114.74	109.32
4	B	602	NAG	O5-C1-C2	-2.08	108.07	111.29
4	B	607	NAG	O3-C3-C2	-2.03	105.18	109.40
4	A	607	NAG	C3-C4-C5	-2.02	106.57	110.23

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	607	NAG	C4-C5-C6-O6
4	A	607	NAG	O5-C5-C6-O6
4	A	602	NAG	C4-C5-C6-O6
4	B	602	NAG	C4-C5-C6-O6
4	B	602	NAG	O5-C5-C6-O6
4	A	602	NAG	O5-C5-C6-O6
4	B	607	NAG	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	602	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	482/510 (94%)	-0.04	11 (2%) 60 59	23, 44, 65, 92	0
1	B	481/510 (94%)	-0.05	10 (2%) 63 62	22, 42, 65, 107	0
All	All	963/1020 (94%)	-0.04	21 (2%) 62 60	22, 43, 65, 107	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	492	SER	6.5
1	B	492	SER	5.8
1	A	418	GLY	4.7
1	B	493	PRO	4.3
1	A	491	LEU	3.6
1	B	338	SER	3.5
1	B	491	LEU	3.1
1	B	455	PRO	3.0
1	B	454	ASP	3.0
1	A	202	SER	2.9
1	B	458	ALA	2.9
1	B	453	THR	2.8
1	A	419	GLN	2.8
1	A	493	PRO	2.7
1	A	416	THR	2.7
1	B	503	ARG	2.5
1	B	502	LEU	2.4
1	A	371	THR	2.2
1	A	417	ALA	2.2
1	A	90	GLY	2.1
1	A	91	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	FGP	A	70	11/12	0.92	0.14	29,40,76,84	0
1	FGP	B	70	11/12	0.93	0.13	24,35,69,75	0

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.79	0.28	50,72,78,81	0
2	NAG	D	2	14/15	0.79	0.29	62,75,77,77	0
2	NAG	D	1	14/15	0.82	0.29	65,75,79,80	0
2	NAG	E	2	14/15	0.89	0.25	59,72,80,82	0
2	NAG	F	2	14/15	0.90	0.21	77,79,85,86	0
2	NAG	F	1	14/15	0.92	0.10	48,56,64,66	0
2	NAG	C	1	14/15	0.96	0.11	37,43,47,61	0
2	NAG	E	1	14/15	0.97	0.09	34,41,48,55	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	602	14/15	0.82	0.17	52,60,65,65	0
4	NAG	B	602	14/15	0.84	0.13	51,71,79,80	0
4	NAG	A	607	14/15	0.88	0.21	64,73,81,83	0
5	MG	B	608	1/1	0.88	0.35	72,72,72,72	0
4	NAG	B	607	14/15	0.89	0.21	46,55,67,68	0
5	MG	B	609	1/1	0.98	0.08	53,53,53,53	0
3	CA	A	601	1/1	0.99	0.05	46,46,46,46	0
3	CA	B	601	1/1	0.99	0.07	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.