



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 15, 2024 – 12:09 PM EST

PDB ID : 9MI2  
Title : Crystal structure of Purine nucleoside phosphorylase from *Trichomonas vaginalis* (C2 form)  
Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2024-12-12  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	91% 5% .
1	B	244	91% 6% .
1	C	244	90% 6% .
1	D	244	91% 6% .
1	E	244	91% 5% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	244	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a small red segment at the beginning labeled '2%', a large green segment in the middle labeled '91%', and a small yellow segment at the end labeled '5%'. The bar is contained within a rectangular frame.</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11678 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Purine nucleoside phosphorylase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	236	1798	1135	304	345	14	0	3	0
1	B	236	1806	1142	303	346	15	0	4	0
1	C	236	1806	1140	303	348	15	0	5	0
1	D	236	1821	1150	307	349	15	0	7	0
1	E	236	1790	1129	304	344	13	0	1	0
1	F	236	1760	1111	296	340	13	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP A2EU62
A	-6	ALA	-	expression tag	UNP A2EU62
A	-5	HIS	-	expression tag	UNP A2EU62
A	-4	HIS	-	expression tag	UNP A2EU62
A	-3	HIS	-	expression tag	UNP A2EU62
A	-2	HIS	-	expression tag	UNP A2EU62
A	-1	HIS	-	expression tag	UNP A2EU62
A	0	HIS	-	expression tag	UNP A2EU62
B	-7	MET	-	initiating methionine	UNP A2EU62
B	-6	ALA	-	expression tag	UNP A2EU62
B	-5	HIS	-	expression tag	UNP A2EU62
B	-4	HIS	-	expression tag	UNP A2EU62
B	-3	HIS	-	expression tag	UNP A2EU62
B	-2	HIS	-	expression tag	UNP A2EU62
B	-1	HIS	-	expression tag	UNP A2EU62
B	0	HIS	-	expression tag	UNP A2EU62
C	-7	MET	-	initiating methionine	UNP A2EU62

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-6	ALA	-	expression tag	UNP A2EU62
C	-5	HIS	-	expression tag	UNP A2EU62
C	-4	HIS	-	expression tag	UNP A2EU62
C	-3	HIS	-	expression tag	UNP A2EU62
C	-2	HIS	-	expression tag	UNP A2EU62
C	-1	HIS	-	expression tag	UNP A2EU62
C	0	HIS	-	expression tag	UNP A2EU62
D	-7	MET	-	initiating methionine	UNP A2EU62
D	-6	ALA	-	expression tag	UNP A2EU62
D	-5	HIS	-	expression tag	UNP A2EU62
D	-4	HIS	-	expression tag	UNP A2EU62
D	-3	HIS	-	expression tag	UNP A2EU62
D	-2	HIS	-	expression tag	UNP A2EU62
D	-1	HIS	-	expression tag	UNP A2EU62
D	0	HIS	-	expression tag	UNP A2EU62
E	-7	MET	-	initiating methionine	UNP A2EU62
E	-6	ALA	-	expression tag	UNP A2EU62
E	-5	HIS	-	expression tag	UNP A2EU62
E	-4	HIS	-	expression tag	UNP A2EU62
E	-3	HIS	-	expression tag	UNP A2EU62
E	-2	HIS	-	expression tag	UNP A2EU62
E	-1	HIS	-	expression tag	UNP A2EU62
E	0	HIS	-	expression tag	UNP A2EU62
F	-7	MET	-	initiating methionine	UNP A2EU62
F	-6	ALA	-	expression tag	UNP A2EU62
F	-5	HIS	-	expression tag	UNP A2EU62
F	-4	HIS	-	expression tag	UNP A2EU62
F	-3	HIS	-	expression tag	UNP A2EU62
F	-2	HIS	-	expression tag	UNP A2EU62
F	-1	HIS	-	expression tag	UNP A2EU62
F	0	HIS	-	expression tag	UNP A2EU62

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	E	1	Total C O 6 3 3	0	0
2	F	1	Total C O 6 3 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	155	Total O 155 155	0	0
3	B	188	Total O 188 188	0	0
3	C	130	Total O 130 130	0	0
3	D	188	Total O 188 188	0	0
3	E	105	Total O 105 105	0	0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	F	89	Total	O	0	0
			89	89		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Purine nucleoside phosphorylase, putative

Chain A:  91% 5%




- Molecule 1: Purine nucleoside phosphorylase, putative

Chain B:  91% 6%



- Molecule 1: Purine nucleoside phosphorylase, putative

Chain C:  90% 6%




- Molecule 1: Purine nucleoside phosphorylase, putative

Chain D:  91% 6%




- Molecule 1: Purine nucleoside phosphorylase, putative


Chain E:  91% 5%



- Molecule 1: Purine nucleoside phosphorylase, putative



Chain F:  2% 91% 5%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	170.27Å 50.36Å 175.49Å 90.00° 98.82° 90.00°	Depositor
Resolution (Å)	48.25 – 1.80 48.25 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.25-1.80) 99.8 (48.25-1.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 1.79Å)	Xtrriage
Refinement program	PHENIX dev_5533	Depositor
R, $R_{free}$	0.164 , 0.197 0.172 , 0.202	Depositor DCC
$R_{free}$ test set	6878 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.1	Xtrriage
Anisotropy	0.620	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11678	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/1840	0.67	0/2495
1	B	0.52	0/1851	0.67	0/2509
1	C	0.47	0/1854	0.66	0/2513
1	D	0.50	0/1875	0.66	0/2539
1	E	0.44	0/1826	0.62	0/2477
1	F	0.44	0/1793	0.61	0/2438
All	All	0.48	0/11039	0.65	0/14971

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	25	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1798	0	1793	8	0
1	B	1806	0	1809	13	0
1	C	1806	0	1806	11	0
1	D	1821	0	1835	8	0
1	E	1790	0	1779	9	0
1	F	1760	0	1721	11	0
2	A	6	0	8	0	0
2	B	12	0	16	0	0
2	C	6	0	8	0	0
2	D	6	0	8	0	0
2	E	6	0	8	0	0
2	F	6	0	8	0	0
3	A	155	0	0	0	0
3	B	188	0	0	2	0
3	C	130	0	0	2	0
3	D	188	0	0	0	0
3	E	105	0	0	0	0
3	F	89	0	0	0	0
All	All	11678	0	10799	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:VAL:CG1	1:A:213:ALA:HB2	2.17	0.74
1:B:89:VAL:HB	1:B:225[B]:MET:HE1	1.76	0.67
1:A:101:VAL:HG12	1:A:213:ALA:HB2	1.82	0.61
1:F:25:ARG:O	1:F:29:ILE:HD12	2.01	0.61
1:B:235:LYS:HD2	3:B:433:HOH:O	2.01	0.59
1:C:225[A]:MET:HE1	3:C:445:HOH:O	2.03	0.58
1:D:229:ALA:O	1:D:233:THR:HG23	2.04	0.56
1:E:8:ALA:HB2	1:E:40:VAL:HG22	1.86	0.56
1:E:12:ASP:HB3	1:E:40:VAL:HG21	1.88	0.55
1:B:235:LYS:CD	3:B:433:HOH:O	2.55	0.54
1:C:95:VAL:HG22	3:C:472:HOH:O	2.08	0.54
1:B:89:VAL:HB	1:B:225[B]:MET:CE	2.37	0.53
1:E:8:ALA:CB	1:E:40:VAL:HG22	2.39	0.52
1:B:20:CYS:O	1:B:62:GLY:HA2	2.10	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:101:VAL:HG13	1:B:213:ALA:HB2	1.91	0.51
1:E:229:ALA:O	1:E:233:THR:HG23	2.11	0.51
1:B:101:VAL:O	1:B:102:ARG:CB	2.60	0.48
1:B:101:VAL:CG1	1:B:213:ALA:HB2	2.43	0.48
1:B:18:LEU:HB2	1:B:60[B]:VAL:HG12	1.96	0.48
1:F:122:LEU:N	1:F:122:LEU:HD12	2.29	0.48
1:F:29:ILE:HD11	1:F:222:PHE:HE1	1.79	0.47
1:D:82:LYS:O	1:D:84:LYS:HE3	2.15	0.47
1:A:39:GLN:OE1	1:A:42:SER:HB3	2.16	0.46
1:D:20:CYS:O	1:D:62:GLY:HA2	2.14	0.46
1:F:18:LEU:O	1:F:60:VAL:HA	2.15	0.46
1:E:20:CYS:O	1:E:62:GLY:HA2	2.15	0.46
1:C:20:CYS:O	1:C:62:GLY:HA2	2.16	0.46
1:E:61:MET:HE1	1:E:77:LEU:HD21	1.98	0.46
1:D:101:VAL:O	1:D:102:ARG:CB	2.64	0.45
1:A:20:CYS:O	1:A:62:GLY:HA2	2.16	0.45
1:E:12:ASP:HB3	1:E:40:VAL:CG2	2.47	0.45
1:C:225[B]:MET:HG3	1:C:226:ILE:N	2.33	0.44
1:C:101:VAL:CG1	1:C:213:ALA:HB2	2.47	0.44
1:F:90:GLY:O	1:F:225:MET:HE1	2.18	0.44
1:A:115[A]:ASN:ND2	1:B:115:ASN:OD1	2.50	0.43
1:C:90:GLY:O	1:C:201[B]:CYS:HA	2.18	0.43
1:F:89:VAL:HG23	1:F:89:VAL:O	2.18	0.43
1:C:89:VAL:HB	1:C:225[B]:MET:HE1	1.99	0.43
1:E:61:MET:CE	1:E:77:LEU:HD21	2.48	0.43
1:C:89:VAL:CG1	1:C:225[B]:MET:HE1	2.49	0.42
1:D:111:GLY:CA	1:F:127:PRO:HD3	2.49	0.42
1:D:127:PRO:HD3	1:F:111:GLY:HA2	2.01	0.42
1:A:104:VAL:HG22	1:A:202:THR:HG22	2.02	0.42
1:C:214:THR:O	1:C:218:ARG:HG3	2.20	0.42
1:B:110[A]:SER:HA	1:B:154:ALA:O	2.20	0.42
1:C:18:LEU:O	1:C:60:VAL:HA	2.20	0.42
1:D:89:VAL:O	1:D:89:VAL:HG23	2.20	0.42
1:F:29:ILE:HD11	1:F:222:PHE:CE1	2.54	0.42
1:F:90:GLY:O	1:F:225:MET:CE	2.67	0.42
1:B:89:VAL:HG23	1:B:89:VAL:O	2.20	0.41
1:B:110[B]:SER:HA	1:B:154:ALA:O	2.20	0.41
1:C:90:GLY:O	1:C:201[A]:CYS:HA	2.21	0.41
1:D:127:PRO:HD3	1:F:111:GLY:CA	2.51	0.41
1:A:122:LEU:HD12	1:A:122:LEU:N	2.36	0.40
1:A:119[A]:MET:SD	1:E:124:HIS:HE1	2.43	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/244 (97%)	232 (98%)	5 (2%)	0	100	100
1	B	238/244 (98%)	234 (98%)	4 (2%)	0	100	100
1	C	239/244 (98%)	236 (99%)	3 (1%)	0	100	100
1	D	241/244 (99%)	236 (98%)	5 (2%)	0	100	100
1	E	235/244 (96%)	230 (98%)	5 (2%)	0	100	100
1	F	234/244 (96%)	230 (98%)	4 (2%)	0	100	100
All	All	1424/1464 (97%)	1398 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/201 (97%)	190 (97%)	5 (3%)	41	29
1	B	197/201 (98%)	195 (99%)	2 (1%)	73	68
1	C	198/201 (98%)	194 (98%)	4 (2%)	50	40
1	D	201/201 (100%)	198 (98%)	3 (2%)	60	53
1	E	193/201 (96%)	190 (98%)	3 (2%)	58	50
1	F	186/201 (92%)	183 (98%)	3 (2%)	58	50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1170/1206 (97%)	1150 (98%)	20 (2%)	58 47

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119[A]	MET
1	A	119[B]	MET
1	A	126	PHE
1	A	180	GLU
1	A	211	GLU
1	B	126	PHE
1	B	180	GLU
1	C	126	PHE
1	C	180	GLU
1	C	201[A]	CYS
1	C	201[B]	CYS
1	D	88	ARG
1	D	126	PHE
1	D	180	GLU
1	E	28	LEU
1	E	126	PHE
1	E	180	GLU
1	F	25	ARG
1	F	126	PHE
1	F	180	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	C	301	-	5,5,5	0.47	0	5,5,5	0.60	0
2	GOL	E	301	-	5,5,5	0.31	0	5,5,5	0.41	0
2	GOL	F	301	-	5,5,5	0.39	0	5,5,5	0.42	0
2	GOL	B	301	-	5,5,5	0.41	0	5,5,5	0.28	0
2	GOL	D	301	-	5,5,5	0.65	0	5,5,5	0.99	0
2	GOL	B	302	-	5,5,5	0.35	0	5,5,5	0.25	0
2	GOL	A	301	-	5,5,5	0.45	0	5,5,5	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	301	-	-	4/4/4/4	-
2	GOL	E	301	-	-	4/4/4/4	-
2	GOL	F	301	-	-	0/4/4/4	-
2	GOL	B	301	-	-	1/4/4/4	-
2	GOL	D	301	-	-	2/4/4/4	-
2	GOL	B	302	-	-	4/4/4/4	-
2	GOL	A	301	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	GOL	O1-C1-C2-C3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	302	GOL	O1-C1-C2-C3
2	C	301	GOL	O1-C1-C2-C3
2	E	301	GOL	O1-C1-C2-C3
2	E	301	GOL	C1-C2-C3-O3
2	B	302	GOL	O2-C2-C3-O3
2	D	301	GOL	O1-C1-C2-O2
2	B	302	GOL	C1-C2-C3-O3
2	D	301	GOL	O1-C1-C2-C3
2	A	301	GOL	O1-C1-C2-O2
2	B	302	GOL	O1-C1-C2-O2
2	C	301	GOL	O1-C1-C2-O2
2	E	301	GOL	O1-C1-C2-O2
2	C	301	GOL	O2-C2-C3-O3
2	E	301	GOL	O2-C2-C3-O3
2	C	301	GOL	C1-C2-C3-O3
2	B	301	GOL	O2-C2-C3-O3
2	A	301	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	236/244 (96%)	-0.10	1 (0%) 89 88	13, 29, 51, 56	3 (1%)
1	B	236/244 (96%)	-0.27	1 (0%) 89 88	13, 24, 39, 61	4 (1%)
1	C	236/244 (96%)	-0.10	0 100 100	14, 30, 49, 59	5 (2%)
1	D	236/244 (96%)	-0.22	0 100 100	14, 25, 40, 59	7 (2%)
1	E	236/244 (96%)	0.17	1 (0%) 89 88	15, 39, 58, 70	1 (0%)
1	F	236/244 (96%)	0.39	6 (2%) 58 57	20, 44, 64, 71	0
All	All	1416/1464 (96%)	-0.02	9 (0%) 85 85	13, 30, 55, 71	20 (1%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	10	VAL	3.2
1	F	1	MET	3.1
1	F	38	LYS	3.1
1	E	38	LYS	2.5
1	F	36	ASN	2.5
1	B	211	GLU	2.4
1	A	119[A]	MET	2.1
1	F	208	ILE	2.1
1	F	37	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	B	302	6/6	0.82	0.14	41,45,51,56	0
2	GOL	E	301	6/6	0.82	0.13	39,52,55,63	0
2	GOL	F	301	6/6	0.83	0.15	40,58,59,61	0
2	GOL	D	301	6/6	0.87	0.13	27,34,43,49	0
2	GOL	A	301	6/6	0.90	0.10	30,37,39,50	0
2	GOL	B	301	6/6	0.90	0.10	33,40,43,51	0
2	GOL	C	301	6/6	0.93	0.09	27,36,39,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.