



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2019 – 01:03 PM EST

PDB ID : 6MLC  
Title : PHD6 domain of MLL3 in complex with histone H4  
Authors : Dong, A.; Liu, Y.; Qin, S.; Lei, M.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.M.; Min, J.; Structural Genomics Consortium (SGC)  
Deposited on : 2018-09-27  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031633  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031633


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	5253 (1.80-1.80)
Clashscore	122126	6077 (1.80-1.80)
Ramachandran outliers	120053	6011 (1.80-1.80)
Sidechain outliers	120020	6010 (1.80-1.80)
RSRZ outliers	108989	5157 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	92	 4% 83% 8% • 8%
1	B	92	 3% 90% •• 8%
1	C	92	 3% 80% 11% • 8%
1	D	92	 82% 8% • 9%
2	E	20	 30% 70%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	20	 30% 70%

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 3059 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase 2C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	85	Total 700	C 420	N 130	O 135	S 15	0	4	1
1	B	85	Total 701	C 421	N 133	O 133	S 14	0	3	0
1	C	85	Total 690	C 415	N 127	O 133	S 15	0	2	0
1	D	84	Total 675	C 407	N 122	O 131	S 15	0	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1053	GLY	-	expression tag	UNP Q8NEZ4
A	1054	SER	-	expression tag	UNP Q8NEZ4
B	1053	GLY	-	expression tag	UNP Q8NEZ4
B	1054	SER	-	expression tag	UNP Q8NEZ4
C	1053	GLY	-	expression tag	UNP Q8NEZ4
C	1054	SER	-	expression tag	UNP Q8NEZ4
D	1053	GLY	-	expression tag	UNP Q8NEZ4
D	1054	SER	-	expression tag	UNP Q8NEZ4


- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	6	Total 49	C 29	N 14	O 6	0	0	0
2	F	6	Total 50	C 29	N 15	O 6	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Zn 2 2	0	0
3	A	5	Total Zn 5 5	0	0
3	D	2	Total Zn 2 2	0	0
3	C	5	Total Zn 5 5	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0

- Molecule 5 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	2	Total X 2 2	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total X 2 2	0	0
5	D	1	Total X 1 1	0	0
5	C	1	Total X 1 1	0	0
5	E	1	Total X 1 1	0	0



- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	28	Total O 30 30	0	2
6	B	36	Total O 36 36	0	0
6	C	40	Total O 41 41	0	1
6	D	33	Total O 34 34	0	1
6	F	2	Total O 2 2	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Histone-lysine N-methyltransferase 2C




- Molecule 1: Histone-lysine N-methyltransferase 2C




- Molecule 1: Histone-lysine N-methyltransferase 2C



- Molecule 1: Histone-lysine N-methyltransferase 2C



- Molecule 2: Histone H4



- Molecule 2: Histone H4



SER	GLY	ARG	GLY	LYS	GLY	GLY	LYS	LEU	GLY	LYS	GLY	GLU14	R1519	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-------	-------	-----



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.94Å 85.94Å 98.73Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.00 – 1.80 42.97 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.00-1.80) 100.0 (42.97-1.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.8.0232	Depositor
R, $R_{free}$	0.194 , 0.216 0.200 , 0.218	Depositor DCC
$R_{free}$ test set	1512 reflections (3.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.8	Xtrriage
Anisotropy	0.154	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 51.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.052 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3059	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.92 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4148e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	0/713	0.75	0/965
1	B	0.75	0/718	0.70	0/972
1	C	0.77	0/704	0.73	0/954
1	D	0.77	0/688	0.72	0/932
2	E	0.74	0/49	0.92	0/62
2	F	0.65	0/50	0.91	0/63
All	All	0.76	0/2922	0.73	0/3948

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	700	0	623	9	0
1	B	701	0	634	4	0
1	C	690	0	618	8	0
1	D	675	0	598	8	0
2	E	49	0	48	0	0
2	F	50	0	53	0	0
3	A	5	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	2	0	0	0	0
3	C	5	0	0	0	0
3	D	2	0	0	0	0
4	A	6	0	8	1	0
4	C	12	0	16	1	0
4	D	12	0	16	2	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
6	A	30	0	0	2	0
6	B	36	0	0	1	0
6	C	41	0	0	2	0
6	D	34	0	0	1	0
6	F	2	0	0	0	0
All	All	3059	0	2614	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1095[B]:ARG:HH21	1:A:1095[B]:ARG:CG	1.78	0.96
1:A:1095[B]:ARG:HG2	1:A:1095[B]:ARG:HH21	1.32	0.92
1:A:1095[B]:ARG:CG	1:A:1095[B]:ARG:NH2	2.50	0.65
1:A:1067:LEU:O	6:A:2101:HOH:O	2.15	0.65
1:C:1059:HIS:O	6:C:2101:HOH:O	2.15	0.63
1:A:1095[B]:ARG:HG2	1:A:1095[B]:ARG:NH2	2.09	0.62
1:A:1095[B]:ARG:HG3	1:A:1095[B]:ARG:HH21	1.66	0.59
1:D:1071:TRP:O	6:D:2101:HOH:O	2.17	0.56
1:B:1095:ARG:HG3	1:D:1095:ARG:HG2	1.88	0.56
4:A:2005:GOL:O1	6:A:2102:HOH:O	2.18	0.54
1:C:1099:LEU:HD13	1:C:1123:VAL:HG21	1.93	0.51
1:B:1095:ARG:HD3	1:D:1095:ARG:HG2	1.93	0.51
1:C:1060:CYS:HB3	4:D:1204:GOL:H12	1.95	0.47
1:A:1099:LEU:HD13	1:A:1123:VAL:HG21	1.95	0.47
1:C:1130:GLY:O	6:C:2102:HOH:O	2.21	0.46
1:A:1096:GLU:O	1:A:1097:GLU:HB2	2.17	0.44
1:C:1079:ALA:N	1:C:1080:PRO:CD	2.80	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1095:ARG:CD	1:D:1095:ARG:HG2	2.46	0.44
1:C:1106:CYS:HB3	1:C:1136:CYS:SG	2.58	0.43
1:B:1102:GLN:OE1	6:B:2101:HOH:O	2.21	0.43
1:D:1106:CYS:HB3	1:D:1136:CYS:SG	2.58	0.43
1:C:1060:CYS:CB	4:D:1204:GOL:H12	2.49	0.43
1:A:1070:GLU:HB2	1:A:1074:ASN:HA	2.01	0.42
1:D:1079:ALA:N	1:D:1080:PRO:CD	2.84	0.41
1:D:1112:ALA:HB1	1:D:1117:LEU:O	2.21	0.41
1:C:1104:ARG:NH2	4:C:2006:GOL:O3	2.49	0.41
1:D:1099:LEU:HA	1:D:1099:LEU:HD23	1.81	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	87/92 (95%)	85 (98%)	2 (2%)	0	100	100
1	B	86/92 (94%)	82 (95%)	4 (5%)	0	100	100
1	C	85/92 (92%)	81 (95%)	4 (5%)	0	100	100
1	D	84/92 (91%)	83 (99%)	1 (1%)	0	100	100
2	E	4/20 (20%)	4 (100%)	0	0	100	100
2	F	4/20 (20%)	4 (100%)	0	0	100	100
All	All	350/408 (86%)	339 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	78/81 (96%)	75 (96%)	3 (4%)	36	20
1	B	79/81 (98%)	78 (99%)	1 (1%)	71	65
1	C	78/81 (96%)	76 (97%)	2 (3%)	49	35
1	D	75/81 (93%)	73 (97%)	2 (3%)	48	34
2	E	3/11 (27%)	3 (100%)	0	100	100
2	F	4/11 (36%)	4 (100%)	0	100	100
All	All	317/346 (92%)	309 (98%)	8 (2%)	49	37

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1054	SER
1	A	1096	GLU
1	A	1099	LEU
1	B	1095	ARG
1	C	1068	ARG
1	C	1099	LEU
1	D	1095	ARG
1	D	1099	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 7 are unknown and 14 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	2005	-	5,5,5	0.22	0	5,5,5	0.29	0
4	GOL	C	2005	-	5,5,5	0.18	0	5,5,5	0.38	0
4	GOL	C	2006	-	5,5,5	0.29	0	5,5,5	0.29	0
4	GOL	D	1203	-	5,5,5	0.27	0	5,5,5	0.35	0
4	GOL	D	1204	-	5,5,5	0.10	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	2005	-	-	0/4/4/4	0/0/0/0
4	GOL	C	2005	-	-	0/4/4/4	0/0/0/0
4	GOL	C	2006	-	-	0/4/4/4	0/0/0/0
4	GOL	D	1203	-	-	0/4/4/4	0/0/0/0
4	GOL	D	1204	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2005	GOL	1	0
4	C	2006	GOL	1	0
4	D	1204	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	85/92 (92%)	0.14	4 (4%) 31 26	22, 28, 44, 53	0
1	B	85/92 (92%)	0.29	3 (3%) 44 38	22, 30, 47, 65	0
1	C	85/92 (92%)	0.17	3 (3%) 44 38	20, 26, 42, 67	0
1	D	84/92 (91%)	0.10	0 100 100	20, 27, 41, 46	0
2	E	6/20 (30%)	0.22	0 100 100	34, 37, 40, 58	0
2	F	6/20 (30%)	0.38	0 100 100	29, 37, 40, 55	0
All	All	351/408 (86%)	0.18	10 (2%) 53 48	20, 28, 46, 67	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1138	PRO	3.9
1	A	1055	VAL	2.6
1	B	1099	LEU	2.6
1	A	1075	TYR	2.2
1	A	1095[A]	ARG	2.2
1	C	1075	TYR	2.2
1	A	1096	GLU	2.1
1	B	1138	PRO	2.1
1	B	1096	GLU	2.1
1	C	1123	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	UNX	B	1204	1/1	0.67	0.30	30,30,30,30	0
4	GOL	C	2006	6/6	0.79	0.17	37,39,42,42	0
5	UNX	E	1601	1/1	0.80	0.15	30,30,30,30	0
4	GOL	D	1203	6/6	0.84	0.12	31,32,35,36	0
5	UNX	A	2006	1/1	0.86	0.33	30,30,30,30	0
5	UNX	B	1203	1/1	0.87	0.19	30,30,30,30	0
5	UNX	C	2007	1/1	0.87	0.34	30,30,30,30	0
5	UNX	A	2007	1/1	0.88	0.15	30,30,30,30	0
4	GOL	D	1204	6/6	0.89	0.14	31,38,43,46	0
5	UNX	D	1205	1/1	0.89	0.37	30,30,30,30	0
4	GOL	A	2005	6/6	0.91	0.14	31,39,46,48	0
4	GOL	C	2005	6/6	0.96	0.08	24,26,28,28	0
3	ZN	C	2003	1/1	0.99	0.08	22,22,22,22	0
3	ZN	C	2001	1/1	0.99	0.12	22,22,22,22	0
3	ZN	D	1202	1/1	0.99	0.12	22,22,22,22	0
3	ZN	C	2002	1/1	0.99	0.10	23,23,23,23	0
3	ZN	D	1201	1/1	0.99	0.07	23,23,23,23	0
3	ZN	A	2002	1/1	0.99	0.07	25,25,25,25	0
3	ZN	A	2001	1/1	0.99	0.10	21,21,21,21	0
3	ZN	C	2008	1/1	0.99	0.11	21,21,21,21	0
3	ZN	A	2004	1/1	1.00	0.10	23,23,23,23	0
3	ZN	B	1201	1/1	1.00	0.08	25,25,25,25	0
3	ZN	B	1202	1/1	1.00	0.09	24,24,24,24	0
3	ZN	C	2004	1/1	1.00	0.08	23,23,23,23	0
3	ZN	A	2003	1/1	1.00	0.10	23,23,23,23	0
3	ZN	A	2008	1/1	1.00	0.12	22,22,22,22	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.