



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 03:18 AM EDT

PDB ID : 7ML5  
Title : Structure of the Starch Branching Enzyme I (BEI) complexed with maltod-  
decaose from *Oryza sativa* L  
Authors : Nayebi Gavgani, H.; Fawaz, R.; Geiger, J.H.  
Deposited on : 2021-04-27  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	701	 85% 11% .
2	B	12	 25% 58% 17%
3	E	4	 25% 50% 25%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5904 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform 2 of 1,4-alpha-glucan-branching enzyme, chloroplastic /amyloplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	678	5528	3529	953	1010	36	25	5	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	MET	LEU	engineered mutation	UNP Q01401
A	280	MET	VAL	engineered mutation	UNP Q01401
A	443	PRO	SER	engineered mutation	UNP Q01401
A	669	ALA	THR	engineered mutation	UNP Q01401

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	12	133	72	61	4	0	0

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	E	4	Total	C	O	2	0	0
			45	24	21			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	198	Total	O	0	0
			198	198		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.67Å 80.11Å 182.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.97 – 2.35 39.97 – 2.35	Depositor EDS
% Data completeness (in resolution range)	95.3 (39.97-2.35) 95.3 (39.97-2.35)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.21 (at 2.34Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.165 , 0.232 0.164 , 0.231	Depositor DCC
$R_{free}$ test set	1457 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.9	Xtrriage
Anisotropy	0.013	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5904	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/5695	0.49	0/7716

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5528	0	5196	50	0
2	B	133	0	111	3	0
3	E	45	0	39	1	0
4	A	198	0	0	11	0
All	All	5904	0	5346	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:CYS:SG	4:A:994:HOH:O	1.96	1.20
1:A:74:PRO:HB3	3:E:1:GLC:O3	1.83	0.79
1:A:619:THR:OG1	4:A:801:HOH:O	2.01	0.77
1:A:438:GLU:OE1	1:A:441:LYS:NZ	2.20	0.74
1:A:9:ASP:OD2	4:A:802:HOH:O	2.03	0.74
1:A:366:GLU:OE1	4:A:803:HOH:O	2.09	0.68
1:A:261:HIS:ND1	4:A:808:HOH:O	2.29	0.65
1:A:595:ASN:ND2	1:A:598:ASP:OD2	2.30	0.64
1:A:619:THR:HG23	1:A:683:PRO:HA	1.81	0.62
1:A:427:PRO:O	1:A:431:ILE:HG13	2.01	0.60
1:A:391:LEU:HD23	1:A:394:ALA:HB2	1.84	0.60
1:A:193:GLU:OE2	4:A:804:HOH:O	2.17	0.58
1:A:509:MET:HG2	1:A:687[B]:CYS:SG	2.43	0.58
1:A:596:GLU:O	1:A:599:LYS:NZ	2.34	0.58
1:A:593:ASP:HB3	1:A:602:VAL:HB	1.86	0.57
1:A:409:ARG:NH1	1:A:457:THR:O	2.39	0.56
2:B:7:GLC:H62	2:B:8:GLC:H5	1.88	0.56
1:A:466:SER:N	1:A:469:GLN:OE1	2.20	0.55
1:A:42:GLU:HG3	1:A:47:GLY:HA2	1.91	0.53
1:A:109:HIS:ND1	4:A:806:HOH:O	2.25	0.52
1:A:438:GLU:OE1	1:A:438:GLU:N	2.41	0.51
1:A:535:TRP:CZ2	1:A:549:LYS:HE2	2.46	0.50
1:A:292:ASN:HD22	1:A:294:HIS:HE1	1.59	0.50
1:A:430:TRP:CE2	1:A:510:ILE:HG23	2.47	0.50
1:A:393:GLU:CD	1:A:393:GLU:H	2.15	0.49
1:A:513:ILE:HB	1:A:612:PHE:CE2	2.46	0.49
1:A:357:ASN:O	1:A:357:ASN:ND2	2.45	0.49
1:A:99:LYS:NZ	4:A:820:HOH:O	2.45	0.49
1:A:509:MET:HA	1:A:687[B]:CYS:SG	2.53	0.49
1:A:393:GLU:N	1:A:393:GLU:OE1	2.46	0.48
1:A:358:LYS:HE3	1:A:358:LYS:HB3	1.59	0.47
1:A:445:SER:HA	1:A:594:MET:SD	2.54	0.47
1:A:185:GLU:HA	1:A:220:GLN:HB3	1.97	0.46
1:A:493:LEU:HD11	2:B:11:GLC:H62	1.98	0.46
1:A:273:HIS:ND1	1:A:379:TYR:OH	2.43	0.46
1:A:406:VAL:HA	1:A:409:ARG:HG2	1.98	0.46
1:A:440:ARG:HD3	4:A:927:HOH:O	2.15	0.46
1:A:530:PHE:HB3	1:A:568:ASN:HB2	1.99	0.45
1:A:610:PHE:HB3	1:A:612:PHE:HE1	1.81	0.45
1:A:475:LYS:HE3	1:A:475:LYS:HB2	1.79	0.45
1:A:439:ASP:HB3	4:A:894:HOH:O	2.16	0.45
1:A:463:TYR:CE2	1:A:466:SER:HB3	2.52	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:451:LEU:HD13	1:A:517:LEU:HB3	1.99	0.44
1:A:484:LYS:NZ	1:A:484:LYS:HB3	2.32	0.44
1:A:632:LYS:HE3	1:A:655:ASP:CG	2.39	0.43
1:A:74:PRO:HB2	1:A:295:GLU:HB3	2.01	0.42
1:A:487:TYR:CE1	2:B:7:GLC:H3	2.54	0.42
1:A:405:PRO:HB3	1:A:455:ARG:CZ	2.51	0.41
1:A:94:LYS:NZ	4:A:835:HOH:O	2.53	0.41
1:A:48:LEU:HD21	1:A:382:LEU:HD22	2.02	0.41
1:A:187:HIS:ND1	1:A:190[A]:MET:HG2	2.36	0.41

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	679/701 (97%)	658 (97%)	19 (3%)	2 (0%)	41 47

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	406	VAL
1	A	472	VAL

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	581/607 (96%)	570 (98%)	11 (2%)	57 68

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	HIS
1	A	198	SER
1	A	233	PHE
1	A	270	ASP
1	A	389	LYS
1	A	449	GLN
1	A	460[A]	CYS
1	A	460[B]	CYS
1	A	555	SER
1	A	652	HIS
1	A	661	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	294	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	GLC	B	1	2	12,12,12	0.60	0	17,17,17	0.64	0
2	GLC	B	10	2	11,11,12	0.91	1 (9%)	15,15,17	1.00	2 (13%)
2	GLC	B	11	2	11,11,12	0.52	0	15,15,17	1.16	2 (13%)
2	GLC	B	12	2	11,11,12	0.51	0	15,15,17	1.09	1 (6%)
2	GLC	B	2	2	11,11,12	0.43	0	15,15,17	1.23	1 (6%)
2	GLC	B	3	2	11,11,12	1.44	2 (18%)	15,15,17	1.35	3 (20%)
2	GLC	B	4	2	11,11,12	0.50	0	15,15,17	0.79	0
2	GLC	B	5	2	11,11,12	0.58	0	15,15,17	0.79	1 (6%)
2	GLC	B	6	2	11,11,12	0.44	0	15,15,17	0.59	0
2	GLC	B	7	2	11,11,12	0.56	0	15,15,17	0.89	0
2	GLC	B	8	2	11,11,12	0.55	0	15,15,17	1.06	1 (6%)
2	GLC	B	9	2	11,11,12	0.54	0	15,15,17	1.21	2 (13%)
3	GLC	E	1	3	12,12,12	0.50	0	17,17,17	1.09	1 (5%)
3	GLC	E	2	3	11,11,12	0.60	0	15,15,17	1.02	1 (6%)
3	GLC	E	3	3	11,11,12	0.49	0	15,15,17	0.77	0
3	GLC	E	4	3	11,11,12	0.73	0	15,15,17	0.98	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	10	2	-	0/2/19/22	0/1/1/1
2	GLC	B	11	2	-	0/2/19/22	0/1/1/1
2	GLC	B	12	2	-	1/2/19/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	2/2/19/22	0/1/1/1
2	GLC	B	4	2	-	2/2/19/22	0/1/1/1
2	GLC	B	5	2	-	2/2/19/22	0/1/1/1
2	GLC	B	6	2	-	1/2/19/22	0/1/1/1
2	GLC	B	7	2	-	0/2/19/22	0/1/1/1
2	GLC	B	8	2	-	0/2/19/22	0/1/1/1
2	GLC	B	9	2	-	0/2/19/22	0/1/1/1
3	GLC	E	1	3	-	1/2/22/22	0/1/1/1
3	GLC	E	2	3	-	0/2/19/22	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	E	3	3	-	0/2/19/22	0/1/1/1
3	GLC	E	4	3	-	2/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	GLC	O5-C1	-4.02	1.37	1.43
2	B	10	GLC	O5-C1	2.58	1.47	1.43
2	B	3	GLC	O5-C5	-2.06	1.39	1.43

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	8	GLC	C1-C2-C3	3.01	113.37	109.67
2	B	3	GLC	O5-C5-C6	2.92	111.77	107.20
2	B	2	GLC	O5-C1-C2	-2.62	106.73	110.77
2	B	3	GLC	C1-C2-C3	2.50	112.74	109.67
2	B	3	GLC	O5-C1-C2	-2.48	106.95	110.77
2	B	12	GLC	O5-C5-C6	2.42	110.99	107.20
2	B	11	GLC	C1-C2-C3	2.39	112.60	109.67
2	B	11	GLC	C1-O5-C5	2.34	115.36	112.19
3	E	4	GLC	O5-C5-C6	2.34	110.87	107.20
2	B	9	GLC	C1-C2-C3	2.20	112.37	109.67
2	B	9	GLC	C1-O5-C5	2.17	115.13	112.19
2	B	10	GLC	C1-C2-C3	2.14	112.30	109.67
3	E	4	GLC	C1-O5-C5	-2.10	109.35	112.19
3	E	2	GLC	C1-C2-C3	2.09	112.23	109.67
3	E	1	GLC	C4-C3-C2	-2.07	107.21	110.82
2	B	5	GLC	O5-C5-C6	2.07	110.44	107.20
2	B	10	GLC	O5-C5-C4	2.05	115.82	110.83

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	5	GLC	O5-C5-C6-O6
2	B	4	GLC	C4-C5-C6-O6
3	E	4	GLC	C4-C5-C6-O6
2	B	4	GLC	O5-C5-C6-O6
2	B	5	GLC	C4-C5-C6-O6
3	E	4	GLC	O5-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	E	1	GLC	O5-C5-C6-O6
2	B	12	GLC	O5-C5-C6-O6
2	B	3	GLC	C4-C5-C6-O6
2	B	6	GLC	C4-C5-C6-O6
2	B	3	GLC	O5-C5-C6-O6


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	7	GLC	2	0
3	E	1	GLC	1	0
2	B	11	GLC	1	0
2	B	8	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	678/701 (96%)	-0.22	5 (0%) <b>87</b> <b>92</b>	25, 35, 51, 74	383 (56%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	672	ASN	3.2
1	A	165	CYS	2.9
1	A	440	ARG	2.6
1	A	9	ASP	2.5
1	A	473	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	B	12	11/12	0.81	0.36	64,69,74,80	8
3	GLC	E	1	12/12	0.82	0.23	39,48,53,58	8
2	GLC	B	1	12/12	0.87	0.24	56,62,75,77	6
2	GLC	B	2	11/12	0.88	0.22	57,61,65,69	5
2	GLC	B	5	11/12	0.90	0.20	50,56,60,61	5
2	GLC	B	3	11/12	0.91	0.19	54,60,64,65	5
2	GLC	B	11	11/12	0.91	0.20	47,50,60,66	7

*Continued on next page...*


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GLC	E	2	11/12	0.91	0.17	37,42,46,50	9
2	GLC	B	4	11/12	0.93	0.18	52,59,64,69	4
2	GLC	B	7	11/12	0.93	0.11	46,53,59,60	5
3	GLC	E	4	11/12	0.93	0.12	38,41,46,47	8
3	GLC	E	3	11/12	0.94	0.13	36,39,44,45	7
2	GLC	B	10	11/12	0.94	0.13	36,39,47,52	7
2	GLC	B	9	11/12	0.95	0.14	33,39,43,43	7
2	GLC	B	6	11/12	0.95	0.09	55,57,61,62	5
2	GLC	B	8	11/12	0.98	0.09	35,39,43,43	5

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.