



# Full wwPDB X-ray Structure Validation Report i

Jun 7, 2022 – 06:03 PM EDT

PDB ID : 7MNP  
Title : Crystal Structure of the ZnF2 of Nucleoporin NUP358/RanBP2 in complex with Ran-GDP  
Authors : Bley, C.J.; Nie, S.; Mobbs, G.W.; Petrovic, S.; Gres, A.T.; Liu, X.; Mukherjee, S.; Harvey, S.; Huber, F.M.; Lin, D.H.; Brown, B.; Tang, A.W.; Rundlet, E.J.; Correia, A.R.; Chen, S.; Regmi, S.G.; Stevens, T.A.; Jette, C.A.; Dasso, M.; Patke, A.; Palazzo, A.F.; Kossiakoff, A.A.; Hoelz, A.  
Deposited on : 2021-05-01  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.28.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.28.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 8131 atoms, of which 3937 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	201	Total	C	H	N	O	S	0	4	0
			3275	1051	1650	281	287	6			
1	C	201	Total	C	H	N	O	S	0	8	0
			3310	1061	1669	284	290	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P62826
A	35	SER	PHE	engineered mutation	UNP P62826
C	0	SER	-	expression tag	UNP P62826
C	35	SER	PHE	engineered mutation	UNP P62826

- Molecule 2 is a protein called E3 SUMO-protein ligase RanBP2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	39	Total	C	H	N	O	S	0	0	0
			598	184	297	57	56	4			
2	D	39	Total	C	H	N	O	S	0	0	0
			598	184	297	57	56	4			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	1402	GLY	-	expression tag	UNP P49792
B	1403	PRO	-	expression tag	UNP P49792
B	1404	LEU	-	expression tag	UNP P49792
B	1405	GLY	-	expression tag	UNP P49792
B	1406	SER	-	expression tag	UNP P49792
D	1402	GLY	-	expression tag	UNP P49792
D	1403	PRO	-	expression tag	UNP P49792
D	1404	LEU	-	expression tag	UNP P49792

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	1405	GLY	-	expression tag	UNP P49792
D	1406	SER	-	expression tag	UNP P49792

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			40	10	12	5	11	2		

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	C	1	Total	C	H	N	O	P	0	0
			40	10	12	5	11	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	Mg					0	0
			1	1						

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	C	1	Total	Mg					0	0
			1	1						

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	B	1	Total	Zn					0	0
			1	1						

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Zn 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	121	Total O 121 121	0	0
6	B	17	Total O 17 17	0	0
6	C	122	Total O 122 122	0	0
6	D	6	Total O 6 6	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GTP-binding nuclear protein Ran




- Molecule 1: GTP-binding nuclear protein Ran



- Molecule 2: E3 SUMO-protein ligase RanBP2



- Molecule 2: E3 SUMO-protein ligase RanBP2



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.23Å   79.84Å   108.75Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	29.02 – 2.05 29.02 – 2.05	Depositor EDS
% Data completeness (in resolution range)	98.8 (29.02-2.05) 99.1 (29.02-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.40 (at 2.06Å)	Xtriage
Refinement program	PHENIX 1.18.2	Depositor
$R$ , $R_{free}$	0.194 , 0.231 0.193 , 0.233	Depositor DCC
$R_{free}$ test set	1620 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.4	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 46.3	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8131	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.27 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.6250e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GDP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.25	0/1680	0.45	0/2276
1	C	0.25	0/1721	0.45	0/2332
2	B	0.25	0/306	0.43	0/411
2	D	0.24	0/306	0.42	0/411
All	All	0.25	0/4013	0.45	0/5430

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1625	1650	1630	3	0
1	C	1641	1669	1629	3	0
2	B	301	297	297	1	0
2	D	301	297	297	2	0
3	A	28	12	11	0	0
3	C	28	12	11	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	1	0	0	0	0
6	A	121	0	0	0	0
6	B	17	0	0	0	0
6	C	122	0	0	0	0
6	D	6	0	0	0	0
All	All	4194	3937	3875	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1415:LYS:HB3	2:B:1418:HIS:HB2	1.84	0.59
1:C:117:ILE:HB	1:C:144:LEU:HD22	1.86	0.57
1:A:117:ILE:HB	1:A:144:LEU:HD22	1.94	0.49
1:C:82:GLN:CG	2:D:1436:ILE:HD11	2.44	0.47
1:A:81:ILE:HG22	1:A:82:GLN:HG2	1.96	0.47
1:A:30:HIS:CG	1:A:50:LEU:HG	2.52	0.44
1:C:82:GLN:HG3	2:D:1436:ILE:HD11	2.01	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	203/217 (94%)	198 (98%)	5 (2%)	0	100 100
1	C	207/217 (95%)	203 (98%)	4 (2%)	0	100 100
2	B	37/42 (88%)	36 (97%)	1 (3%)	0	100 100
2	D	37/42 (88%)	36 (97%)	1 (3%)	0	100 100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	484/518 (93%)	473 (98%)	11 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	178/186 (96%)	178 (100%)	0	100 100
1	C	182/186 (98%)	182 (100%)	0	100 100
2	B	35/37 (95%)	35 (100%)	0	100 100
2	D	35/37 (95%)	35 (100%)	0	100 100
All	All	430/446 (96%)	430 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GDP	A	301	4	24,30,30	5.79	13 (54%)	31,47,47	2.39	10 (32%)
3	GDP	C	301	4	24,30,30	5.82	13 (54%)	31,47,47	2.36	9 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	301	4	-	1/12/32/32	0/3/3/3
3	GDP	C	301	4	-	1/12/32/32	0/3/3/3

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	GDP	C2'-C1'	-17.34	1.27	1.53
3	C	301	GDP	C2'-C1'	-17.26	1.27	1.53
3	C	301	GDP	C3'-C4'	-10.44	1.26	1.53
3	A	301	GDP	C3'-C4'	-10.41	1.26	1.53
3	C	301	GDP	O4'-C1'	10.25	1.55	1.41
3	A	301	GDP	O4'-C1'	9.92	1.54	1.41
3	C	301	GDP	C4-N3	7.51	1.47	1.35
3	A	301	GDP	C4-N3	7.38	1.47	1.35
3	C	301	GDP	C5-C6	7.33	1.54	1.41
3	A	301	GDP	C5-C6	7.28	1.53	1.41
3	C	301	GDP	C2-N2	6.65	1.47	1.33
3	A	301	GDP	C2-N2	6.57	1.47	1.33
3	A	301	GDP	C6-N1	6.28	1.44	1.33
3	C	301	GDP	C6-N1	6.17	1.43	1.33
3	A	301	GDP	O4'-C4'	5.18	1.56	1.45
3	C	301	GDP	O4'-C4'	5.17	1.56	1.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	GDP	C2-N1	5.08	1.44	1.35
3	C	301	GDP	C2-N1	5.03	1.44	1.35
3	C	301	GDP	C2'-C3'	5.00	1.67	1.53
3	A	301	GDP	C2'-C3'	4.89	1.66	1.53
3	C	301	GDP	C2-N3	2.51	1.46	1.34
3	A	301	GDP	O3'-C3'	2.47	1.48	1.43
3	C	301	GDP	O3'-C3'	2.45	1.48	1.43
3	A	301	GDP	C2-N3	2.40	1.45	1.34
3	C	301	GDP	C5'-C4'	2.15	1.58	1.51
3	A	301	GDP	C5'-C4'	2.13	1.58	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	301	GDP	C1'-N9-C4	8.03	140.76	126.64
3	A	301	GDP	C1'-N9-C4	7.29	139.44	126.64
3	A	301	GDP	N3-C2-N1	-6.11	119.07	127.22
3	C	301	GDP	N3-C2-N1	-5.78	119.51	127.22
3	A	301	GDP	C2-N3-C4	4.21	120.16	115.36
3	A	301	GDP	N2-C2-N1	4.18	123.75	117.25
3	C	301	GDP	C2-N3-C4	4.04	119.97	115.36
3	C	301	GDP	N2-C2-N1	3.39	122.52	117.25
3	A	301	GDP	C4-C5-C6	-2.82	118.11	120.80
3	A	301	GDP	C2-N1-C6	2.72	120.26	115.93
3	C	301	GDP	C5-C6-N1	-2.66	119.79	123.43
3	A	301	GDP	PA-O3A-PB	-2.63	123.79	132.83
3	C	301	GDP	PA-O3A-PB	-2.63	123.81	132.83
3	C	301	GDP	C2-N1-C6	2.62	120.09	115.93
3	A	301	GDP	C5-C6-N1	-2.53	119.97	123.43
3	A	301	GDP	C2'-C3'-C4'	2.13	106.77	102.64
3	C	301	GDP	C4-C5-C6	-2.10	118.79	120.80
3	A	301	GDP	C4-C5-N7	-2.09	107.22	109.40
3	C	301	GDP	C2'-C3'-C4'	2.04	106.60	102.64

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	301	GDP	C5'-O5'-PA-O1A
3	A	301	GDP	PA-O3A-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	201/217 (92%)	0.50	18 (8%) 9 10	22, 35, 69, 101	0
1	C	201/217 (92%)	0.79	32 (15%) 1 1	25, 37, 74, 114	0
2	B	39/42 (92%)	2.01	14 (35%) 0 0	30, 56, 98, 99	0
2	D	39/42 (92%)	2.36	19 (48%) 0 0	38, 70, 114, 120	0
All	All	480/518 (92%)	0.90	83 (17%) 1 1	22, 39, 88, 120	0

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	1432	VAL	7.8
2	D	1443	SER	7.1
1	C	206	THR	7.0
2	B	1412	THR	6.3
2	D	1412	THR	5.9
1	A	206	THR	5.8
2	D	1431	THR	5.5
2	B	1432	VAL	5.3
2	B	1405	GLY	5.1
2	B	1406	SER	5.1
2	B	1407	ARG	5.0
2	D	1413	PRO	4.9
2	B	1408	PHE	4.8
2	B	1431	THR	4.7
1	C	141	LYS	4.6
1	A	207	THR	4.6
2	D	1407	ARG	4.4
1	C	207	THR	4.3
2	D	1406	SER	4.3
1	C	87	ILE	4.2
2	D	1408	PHE	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	15	LEU	3.9
2	D	1405	GLY	3.8
1	C	149[A]	ILE	3.8
1	C	63	VAL	3.6
1	A	188	VAL	3.6
1	C	205	GLN	3.5
1	A	142	LYS	3.5
1	A	141	LYS	3.5
2	D	1414	LYS	3.5
2	D	1415	LYS	3.5
1	C	188	VAL	3.4
2	B	1415	LYS	3.4
2	B	1443	SER	3.4
1	C	203	VAL	3.4
2	B	1413	PRO	3.3
2	B	1414	LYS	3.2
1	C	144	LEU	3.2
1	A	15	LEU	3.1
2	D	1433	SER	3.1
1	C	13	LEU	3.0
2	B	1416	GLU	3.0
1	C	202	GLU	3.0
1	C	26	PHE	2.9
1	C	161	PHE	2.9
2	D	1418	HIS	2.9
1	C	142	LYS	2.9
1	A	118	VAL	2.8
1	A	120	CYS	2.7
1	C	120	CYS	2.7
1	C	27	VAL	2.7
1	C	86	ALA	2.7
1	C	118	VAL	2.6
1	C	160	PRO	2.6
2	D	1416	GLU	2.6
1	C	89	MET	2.6
1	C	8	GLN	2.5
2	D	1430	PRO	2.5
2	D	1411	VAL	2.5
1	C	187	VAL	2.5
2	B	1411	VAL	2.5
2	D	1419	TRP	2.4
1	A	202	GLU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	22	GLY	2.4
1	C	140	ARG	2.4
1	A	161	PHE	2.3
1	C	50	LEU	2.3
1	C	7	PRO	2.3
2	B	1433	SER	2.3
2	D	1429	GLU	2.3
2	D	1417	GLY	2.3
1	A	63	VAL	2.3
1	A	86	ALA	2.2
1	A	140	ARG	2.2
1	A	143	ASN	2.2
1	C	61	PHE	2.2
1	C	16	VAL	2.2
1	A	13	LEU	2.1
1	C	157	PHE	2.1
1	A	149[A]	ILE	2.1
1	A	203	VAL	2.1
1	C	65	ASP	2.1
1	A	14	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GDP	C	301	28/28	0.95	0.15	27,36,51,53	0
3	GDP	A	301	28/28	0.96	0.14	25,35,51,52	0
4	MG	C	302	1/1	0.97	0.19	30,30,30,30	0


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MG	A	302	1/1	0.98	0.20	29,29,29,29	0
5	ZN	D	1501	1/1	0.98	0.06	39,39,39,39	0
5	ZN	B	1501	1/1	0.99	0.09	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.