



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 02:36 am BST

PDB ID : 3MP7
Title : Lateral opening of a translocon upon entry of protein suggests the mechanism of insertion into membranes
Authors : Egea, P.F.; Stroud, R.M.; Center for Structures of Membrane Proteins (CSMP)
Deposited on : 2010-04-26
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 56%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 22%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 20%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="font-size: small; margin-top: 5px;">4% 56% 22% • 20%</p>
2	B	61	<div style="display: flex; align-items: center;"> <div style="width: 11%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 66%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="font-size: small; margin-top: 5px;">11% 66% 16% 7% 11%</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3362 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Preprotein translocase subunit secY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	387	2942	1969	468	497	8	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	469	GLY	-	EXPRESSION TAG	UNP Q8U019
A	470	THR	-	EXPRESSION TAG	UNP Q8U019
A	471	LEU	-	EXPRESSION TAG	UNP Q8U019
A	472	VAL	-	EXPRESSION TAG	UNP Q8U019
A	473	PRO	-	EXPRESSION TAG	UNP Q8U019
A	474	ARG	-	EXPRESSION TAG	UNP Q8U019
A	475	GLY	-	EXPRESSION TAG	UNP Q8U019
A	476	SER	-	EXPRESSION TAG	UNP Q8U019
A	477	HIS	-	EXPRESSION TAG	UNP Q8U019
A	478	HIS	-	EXPRESSION TAG	UNP Q8U019
A	479	HIS	-	EXPRESSION TAG	UNP Q8U019
A	480	HIS	-	EXPRESSION TAG	UNP Q8U019
A	481	HIS	-	EXPRESSION TAG	UNP Q8U019
A	482	HIS	-	EXPRESSION TAG	UNP Q8U019

- Molecule 2 is a protein called Preprotein translocase subunit secE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	54	420	278	78	63	1	0	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	71.24Å 141.57Å 235.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.43 – 2.90 49.43 – 2.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (49.43-2.90) 61.9 (49.43-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.67	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.57 (at 2.91Å)	Xtrriage
Refinement program	ELVES, BUSTER 2.8.0	Depositor
R, R_{free}	0.277 , 0.317 0.308 , 0.358	Depositor DCC
R_{free} test set	1375 reflections (8.29%)	wwPDB-VP
Wilson B-factor (Å ²)	76.3	Xtrriage
Anisotropy	0.048	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 66.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	3362	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/3011	0.79	2/4107 (0.0%)
2	B	0.52	0/428	0.83	1/574 (0.2%)
All	All	0.52	0/3439	0.80	3/4681 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	24	PRO	C-N-CA	5.81	136.23	121.70
1	A	389	PRO	N-CA-CB	5.72	110.16	103.30
1	A	395	PRO	N-CA-CB	5.36	109.73	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2942	0	2987	43	0
2	B	420	0	454	8	0
All	All	3362	0	3441	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:SER:HB3	1:A:238:ALA:HA	1.57	0.86
2:B:24:PRO:HA	2:B:25:ASN:HB2	1.62	0.79
1:A:64:VAL:HG11	1:A:432:THR:HG21	1.71	0.72
1:A:438:LEU:O	1:A:442:ILE:HG12	1.95	0.66
2:B:39:LEU:HA	2:B:42:ILE:HG22	1.79	0.63
1:A:289:ILE:HD13	1:A:418:ALA:HA	1.82	0.62
1:A:420:ILE:HG23	1:A:437:LEU:HD11	1.81	0.61
1:A:262:VAL:HB	1:A:279:ILE:HB	1.83	0.61
1:A:238:ALA:HB1	1:A:239:PRO:CD	2.31	0.60
2:B:24:PRO:HA	2:B:25:ASN:CB	2.33	0.58
1:A:221:GLN:HE21	1:A:225:LYS:HE3	1.70	0.56
1:A:431:GLY:HA3	1:A:436:ILE:HG13	1.87	0.55
1:A:153:GLN:HA	1:A:156:MET:HG2	1.88	0.54
1:A:47:PRO:HA	1:A:72:SER:HA	1.92	0.52
2:B:29:THR:HA	2:B:32:ARG:HD2	1.91	0.52
1:A:178:SER:HB2	1:A:442:ILE:HD12	1.93	0.51
2:B:35:LYS:O	2:B:39:LEU:HB2	2.12	0.50
1:A:165:ASP:HA	1:A:176:GLY:HA3	1.92	0.50
1:A:461:ALA:HA	1:A:464:LYS:HE2	1.94	0.50
1:A:332:VAL:O	1:A:351:TYR:HE1	1.95	0.49
1:A:160:VAL:HA	1:A:163:ILE:HD12	1.93	0.49
1:A:56:GLN:HB2	1:A:134:ALA:HA	1.95	0.49
1:A:456:THR:HA	1:A:459:PHE:HD2	1.78	0.48
1:A:280:ARG:HB3	1:A:283:TYR:HA	1.95	0.48
1:A:238:ALA:HB1	1:A:239:PRO:HD3	1.95	0.47
1:A:449:GLU:HA	1:A:452:ARG:HG2	1.96	0.47
1:A:246:ALA:O	1:A:250:VAL:HG23	2.15	0.47
1:A:78:ILE:HD13	1:A:177:ILE:HG23	1.96	0.46
1:A:66:LEU:HD22	1:A:79:GLY:HA2	1.97	0.46
1:A:43:LEU:HA	1:A:46:ILE:HG12	1.97	0.46
1:A:452:ARG:HA	1:A:455:ILE:HG22	1.96	0.46
2:B:22:LYS:HB3	2:B:24:PRO:HD3	1.98	0.46
1:A:190:LEU:HD21	2:B:42:ILE:HA	1.97	0.46
1:A:247:THR:HG23	1:A:436:ILE:HD11	2.00	0.44
1:A:332:VAL:O	1:A:351:TYR:CE1	2.71	0.43
1:A:468:ALA:HB3	1:A:469:GLY:HA3	2.00	0.43
1:A:315:GLY:HA2	1:A:327:GLY:HA3	2.00	0.43
1:A:280:ARG:HB2	1:A:280:ARG:HE	1.67	0.43
1:A:317:PHE:CD1	1:A:324:PRO:HB2	2.54	0.43
1:A:215:ALA:HB2	1:A:239:PRO:HB2	2.00	0.43
1:A:206:LEU:HD11	1:A:225:LYS:HD3	2.00	0.43
1:A:468:ALA:HB3	1:A:469:GLY:CA	2.49	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:THR:HB	1:A:210:PRO:HB2	2.02	0.42
1:A:181:ILE:O	1:A:185:VAL:HG23	2.19	0.42
1:A:446:PHE:O	1:A:450:ILE:HG12	2.20	0.41
1:A:191:THR:HG21	2:B:52:ARG:HD2	2.01	0.41
1:A:89:GLN:HB3	1:A:90:LEU:HD22	2.01	0.41
1:A:35:VAL:HA	1:A:38:ILE:HD12	2.03	0.41
1:A:62:LEU:HB3	1:A:66:LEU:HD12	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	373/482 (77%)	299 (80%)	53 (14%)	21 (6%)	2 5
2	B	52/61 (85%)	46 (88%)	4 (8%)	2 (4%)	3 13
All	All	425/543 (78%)	345 (81%)	57 (13%)	23 (5%)	2 6

All (23) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	GLY
1	A	232	ILE
1	A	326	SER
1	A	371	GLY
1	A	389	PRO
1	A	395	PRO
1	A	432	THR
2	B	25	ASN
1	A	134	ALA
1	A	280	ARG
1	A	309	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	26	TRP
1	A	130	ILE
1	A	210	PRO
1	A	336	ARG
1	A	358	PHE
1	A	199	ASP
1	A	281	PHE
1	A	57	ASP
1	A	211	ALA
1	A	229	TRP
1	A	377	ILE
1	A	466	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/402 (74%)	260 (87%)	38 (13%)	4	13
2	B	40/49 (82%)	34 (85%)	6 (15%)	3	9
All	All	338/451 (75%)	294 (87%)	44 (13%)	4	12

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	TRP
1	A	37	LEU
1	A	39	LEU
1	A	51	ILE
1	A	72	SER
1	A	121	MET
1	A	130	ILE
1	A	154	LEU
1	A	162	ILE
1	A	171	TRP
1	A	194	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	219	PHE
1	A	223	ILE
1	A	228	LEU
1	A	291	LEU
1	A	292	THR
1	A	295	LEU
1	A	300	GLN
1	A	306	LEU
1	A	309	LEU
1	A	317	PHE
1	A	325	ILE
1	A	347	ARG
1	A	360	LEU
1	A	368	GLU
1	A	370	THR
1	A	398	LEU
1	A	415	LEU
1	A	417	VAL
1	A	419	LEU
1	A	425	ASP
1	A	437	LEU
1	A	447	TYR
1	A	454	GLN
1	A	455	ILE
1	A	457	GLU
1	A	462	LEU
1	A	463	ARG
2	B	22	LYS
2	B	43	LEU
2	B	44	ILE
2	B	46	LEU
2	B	47	ILE
2	B	52	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	153	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/482 (80%)	0.20	20 (5%) 27 23	21, 73, 121, 144	0
2	B	54/61 (88%)	0.53	7 (12%) 3 2	49, 86, 150, 154	0
All	All	441/543 (81%)	0.24	27 (6%) 21 17	21, 74, 129, 154	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	321	THR	10.1
2	B	9	HIS	4.9
1	A	390	GLY	4.8
1	A	388	ILE	4.2
1	A	391	PHE	3.8
1	A	389	PRO	3.8
2	B	10	PHE	3.6
1	A	138	VAL	3.3
1	A	313	TRP	3.2
1	A	375	ARG	3.0
2	B	61	GLY	2.9
1	A	408	TYR	2.8
1	A	28	GLU	2.6
1	A	335	PRO	2.6
2	B	11	TRP	2.6
1	A	116	VAL	2.6
1	A	51	ILE	2.5
2	B	20	VAL	2.4
1	A	262	VAL	2.4
2	B	22	LYS	2.4
1	A	146	ILE	2.4
1	A	309	LEU	2.3
1	A	397	THR	2.3
1	A	27	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	163	ILE	2.1
2	B	12	LYS	2.0
1	A	233	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.