

Full wwPDB X-ray Structure Validation Report (i)

Oct 2, 2023 – 01:39 AM EDT

PDB ID	:	6MQK
Title	:	Crystal Structure of GTPase Domain of Human Septin 12 in complex with
		GDP
Authors	:	Castro, D.K.S.V.; Pereira, H.M.; Brandao-Neto, J.; Ulian, A.P.U.; Garratt,
		R.C.
Deposited on	:	2018-10-10
Resolution	:	2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.19 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



6MQK

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	961	Total	С	Ν	0	S	0	1	0
	A	261	2092	1322	378	377	15	0	1	0
1	В	265	Total	С	Ν	0	S	0	0	0
	ГБ	200	2133	1349	383	387	14	0		
1	С	262	Total	С	Ν	0	S	0	1	0
			2133	1346	385	388	14	0		0
1	1 D	267	Total	С	Ν	0	S	0	4	0
		207	2168	1370	395	387	16	0	4	0

• Molecule 1 is a protein called Septin-12.

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	26	MET	-	expression tag	UNP Q8IYM1
А	27	GLY	-	expression tag	UNP Q8IYM1
А	28	SER	-	expression tag	UNP Q8IYM1
A	29	SER	-	expression tag	UNP Q8IYM1
A	30	HIS	-	expression tag	UNP Q8IYM1
A	31	HIS	-	expression tag	UNP Q8IYM1
А	32	HIS	-	expression tag	UNP Q8IYM1
A	33	HIS	-	expression tag	UNP Q8IYM1
А	34	HIS	-	expression tag	UNP Q8IYM1
А	35	HIS	-	expression tag	UNP Q8IYM1
А	36	SER	-	expression tag	UNP Q8IYM1
А	37	SER	-	expression tag	UNP Q8IYM1
A	38	GLY	-	expression tag	UNP Q8IYM1
А	39	LEU	-	expression tag	UNP Q8IYM1
А	40	VAL	-	expression tag	UNP Q8IYM1
А	41	PRO	-	expression tag	UNP Q8IYM1
А	42	ARG	-	expression tag	UNP Q8IYM1
А	43	GLY	-	expression tag	UNP Q8IYM1
А	44	SER	-	expression tag	UNP Q8IYM1
А	45	HIS	-	expression tag	UNP Q8IYM1
В	26	MET	-	expression tag	UNP Q8IYM1



leference
IP Q8IYM1
IP Q8IYM1
IP Q8IYM1
P Q8IYM1
ID OSIVM1

Continued from previous page...

Chain	Residue	Modelled	lelled Actual Con		Reference
В	27	GLY	-	expression tag	UNP Q8IYM1
В	28	SER	-	expression tag	UNP Q8IYM1
В	29	SER	-	expression tag	UNP Q8IYM1
В	30	HIS	-	expression tag	UNP Q8IYM1
В	31	HIS	-	expression tag	UNP Q8IYM1
В	32	HIS	-	expression tag	UNP Q8IYM1
В	33	HIS	-	expression tag	UNP Q8IYM1
В	34	HIS	-	expression tag	UNP Q8IYM1
В	35	HIS	-	expression tag	UNP Q8IYM1
В	36	SER	-	expression tag	UNP Q8IYM1
В	37	SER	-	expression tag	UNP Q8IYM1
В	38	GLY	-	expression tag	UNP Q8IYM1
В	39	LEU	-	expression tag	UNP Q8IYM1
В	40	VAL	-	expression tag	UNP Q8IYM1
В	41	PRO	-	expression tag	UNP Q8IYM1
В	42	ARG	_	expression tag	UNP Q8IYM1
В	43	GLY	_	expression tag	UNP Q8IYM1
В	44	SER	-	expression tag	UNP Q8IYM1
В	45	HIS	-	expression tag	UNP Q8IYM1
С	26	MET	-	expression tag	UNP Q8IYM1
С	27	GLY	-	expression tag	UNP Q8IYM1
С	28	SER	-	expression tag	UNP Q8IYM1
С	29	SER	-	expression tag	UNP Q8IYM1
С	30	HIS	-	expression tag	UNP Q8IYM1
С	31	HIS	-	expression tag	UNP Q8IYM1
С	32	HIS	-	expression tag	UNP Q8IYM1
С	33	HIS	-	expression tag	UNP Q8IYM1
С	34	HIS	-	expression tag	UNP Q8IYM1
С	35	HIS	-	expression tag	UNP Q8IYM1
С	36	SER	-	expression tag	UNP Q8IYM1
С	37	SER	-	expression tag	UNP Q8IYM1
С	38	GLY	-	expression tag	UNP Q8IYM1
С	39	LEU	-	expression tag	UNP Q8IYM1
С	40	VAL	-	expression tag	UNP Q8IYM1
С	41	PRO	-	expression tag	UNP Q8IYM1
С	42	ARG	-	expression tag	UNP Q8IYM1
С	43	GLY	-	expression tag	UNP Q8IYM1
С	44	SER	-	expression tag	UNP Q8IYM1
С	45	HIS	-	expression tag	UNP Q8IYM1
D	26	MET	-	expression tag	UNP Q8IYM1
D	27	GLY	-	expression tag	UNP Q8IYM1
D	28	SER	-	expression tag	UNP Q8IYM1




6MQK

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	SER	-	expression tag	UNP Q8IYM1
D	30	HIS	-	expression tag	UNP Q8IYM1
D	31	HIS	-	expression tag	UNP Q8IYM1
D	32	HIS	-	expression tag	UNP Q8IYM1
D	33	HIS	-	expression tag	UNP Q8IYM1
D	34	HIS	-	expression tag	UNP Q8IYM1
D	35	HIS	-	expression tag	UNP Q8IYM1
D	36	SER	-	expression tag	UNP Q8IYM1
D	37	SER	-	expression tag	UNP Q8IYM1
D	38	GLY	-	expression tag	UNP Q8IYM1
D	39	LEU	-	expression tag	UNP Q8IYM1
D	40	VAL	-	expression tag	UNP Q8IYM1
D	41	PRO	-	expression tag	UNP Q8IYM1
D	42	ARG	-	expression tag	UNP Q8IYM1
D	43	GLY	-	expression tag	UNP Q8IYM1
D	44	SER	-	expression tag	UNP Q8IYM1
D	45	HIS	-	expression tag	UNP Q8IYM1

Continued from previous page...

• Molecule 2 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	٨	1	Total	С	Ν	Ο	Р	0	0
	Z A	1	28	10	5	11	2	0	0
0	р	1	Total	С	Ν	Ο	Р	0	0
	2 B	1	28	10	5	11	2	U	U



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
0	C	1	Total	С	Ν	Ο	Р	0	0
		1	28	10	5	11	2	0	0
0	Л	1	Total	С	Ν	Ο	Р	0	0
	D	DI		10	5	11	2	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	31	Total O 31 31	0	0
4	В	50	Total O 50 50	0	0
4	С	46	Total O 46 46	0	0
4	D	45	Total O 45 45	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1	Depositor	
Cell constants	46.96Å 69.25Å 88.23Å	Depositor	
a, b, c, α , β , γ	75.45° 89.46° 76.43°	Depositor	
Resolution (Å)	45.59 - 2.19	Depositor	
% Data completeness	96.7 (45.59-2.19)	Depositor	
(in resolution range)		Depositor	
R_{merge}	0.11	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$1.51 (at 2.18 \text{\AA})$	Xtriage	
Refinement program	PHENIX 1.10.1_2155	Depositor	
R, R_{free}	0.184 , 0.229	Depositor	
Wilson B-factor ($Å^2$)	42.2	Xtriage	
Anisotropy	0.330	Xtriage	
L-test for twinning ²	$ \langle L \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	8814	wwPDB-VP	
Average B, all atoms $(Å^2)$	56.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.53% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mol Type	Turne	Chain	Res	Link	Bond lengths			Bond angles		
	Ullalli	nes	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	GDP	В	401	3	24,30,30	0.95	1 (4%)	30,47,47	1.18	3 (10%)
2	GDP	D	401	3	24,30,30	0.97	1 (4%)	30,47,47	1.02	2 (6%)
2	GDP	С	401	3	24,30,30	0.91	1 (4%)	$30,\!47,\!47$	1.24	4 (13%)
2	GDP	А	401	3	24,30,30	0.95	1 (4%)	30,47,47	1.14	3 (10%)

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GDP	В	401	3	-	0/12/32/32	0/3/3/3
2	GDP	D	401	3	-	0/12/32/32	0/3/3/3
2	GDP	С	401	3	-	0/12/32/32	0/3/3/3
2	GDP	А	401	3	-	0/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
2	D	401	GDP	C6-N1	-2.55	1.34	1.37
2	В	401	GDP	O4'-C1'	2.44	1.44	1.41
2	С	401	GDP	C6-N1	-2.22	1.34	1.37
2	А	401	GDP	C6-N1	-2.19	1.34	1.37

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	401	GDP	PA-O3A-PB	-3.41	121.14	132.83
2	А	401	GDP	PA-O3A-PB	-3.00	122.53	132.83
2	В	401	GDP	C5-C6-N1	2.71	118.74	113.95
2	D	401	GDP	PA-O3A-PB	-2.61	123.86	132.83
2	С	401	GDP	C5-C6-N1	2.55	118.46	113.95
2	D	401	GDP	C8-N7-C5	2.46	107.68	102.99
2	В	401	GDP	O6-C6-C5	-2.41	119.67	124.37
2	А	401	GDP	C8-N7-C5	2.23	107.23	102.99
2	С	401	GDP	C8-N7-C5	2.20	107.17	102.99
2	В	401	GDP	PA-O3A-PB	-2.14	125.47	132.83



	Chain	1	1 0		Ζ	Observed(°)	$Ideal(^{o})$
2	С	401	GDP	O6-C6-C5	-2.12	120.23	124.37
2	А	401	GDP	O3B-PB-O3A	2.09	111.65	104.64

Continued from previous page...


There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

