



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 20, 2024 – 06:11 AM EST

PDB ID : 4MSX
Title : Crystal structure of an essential yeast splicing factor
Authors : Hadjivassiliou, H.; Guthrie, C.; Rosenberg, O.S.
Deposited on : 2013-09-18
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	502	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7108 atoms, of which 3414 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor SAD1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	395	6779	2204	3411	558	593	13	0	19	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	expression tag	UNP P43589
A	-32	GLY	-	expression tag	UNP P43589
A	-31	SER	-	expression tag	UNP P43589
A	-30	SER	-	expression tag	UNP P43589
A	-29	HIS	-	expression tag	UNP P43589
A	-28	HIS	-	expression tag	UNP P43589
A	-27	HIS	-	expression tag	UNP P43589
A	-26	HIS	-	expression tag	UNP P43589
A	-25	HIS	-	expression tag	UNP P43589
A	-24	HIS	-	expression tag	UNP P43589
A	-23	SER	-	expression tag	UNP P43589
A	-22	SER	-	expression tag	UNP P43589
A	-21	GLY	-	expression tag	UNP P43589
A	-20	LEU	-	expression tag	UNP P43589
A	-19	VAL	-	expression tag	UNP P43589
A	-18	PRO	-	expression tag	UNP P43589
A	-17	ARG	-	expression tag	UNP P43589
A	-16	GLY	-	expression tag	UNP P43589
A	-15	SER	-	expression tag	UNP P43589
A	-14	HIS	-	expression tag	UNP P43589
A	-13	MET	-	expression tag	UNP P43589
A	-12	ALA	-	expression tag	UNP P43589
A	-11	SER	-	expression tag	UNP P43589
A	-10	MET	-	expression tag	UNP P43589
A	-9	THR	-	expression tag	UNP P43589
A	-8	GLY	-	expression tag	UNP P43589
A	-7	GLY	-	expression tag	UNP P43589

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLN	-	expression tag	UNP P43589
A	-5	GLN	-	expression tag	UNP P43589
A	-4	MET	-	expression tag	UNP P43589
A	-3	GLY	-	expression tag	UNP P43589
A	-2	ARG	-	expression tag	UNP P43589
A	-1	ASP	-	expression tag	UNP P43589
A	0	PRO	-	expression tag	UNP P43589

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 7 2 3 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	321	Total O 321 321	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor SAD1



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	139.31Å 139.31Å 86.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.83 – 1.87 42.83 – 1.87	Depositor EDS
% Data completeness (in resolution range)	99.5 (42.83-1.87) 99.5 (42.83-1.87)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	10.50	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.87Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.184 , 0.220 0.177 , 0.206	Depositor DCC
R_{free} test set	1997 reflections (2.84%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtrriage
Anisotropy	0.454	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7108	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	1/3507 (0.0%)	0.78	1/4748 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	49	CYS	CB-SG	5.86	1.92	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	ARG	NE-CZ-NH2	-5.10	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3368	3411	3421	27	0
2	A	1	0	0	0	0
3	A	4	3	3	3	0
4	A	321	0	0	4	1
All	All	3694	3414	3424	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:502:ACT:H1	4:A:909:HOH:O	1.67	0.93
1:A:359[A]:ARG:NH1	1:A:438:GLU:OE1	2.10	0.85
1:A:349[B]:ASP:OD2	3:A:502:ACT:H2	1.81	0.81
1:A:188:GLU:OE1	1:A:222[A]:ARG:NH1	2.25	0.69
1:A:186[A]:GLN:NE2	4:A:817:HOH:O	2.29	0.66
1:A:386:VAL:HG21	1:A:407[B]:ILE:HD12	1.81	0.62
1:A:168[B]:ILE:HD12	1:A:239:LEU:HD23	1.87	0.57
1:A:349[B]:ASP:CG	3:A:502:ACT:H2	2.26	0.55
1:A:233:ARG:HD2	1:A:344:HIS:CG	2.43	0.53
1:A:72:GLU:O	1:A:73:LYS:HB2	2.11	0.50
1:A:168[B]:ILE:CD1	1:A:239:LEU:HD23	2.41	0.50
1:A:35:VAL:HG11	1:A:117:ILE:HD11	1.95	0.49
1:A:414:LYS:HD2	4:A:912:HOH:O	2.15	0.47
1:A:311:ILE:HD11	1:A:315:LYS:HG2	1.97	0.46
1:A:280:LYS:HE3	1:A:282:ILE:HD11	1.96	0.46
1:A:418:TRP:CD2	1:A:434:LEU:HD11	2.51	0.45
1:A:134:PRO:HD2	1:A:414:LYS:HE2	1.99	0.43
1:A:61:LEU:HD12	1:A:87:VAL:HG12	2.00	0.43
1:A:175:ARG:HD2	1:A:201:TRP:CE2	2.54	0.43
1:A:414:LYS:CD	4:A:912:HOH:O	2.67	0.42
1:A:370:LEU:C	1:A:370:LEU:HD23	2.40	0.42
1:A:370:LEU:HD21	1:A:372:ILE:HG13	2.03	0.41
1:A:218:TYR:HE1	1:A:222[A]:ARG:HD3	1.85	0.41
1:A:109:GLY:HA2	1:A:115:ASN:OD1	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:734:HOH:O	4:A:735:HOH:O[8_665]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	406/482 (84%)	397 (98%)	9 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	391/449 (87%)	390 (100%)	1 (0%)	92 92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	208	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	A	502	-	3,3,3	0.77	0	3,3,3	1.47	1 (33%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	ACT	OXT-C-CH3	2.04	123.63	115.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	ACT	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/482 (81%)	0.30	22 (5%) 24 26	23, 37, 67, 101	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	278	THR	5.7
1	A	107	HIS	5.3
1	A	448	GLU	5.1
1	A	321	THR	5.0
1	A	373	LEU	4.3
1	A	274	SER	4.2
1	A	374	HIS	4.2
1	A	55[A]	LEU	3.9
1	A	299[A]	PHE	3.6
1	A	185	ASN	3.6
1	A	388	ILE	3.6
1	A	29	TYR	3.5
1	A	109	GLY	3.5
1	A	106	LEU	3.5
1	A	350[A]	ARG	3.5
1	A	277	VAL	3.3
1	A	276	SER	2.9
1	A	322	LYS	2.9
1	A	331	VAL	2.7
1	A	266	LYS	2.5
1	A	275	GLU	2.3
1	A	349[A]	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	A	502	4/4	0.86	0.15	37,45,52,53	0
2	ZN	A	501	1/1	0.99	0.11	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.