



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 01:08 PM BST

PDB ID : 6MSY
Title : Anti-HIV-1 Fab Fab 2G12 + Man4 re-refinement
Authors : Calarese, D.A.; Stanfield, R.L.; Wilson, I.A.
Deposited on : 2018-10-18
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	213	
2	H	225	
3	A	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UNX	H	301	-	-	-	X
4	UNX	H	303	-	-	-	X
5	ACT	H	308	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3434 atoms, of which 3 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 2G12, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	210	1608	1013	268	322	5	0	0	0

- Molecule 2 is a protein called Fab 2g12, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	213	1609	1015	273	313	8	0	2	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
3	A	4	45	24	21	0	0	0

- Molecule 4 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	X		
4	H	3	3	3	0	0
4	L	1	1	1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	H	1	7	2	3	2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	72	Total	O	0	0
			72	72		
6	H	89	Total	O	0	0
			89	89		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab 2G12, light chain



- Molecule 2: Fab 2g12, heavy chain



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	144.65Å 148.32Å 54.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.39 – 2.00 29.18 – 2.00	Depositor EDS
% Data completeness (in resolution range)	93.9 (28.39-2.00) 93.9 (29.18-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.12_2829)	Depositor
R, R_{free}	0.206 , 0.242 0.217 , 0.254	Depositor DCC
R_{free} test set	1884 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	33.7	Xtrriage
Anisotropy	0.438	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.006 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3434	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, ACT, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.79	0/1644	0.83	0/2234
2	H	0.84	0/1652	0.83	0/2249
All	All	0.82	0/3296	0.83	0/4483

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1608	0	1569	21	0
2	H	1609	0	1575	30	0
3	A	45	0	38	2	0
4	H	3	0	0	0	0
4	L	1	0	0	0	0
5	H	4	3	3	2	0
6	H	89	0	0	0	0
6	L	72	0	0	2	0
All	All	3431	3	3185	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:58:ASP:OD1	5:H:308:ACT:C	2.17	0.92
2:H:12:VAL:HG11	2:H:82(C):MET:HE3	1.50	0.91
2:H:58:ASP:OD1	5:H:308:ACT:O	1.90	0.89
2:H:203:GLN:OE1	2:H:205:THR:N	2.09	0.86
1:L:183:LYS:O	1:L:187:GLU:HG2	1.81	0.80
2:H:12:VAL:HG11	2:H:82(C):MET:CE	2.15	0.77
2:H:147:TYR:CE2	2:H:152:VAL:HG23	2.27	0.69
1:L:169:LYS:HE2	1:L:169:LYS:HA	1.80	0.64
1:L:125:LEU:HD11	1:L:186:TYR:CD2	2.32	0.64
2:H:205:THR:HB	2:H:222:LYS:CE	2.29	0.62
2:H:100(B)[B]:ASP:OD1	3:A:1:MAN:H2	2.00	0.62
1:L:203:SER:HB2	1:L:204:PRO:HD2	1.83	0.60
2:H:184:LEU:HD12	2:H:184:LEU:N	2.17	0.59
1:L:166:GLN:HG3	1:L:173:TYR:CZ	2.39	0.58
2:H:205:THR:HB	2:H:222:LYS:HE3	1.85	0.58
1:L:147:GLN:HB3	1:L:154:LEU:CD1	2.35	0.57
1:L:125:LEU:HD11	1:L:186:TYR:HD2	1.70	0.56
2:H:1:GLU:O	2:H:26:ASN:ND2	2.27	0.56
2:H:182:SER:O	2:H:184:LEU:HD12	2.07	0.55
1:L:189:HIS:CE1	6:L:406:HOH:O	2.63	0.51
1:L:151:ASP:OD1	1:L:191:VAL:HG12	2.12	0.50
2:H:182:SER:OG	2:H:184:LEU:HD13	2.11	0.49
2:H:205:THR:HB	2:H:222:LYS:HE2	1.94	0.49
2:H:6:GLU:OE1	2:H:104:GLY:HA3	2.14	0.47
2:H:12:VAL:HG21	2:H:82(C):MET:HE1	1.97	0.47
2:H:136:GLY:H	2:H:195:SER:HB3	1.80	0.46
1:L:103:ARG:HD2	6:L:401:HOH:O	2.15	0.46
2:H:18:LEU:HB2	2:H:82(C):MET:HE1	1.98	0.46
1:L:33:LEU:HG	1:L:34:ALA:N	2.31	0.46
1:L:151:ASP:HA	1:L:191:VAL:HG13	1.98	0.45
2:H:152:VAL:HG12	2:H:153:THR:N	2.32	0.45
1:L:4:MET:HE3	1:L:23:CYS:SG	2.57	0.45
1:L:120:PRO:HG3	1:L:130:ALA:HB1	1.98	0.45
2:H:226:GLU:HB2	2:H:227:PRO:HD2	1.99	0.44
1:L:78:LEU:HD22	1:L:82:ASP:HB2	2.01	0.43
2:H:147:TYR:CD2	2:H:152:VAL:HG23	2.53	0.43
1:L:175:LEU:C	1:L:175:LEU:HD23	2.39	0.43
2:H:19:ILE:HD13	2:H:81:GLN:HG2	2.00	0.43
2:H:14:ALA:O	2:H:115:SER:HB3	2.19	0.42
2:H:121:VAL:O	2:H:221:LYS:HE3	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:100(B)[B]:ASP:CG	3:A:1:MAN:H2	2.40	0.42
2:H:35:ASN:OD1	2:H:50:SER:HB2	2.18	0.42
1:L:4:MET:CE	1:L:23:CYS:SG	3.07	0.42
2:H:152:VAL:HG11	2:H:210:VAL:HG13	2.01	0.41
1:L:75:ILE:HG21	1:L:78:LEU:HD23	2.02	0.41
2:H:12:VAL:HG21	2:H:82(C):MET:CE	2.51	0.41
1:L:142:ARG:HB2	1:L:173:TYR:CE2	2.55	0.41
2:H:152:VAL:CG1	2:H:210:VAL:HG13	2.51	0.41
1:L:163:VAL:HG22	1:L:175:LEU:HD12	2.03	0.41
2:H:100(D):ASP:HB3	2:H:100(E):PRO:HD2	2.03	0.41
1:L:81:ASP:N	1:L:81:ASP:OD1	2.43	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	208/213 (98%)	198 (95%)	10 (5%)	0	100	100
2	H	209/225 (93%)	202 (97%)	7 (3%)	0	100	100
All	All	417/438 (95%)	400 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	181/184 (98%)	178 (98%)	3 (2%)	60	65
2	H	181/190 (95%)	175 (97%)	6 (3%)	38	37
All	All	362/374 (97%)	353 (98%)	9 (2%)	52	49

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	55	LYS
1	L	81	ASP
1	L	169	LYS
2	H	2	VAL
2	H	5	VAL
2	H	19	ILE
2	H	100(B)[A]	ASP
2	H	100(B)[B]	ASP
2	H	203	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	A	1	3	12,12,12	2.54	5 (41%)	17,17,17	2.59	7 (41%)
3	MAN	A	2	3	11,11,12	1.76	5 (45%)	15,15,17	1.32	2 (13%)
3	MAN	A	3	3	11,11,12	1.78	6 (54%)	15,15,17	1.39	2 (13%)
3	MAN	A	4	3	11,11,12	2.35	4 (36%)	15,15,17	1.07	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	A	1	3	-	0/2/22/22	0/1/1/1
3	MAN	A	2	3	-	0/2/19/22	0/1/1/1
3	MAN	A	3	3	-	0/2/19/22	0/1/1/1
3	MAN	A	4	3	-	0/2/19/22	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	4	MAN	O4-C4	4.80	1.54	1.43
3	A	1	MAN	O1-C1	4.70	1.54	1.39
3	A	1	MAN	C1-C2	4.49	1.63	1.52
3	A	4	MAN	C2-C3	4.12	1.58	1.52
3	A	1	MAN	O5-C5	-3.82	1.35	1.44
3	A	1	MAN	O3-C3	3.01	1.50	1.43
3	A	3	MAN	C6-C5	2.94	1.61	1.51
3	A	2	MAN	C4-C5	2.81	1.59	1.53
3	A	2	MAN	C4-C3	2.73	1.59	1.52
3	A	4	MAN	O2-C2	-2.50	1.38	1.43
3	A	2	MAN	C2-C3	2.47	1.56	1.52
3	A	3	MAN	C1-C2	2.38	1.57	1.52
3	A	2	MAN	C6-C5	2.29	1.59	1.51
3	A	4	MAN	O5-C5	2.28	1.48	1.43
3	A	1	MAN	O2-C2	-2.18	1.37	1.43
3	A	3	MAN	O2-C2	2.14	1.47	1.43
3	A	3	MAN	O5-C1	-2.05	1.40	1.43
3	A	3	MAN	C4-C3	2.05	1.57	1.52
3	A	2	MAN	O5-C5	2.03	1.47	1.43
3	A	3	MAN	C2-C3	-2.02	1.49	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1	MAN	C6-C5-C4	-5.35	100.48	113.00
3	A	1	MAN	O3-C3-C2	5.17	122.29	110.35
3	A	1	MAN	C4-C3-C2	3.79	117.43	110.82
3	A	1	MAN	C1-O5-C5	3.38	120.05	113.66
3	A	3	MAN	C1-O5-C5	3.24	116.58	112.19
3	A	2	MAN	O2-C2-C3	-3.00	104.12	110.14
3	A	1	MAN	C3-C4-C5	2.62	114.90	110.24
3	A	2	MAN	O5-C5-C6	2.58	111.25	107.20
3	A	1	MAN	O4-C4-C3	-2.56	104.42	110.35
3	A	1	MAN	O3-C3-C4	-2.35	104.92	110.35
3	A	4	MAN	O2-C2-C3	-2.09	105.95	110.14
3	A	3	MAN	O3-C3-C2	2.06	113.94	109.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	MAN	2	0

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are unknown - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	H	308	-	1,3,3	10.49	1 (100%)	0,3,3	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	308	ACT	CH3-C	10.49	1.62	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	308	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	L	210/213 (98%)	0.33	8 (3%) 40 39	25, 39, 66, 77	0
2	H	213/225 (94%)	0.73	21 (9%) 7 6	24, 40, 73, 85	0
All	All	423/438 (96%)	0.53	29 (6%) 16 16	24, 40, 71, 85	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	135	GLY	4.4
2	H	165	ALA	4.3
2	H	213	LYS	4.1
2	H	116	THR	4.0
1	L	186	TYR	3.7
2	H	214	PRO	3.7
2	H	37	VAL	3.4
2	H	216	ASN	3.3
2	H	92[A]	CYS	3.1
2	H	114	ALA	3.0
2	H	79	TYR	2.9
2	H	167	THR	2.7
1	L	191	VAL	2.7
2	H	142	CYS	2.7
2	H	122	PHE	2.7
2	H	209	ASN	2.7
2	H	1	GLU	2.6
2	H	36	TRP	2.4
1	L	202	SER	2.4
1	L	203	SER	2.2
2	H	143	LEU	2.2
1	L	47	LEU	2.2
1	L	190	LYS	2.2
2	H	115	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	212	HIS	2.1
1	L	46	LEU	2.1
1	L	154	LEU	2.1
2	H	205	THR	2.1
2	H	227	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	A	1	12/12	0.90	0.12	34,43,58,60	0
3	MAN	A	2	11/12	0.94	0.09	31,34,39,52	0
3	MAN	A	3	11/12	0.94	0.11	26,30,34,36	0
3	MAN	A	4	11/12	0.95	0.12	24,27,30,30	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	UNX	H	301	1/1	0.73	0.60	31,31,31,31	0
4	UNX	H	303	1/1	0.77	0.73	63,63,63,63	0
4	UNX	L	301	1/1	0.83	0.63	35,35,35,35	0
4	UNX	H	302	1/1	0.83	0.63	56,56,56,56	0
5	ACT	H	308	4/4	0.96	0.06	32,40,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.