



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2023 – 06:34 AM EDT

PDB ID : 7MWX
Title : Structure of the core ectodomain of the hepatitis C virus envelope glycoprotein 2 with tamarin CD81
Authors : Kumar, A.; Hossain, R.A.; Yost, S.A.; Bu, W.; Wang, Y.; Dearborn, A.D.; Grakoui, A.; Cohen, J.I.; Marcotrigiano, J.
Deposited on : 2021-05-17
Resolution : 3.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1089 (3.36-3.28)
Clashscore	141614	1137 (3.36-3.28)
Ramachandran outliers	138981	1115 (3.36-3.28)
Sidechain outliers	138945	1114 (3.36-3.28)
RSRZ outliers	127900	1059 (3.36-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	259	
1	G	259	
2	A	218	
2	E	218	
3	B	218	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	218	 5% 88% 11%
4	D	93	 4% 59% 26% 13%
4	H	93	 18% 67% 23% 10%
5	I	3	 67% 33%
6	J	2	 100%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Core protein precursor eE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	172	Total	C	N	O	S	0	0	0
			1289	828	214	233	14			
1	G	180	Total	C	N	O	S	0	0	0
			1328	845	222	247	14			

- Molecule 2 is a protein called 2A12 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	218	Total	C	N	O	S	0	0	0
			1607	1009	264	329	5			
2	E	218	Total	C	N	O	S	0	0	0
			1628	1020	266	337	5			

- Molecule 3 is a protein called 2A12 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	B	218	Total	C	N	O	S	0	0	0
			1659	1030	277	344	8			
3	F	218	Total	C	N	O	S	0	0	0
			1676	1040	282	346	8			


- Molecule 4 is a protein called CD81 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	81	Total	C	N	O	S	0	0	0
			555	338	94	118	5			
4	H	84	Total	C	N	O	S	0	0	0
			608	374	102	127	5			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	7	THR	-	cloning artifact	UNP Q9N0J9
D	8	GLY	-	cloning artifact	UNP Q9N0J9
D	9	SER	-	cloning artifact	UNP Q9N0J9
H	7	THR	-	cloning artifact	UNP Q9N0J9
H	8	GLY	-	cloning artifact	UNP Q9N0J9
H	9	SER	-	cloning artifact	UNP Q9N0J9

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	I	3	39	22	2	15	0	0	0

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	J	2	28	16	2	10	0	0	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	C	1	Total 14	8	1	5	0	0
7	C	1	Total 14	8	1	5	0	0
7	G	1	Total 14	8	1	5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Core protein precursor eE2




- Molecule 1: Core protein precursor eE2





- Molecule 2: 2A12 Fab Heavy Chain









• Molecule 2: 2A12 Fab Heavy Chain



• Molecule 3: 2A12 Fab Light Chain



• Molecule 3: 2A12 Fab Light Chain




• Molecule 4: CD81 protein



• Molecule 4: CD81 protein





- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 67% 33%



- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.95Å 127.77Å 212.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.09 – 3.32 54.74 – 3.32	Depositor EDS
% Data completeness (in resolution range)	99.4 (53.09-3.32) 99.5 (54.74-3.32)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 3.33Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.239 , 0.283 0.242 , 0.285	Depositor DCC
R_{free} test set	1593 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	124.4	Xtrriage
Anisotropy	0.499	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 103.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	10459	wwPDB-VP
Average B, all atoms (Å ²)	168.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.26	0/1335	0.54	0/1837
1	G	0.27	0/1372	0.56	0/1889
2	A	0.25	0/1647	0.46	0/2263
2	E	0.25	0/1668	0.50	1/2288 (0.0%)
3	B	0.24	0/1696	0.42	0/2310
3	F	0.25	0/1713	0.42	0/2329
4	D	0.26	0/559	0.47	0/764
4	H	0.25	0/614	0.45	0/834
All	All	0.25	0/10604	0.48	1/14514 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	100	ASP	CB-CG-OD2	5.19	122.97	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	558	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1289	0	1167	42	0
1	G	1328	0	1189	51	0
2	A	1607	0	1532	12	0
2	E	1628	0	1564	31	0
3	B	1659	0	1543	12	0
3	F	1676	0	1578	15	0
4	D	555	0	495	21	0
4	H	608	0	561	18	0
5	I	39	0	34	2	0
6	J	28	0	25	0	0
7	C	28	0	26	0	0
7	G	14	0	13	0	0
All	All	10459	0	9727	189	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 189 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:29:ILE:HD13	2:E:53:PRO:CB	1.93	0.98
1:C:421:HIS:HB3	4:D:80:SER:H	1.31	0.95
1:G:501:ALA:H	1:G:538:VAL:HG22	1.34	0.92
2:A:91:THR:HG22	2:A:116:VAL:H	1.38	0.89
1:G:538:VAL:HG12	1:G:539:PHE:H	1.37	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	166/259 (64%)	142 (86%)	20 (12%)	4 (2%)	6	29
1	G	174/259 (67%)	146 (84%)	24 (14%)	4 (2%)	6	30
2	A	216/218 (99%)	207 (96%)	9 (4%)	0	100	100
2	E	216/218 (99%)	201 (93%)	13 (6%)	2 (1%)	17	49
3	B	216/218 (99%)	207 (96%)	9 (4%)	0	100	100
3	F	216/218 (99%)	207 (96%)	9 (4%)	0	100	100
4	D	79/93 (85%)	69 (87%)	10 (13%)	0	100	100
4	H	82/93 (88%)	75 (92%)	7 (8%)	0	100	100
All	All	1365/1576 (87%)	1254 (92%)	101 (7%)	10 (1%)	22	55

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	512	THR
1	C	526	ALA
1	G	512	THR
1	G	570	PRO
2	E	138	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	137/227 (60%)	135 (98%)	2 (2%)	65	81
1	G	140/227 (62%)	137 (98%)	3 (2%)	53	76
2	A	180/187 (96%)	177 (98%)	3 (2%)	60	79
2	E	186/187 (100%)	181 (97%)	5 (3%)	44	72
3	B	187/195 (96%)	185 (99%)	2 (1%)	73	85
3	F	191/195 (98%)	190 (100%)	1 (0%)	88	93

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	D	58/82 (71%)	55 (95%)	3 (5%)	23	55
4	H	66/82 (80%)	65 (98%)	1 (2%)	65	81
All	All	1145/1382 (83%)	1125 (98%)	20 (2%)	60	79

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	96	CYS
2	E	201	ASN
4	H	66	MET
3	F	217	ARG
2	A	106	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	I	1	5,1	14,14,15	0.51	0	17,19,21	1.30	2 (11%)
5	NAG	I	2	5	14,14,15	0.30	0	17,19,21	0.43	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BMA	I	3	5	11,11,12	0.62	0	15,15,17	0.77	0
6	NAG	J	1	6,1	14,14,15	0.25	0	17,19,21	0.43	0
6	NAG	J	2	6	14,14,15	0.26	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	I	1	5,1	-	5/6/23/26	0/1/1/1
5	NAG	I	2	5	-	4/6/23/26	0/1/1/1
5	BMA	I	3	5	-	0/2/19/22	0/1/1/1
6	NAG	J	1	6,1	-	2/6/23/26	0/1/1/1
6	NAG	J	2	6	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	1	NAG	C2-N2-C7	4.33	129.07	122.90
5	I	1	NAG	C1-C2-N2	2.03	113.96	110.49

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	I	2	NAG	O5-C5-C6-O6
5	I	1	NAG	C8-C7-N2-C2
5	I	1	NAG	O7-C7-N2-C2
5	I	2	NAG	C8-C7-N2-C2
5	I	2	NAG	O7-C7-N2-C2


There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	I	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	NAG	C	701	1	14,14,15	0.40	0	17,19,21	0.52	0
7	NAG	G	701	1	14,14,15	0.24	0	17,19,21	0.43	0
7	NAG	C	702	1	14,14,15	0.27	0	17,19,21	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	C	701	1	-	3/6/23/26	0/1/1/1
7	NAG	G	701	1	-	2/6/23/26	0/1/1/1
7	NAG	C	702	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	C	702	NAG	O5-C5-C6-O6
7	C	702	NAG	C4-C5-C6-O6
7	G	701	NAG	C4-C5-C6-O6
7	C	701	NAG	C4-C5-C6-O6
7	C	701	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	C	172/259 (66%)	1.33	52 (30%) 0 0	121, 209, 273, 347	0
1	G	180/259 (69%)	1.01	46 (25%) 0 0	119, 214, 290, 351	0
2	A	218/218 (100%)	0.44	15 (6%) 16 17	90, 121, 164, 222	0
2	E	218/218 (100%)	0.31	13 (5%) 21 22	80, 136, 205, 257	0
3	B	218/218 (100%)	0.36	16 (7%) 15 15	94, 139, 204, 301	0
3	F	218/218 (100%)	0.24	11 (5%) 28 28	108, 143, 202, 226	0
4	D	81/93 (87%)	-0.01	4 (4%) 29 29	163, 228, 294, 355	0
4	H	84/93 (90%)	0.76	17 (20%) 1 0	169, 219, 288, 328	0
All	All	1389/1576 (88%)	0.55	174 (12%) 3 3	80, 155, 264, 355	0

The worst 5 of 174 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	509	TYR	10.5
1	C	613	ILE	9.1
1	C	511	PHE	7.6
1	C	518	VAL	7.1
1	G	506	GLY	6.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	J	2	14/15	0.72	0.24	272,316,333,333	0
5	BMA	I	3	11/12	0.76	0.18	207,250,262,265	0
5	NAG	I	2	14/15	0.80	0.24	232,276,302,307	0
6	NAG	J	1	14/15	0.89	0.20	246,278,321,325	0
5	NAG	I	1	14/15	0.90	0.25	190,227,241,265	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	NAG	C	702	14/15	0.51	0.29	156,207,230,240	0
7	NAG	C	701	14/15	0.59	0.24	218,249,259,260	0
7	NAG	G	701	14/15	0.67	0.20	150,226,238,249	0

6.5 Other polymers [i](#)

There are no such residues in this entry.