



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2022 – 05:59 PM EDT

PDB ID : 4MYV  
Title : Free HSV-2 gD structure  
Authors : Lu, G.; Zhang, N.; Qi, J.; Li, Y.; Chen, Z.; Zheng, C.; Yan, J.; Gao, G.F.  
Deposited on : 2013-09-28  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	
1	B	294	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3240 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	176	1406	904	240	253	9	0	1	0
1	B	176	1399	899	238	253	9	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	ALA	-	expression tag	UNP P03172
A	-1	ASP	-	expression tag	UNP P03172
A	0	LEU	-	expression tag	UNP P03172
A	27	GLN	ARG	SEE REMARK 999	UNP P03172
A	286	HIS	-	expression tag	UNP P03172
A	287	HIS	-	expression tag	UNP P03172
A	288	HIS	-	expression tag	UNP P03172
A	289	HIS	-	expression tag	UNP P03172
A	290	HIS	-	expression tag	UNP P03172
A	291	HIS	-	expression tag	UNP P03172
B	-2	ALA	-	expression tag	UNP P03172
B	-1	ASP	-	expression tag	UNP P03172
B	0	LEU	-	expression tag	UNP P03172
B	27	GLN	ARG	SEE REMARK 999	UNP P03172
B	286	HIS	-	expression tag	UNP P03172
B	287	HIS	-	expression tag	UNP P03172
B	288	HIS	-	expression tag	UNP P03172
B	289	HIS	-	expression tag	UNP P03172
B	290	HIS	-	expression tag	UNP P03172
B	291	HIS	-	expression tag	UNP P03172

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
2	A	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	202	202	202	0	0
3	B	205	205	205	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.56Å 55.35Å 65.33Å 90.00° 96.35° 90.00°	Depositor
Resolution (Å)	36.27 – 1.80 36.27 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.8 (36.27-1.80) 96.8 (36.27-1.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.01 (at 1.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.189 , 0.230 0.189 , 0.230	Depositor DCC
$R_{free}$ test set	2038 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.7	Xtrriage
Anisotropy	0.429	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 47.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.024 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3240	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1450	0.63	1/1981 (0.1%)
1	B	0.39	0/1439	0.69	4/1966 (0.2%)
All	All	0.39	0/2889	0.66	5/3947 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	ALA	N-CA-C	-7.78	89.98	111.00
1	B	87	GLU	N-CA-C	6.04	127.32	111.00
1	B	86	ASP	N-CA-C	5.71	126.42	111.00
1	B	88	ALA	N-CA-CB	5.69	118.06	110.10
1	A	222	ARG	NE-CZ-NH1	5.29	122.95	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1406	0	1379	27	0
1	B	1399	0	1370	20	0
2	A	14	0	13	2	0
2	B	14	0	13	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	202	0	0	6	0
3	B	205	0	0	3	0
All	All	3240	0	2775	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:SER:O	3:A:638:HOH:O	1.62	1.17
1:B:101:ARG:NH1	1:B:159:GLU:O	2.09	0.85
1:A:64:ARG:HD3	1:A:67:ARG:HG2	1.62	0.81
1:A:94:ASN:HD21	2:A:401:NAG:C1	1.94	0.80
1:A:119:PRO:HG2	1:A:122:LYS:HG2	1.66	0.76
1:A:69:VAL:HG23	1:A:156:PRO:HG2	1.74	0.69
1:A:62:LEU:O	1:A:64:ARG:NE	2.23	0.68
1:B:94:ASN:HD21	2:B:401:NAG:C1	2.11	0.63
1:A:126:VAL:N	3:A:638:HOH:O	2.22	0.61
1:B:94:ASN:HD21	2:B:401:NAG:C2	2.15	0.59
1:A:64:ARG:HD3	1:A:67:ARG:CG	2.32	0.58
1:B:62:LEU:HD11	1:B:160:THR:HG22	1.86	0.57
1:B:117:GLU:HG3	1:B:129:ILE:HD12	1.86	0.57
1:B:91:HIS:O	3:B:583:HOH:O	2.17	0.57
1:A:151:PHE:HE2	1:A:153:MET:HG3	1.72	0.55
1:A:184:ARG:CZ	1:A:184:ARG:HB3	2.38	0.54
1:A:63:GLU:HG3	1:A:64:ARG:H	1.73	0.54
1:A:123:SER:OG	1:A:124:LEU:N	2.42	0.53
1:A:196:ARG:HG2	3:A:654:HOH:O	2.10	0.52
1:A:113:MET:HG2	1:A:133:PRO:HG3	1.91	0.52
1:B:76:GLU:OE1	3:B:666:HOH:O	2.19	0.52
1:A:91:HIS:O	3:A:608:HOH:O	2.19	0.51
1:B:84:ALA:HB1	1:B:88:ALA:HB3	1.92	0.50
1:A:154[B]:HIS:NE2	3:A:657:HOH:O	2.27	0.49
1:A:222:ARG:HD2	3:A:568:HOH:O	2.12	0.49
1:A:183:HIS:O	1:A:184:ARG:HB2	2.14	0.48
1:B:158:PHE:HB3	1:B:197:ILE:HD12	1.96	0.47
1:B:101:ARG:NH1	1:B:164:TYR:OH	2.47	0.47
1:A:94:ASN:ND2	2:A:401:NAG:C1	2.73	0.47
1:B:62:LEU:HD21	1:B:69:VAL:HG21	1.98	0.45
1:B:195:LEU:HD22	1:B:195:LEU:HA	1.83	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:ARG:HH21	1:B:121:ASN:HB2	1.83	0.44
1:A:79:GLN:OE1	1:A:82:ARG:NH1	2.37	0.43
1:A:51:PRO:HG3	1:A:55:ILE:HG13	2.00	0.43
1:A:62:LEU:HD13	1:A:183:HIS:ND1	2.33	0.42
1:A:64:ARG:HB2	1:A:65:ALA:H	1.61	0.42
1:A:88:ALA:HB1	1:A:91:HIS:HB2	2.02	0.42
1:A:206:LYS:HE2	1:A:206:LYS:HB3	1.80	0.42
1:B:85:SER:O	1:B:88:ALA:N	2.53	0.41
1:B:130:ARG:NH2	3:B:545:HOH:O	2.51	0.41
1:B:88:ALA:O	1:B:91:HIS:HB2	2.20	0.41
1:B:94:ASN:HD21	2:B:401:NAG:H2	1.86	0.41
1:A:87:GLU:HA	1:A:88:ALA:HA	1.55	0.41
1:B:151:PHE:CE2	1:B:153:MET:HB2	2.56	0.41
1:B:153:MET:CE	1:B:158:PHE:HE2	2.34	0.41
1:A:82:ARG:HA	1:A:83:GLY:HA3	1.92	0.41
1:B:81:VAL:HG13	1:B:170:ILE:HD13	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/294 (59%)	160 (92%)	12 (7%)	1 (1%)	25	12
1	B	172/294 (58%)	164 (95%)	7 (4%)	1 (1%)	25	12
All	All	345/588 (59%)	324 (94%)	19 (6%)	2 (1%)	25	12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	65	ALA
1	A	86	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/255 (60%)	146 (95%)	7 (5%)	27	13
1	B	152/255 (60%)	145 (95%)	7 (5%)	27	13
All	All	305/510 (60%)	291 (95%)	14 (5%)	27	13

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	VAL
1	A	64	ARG
1	A	90	LYS
1	A	123	SER
1	A	184	ARG
1	A	195	LEU
1	A	196	ARG
1	B	64	ARG
1	B	66	CYS
1	B	90	LYS
1	B	105	ASN
1	B	117	GLU
1	B	124	LEU
1	B	195	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	209	GLN
1	B	94	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	401	-	14,14,15	0.79	1 (7%)	17,19,21	0.61	0
2	NAG	A	401	-	14,14,15	0.79	1 (7%)	17,19,21	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	401	-	-	0/6/23/26	0/1/1/1
2	NAG	A	401	-	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAG	O5-C1	-2.83	1.39	1.43
2	A	401	NAG	O5-C1	-2.69	1.39	1.43

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	NAG	3	0
2	A	401	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	176/294 (59%)	0.67	29 (16%) <b>1</b> <b>1</b>	9, 20, 62, 86	0
1	B	176/294 (59%)	0.71	28 (15%) <b>1</b> <b>1</b>	9, 21, 61, 72	0
All	All	352/588 (59%)	0.69	57 (16%) <b>1</b> <b>1</b>	9, 20, 62, 86	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	66	CYS	20.6
1	B	195	LEU	15.5
1	A	66	CYS	12.3
1	B	65	ALA	9.3
1	A	157	ALA	9.0
1	A	195	LEU	8.9
1	A	155	ALA	8.0
1	A	158	PHE	7.3
1	A	84	ALA	7.2
1	B	87	GLU	6.5
1	B	85	SER	6.3
1	A	64	ARG	6.2
1	B	88	ALA	6.1
1	A	65	ALA	5.8
1	A	87	GLU	5.7
1	A	88	ALA	5.5
1	B	83	GLY	5.4
1	A	196	ARG	5.3
1	B	196	ARG	5.2
1	A	63	GLU	4.8
1	A	86	ASP	4.7
1	B	89	ARG	4.6
1	A	159	GLU	4.5
1	A	83	GLY	4.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	38	TYR	4.5
1	A	89	ARG	4.4
1	B	90	LYS	4.3
1	B	37	VAL	4.2
1	A	156	PRO	4.2
1	B	84	ALA	3.9
1	A	184	ARG	3.9
1	A	85	SER	3.8
1	A	67	ARG	3.8
1	B	86	ASP	3.7
1	B	82	ARG	3.7
1	B	64	ARG	3.5
1	B	91	HIS	3.5
1	B	63	GLU	3.4
1	B	159	GLU	3.1
1	A	68	SER	3.1
1	B	155	ALA	3.0
1	A	121	ASN	2.9
1	A	183	HIS	2.9
1	B	67	ARG	2.8
1	A	154[A]	HIS	2.7
1	A	82	ARG	2.6
1	B	157	ALA	2.5
1	A	62	LEU	2.5
1	B	62	LEU	2.5
1	B	158	PHE	2.4
1	A	122	LYS	2.2
1	A	123	SER	2.2
1	B	156	PRO	2.2
1	A	91	HIS	2.1
1	B	39	HIS	2.1
1	B	68	SER	2.1
1	B	154	HIS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	401	14/15	0.77	0.21	31,45,49,52	0
2	NAG	A	401	14/15	0.78	0.17	28,36,38,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.