

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 29, 2024 – 01:48 AM EDT

PDB ID	:	3MZ1
Title	:	The crystal structure of a possible TRANSCRIPTION REGULATOR PRO-
		TEIN from Sinorhizobium meliloti 1021
Authors	:	Tan, K.; Xu, X.; Cui, H.; Chin, S.; Savchenko, A.; Edwards, A.; Joachimiak,
		A.; Midwest Center for Structural Genomics (MCSG)
Deposited on	:	2010-05-11
Resolution	:	1.88  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	164625	1090 (1.88-1.88)
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)
RSRZ outliers	164620	1090 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain								
1	А	300	<u>6%</u> 53%	15% •	31%						
1	В	300	5%	11% •	29%						
1	С	300	5%	13% •	32%						
1	D	300	4%	15%	29%						



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7104 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Atoms						AltConf	Trace
1	Λ	208	Total	С	Ν	0	S	Se	0	2	0
1	Л	208	1685	1074	303	298	3	7	0	5	0
1	В	212	Total	С	Ν	0	S	Se	0	3	0
1	D		1704	1088	302	304	3	7	0	5	
1	С	204	Total	С	Ν	0	S	Se	0	9	0
		204	1629	1042	288	288	4	7	0	Δ	0
1	1 D	919	Total	С	Ν	0	S	Se	0	5	0
	212	1722	1099	308	305	3	7		5	0	

• Molecule 1 is a protein called Putative transcriptional regulator.

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-1	GLN	-	expression tag	UNP Q92SG7
А	0	GLY	-	expression tag	UNP Q92SG7
А	215	MSE	VAL	cloning artifact	UNP Q92SG7
А	297	GLY	-	expression tag	UNP Q92SG7
А	298	SER	-	expression tag	UNP Q92SG7
В	-1	GLN	-	expression tag	UNP Q92SG7
В	0	GLY	-	expression tag	UNP Q92SG7
В	215	MSE	VAL	cloning artifact	UNP Q92SG7
В	297	GLY	-	expression tag	UNP Q92SG7
В	298	SER	-	expression tag	UNP Q92SG7
С	-1	GLN	-	expression tag	UNP Q92SG7
С	0	GLY	-	expression tag	UNP Q92SG7
С	215	MSE	VAL	cloning artifact	UNP Q92SG7
С	297	GLY	-	expression tag	UNP Q92SG7
С	298	SER	-	expression tag	UNP Q92SG7
D	-1	GLN	-	expression tag	UNP Q92SG7
D	0	GLY	-	expression tag	UNP $Q92SG7$
D	215	MSE	VAL	cloning artifact	UNP Q92SG7
D	297	GLY	-	expression tag	UNP Q92SG7
D	298	SER	_	expression tag	UNP Q92SG7





• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Cl 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	103	Total O 103 103	0	0
3	В	90	Total O 90 90	0	0
3	С	73	Total         O           73         73	0	0
3	D	97	Total O 97 97	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Putative transcriptional regulator







## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	67.34Å 83.33Å 85.07Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $91.65^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	36.42 - 1.88	Depositor
Resolution (A)	36.42 - 1.88	EDS
% Data completeness	93.8 (36.42-1.88)	Depositor
(in resolution range)	99.2 (36.42-1.88)	EDS
R <sub>merge</sub>	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.01 (at 1.88 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
B B.	0.184 , $0.224$	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.184 , $0.222$	DCC
$R_{free}$ test set	3817 reflections $(5.02%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	26.4	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37, 57.1	EDS
L-test for $twinning^2$	$< L >=0.49, < L^2>=0.32$	Xtriage
	0.000 for -h,l,k	
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
	0.027 for h,-k,-l	
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7104	wwPDB-VP
Average B, all atoms $(Å^2)$	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 74.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4630e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.36	0/1720	0.56	0/2322	
1	В	0.41	0/1747	0.57	0/2364	
1	С	0.37	0/1666	0.56	0/2248	
1	D	0.37	0/1768	0.58	0/2390	
All	All	0.38	0/6901	0.57	0/9324	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1685	0	1690	36	0
1	В	1704	0	1720	35	0
1	С	1629	0	1631	45	0
1	D	1722	0	1744	40	0
2	D	1	0	0	0	0
3	А	103	0	0	1	0
3	В	90	0	0	1	0
3	С	73	0	0	2	0
3	D	97	0	0	1	0
All	All	7104	0	6785	139	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (139)	$\operatorname{close}$	$\operatorname{contacts}$	within	the	same	asymmetric	unit	are	listed	below,	sorted	by	their	$\operatorname{clash}$
magnitud	e.													

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap (Å)
1:A:98:ASN:HD21	1:B:222[B]:THR:HG22	1.01	1.08
1:B:272:ARG:HG2	1:B:272:ARG:HH11	1.10	1.08
1:C:98:ASN:HD21	1:D:222:THR:HG22	1.23	1.03
1:D:224:LEU:HD12	1:D:240:MSE:HE2	1.43	1.00
1:A:98:ASN:ND2	1:B:222[B]:THR:HG22	1.77	0.98
1:B:272:ARG:HH11	1:B:272:ARG:CG	1.88	0.85
1:D:151[A]:ARG:NH2	1:D:154:GLU:HG2	1.92	0.84
1:C:98:ASN:ND2	1:D:222:THR:HG22	1.91	0.84
1:C:224:LEU:HD12	1:C:240:MSE:HE2	1.56	0.84
1:D:151[A]:ARG:HH21	1:D:154:GLU:HG2	1.40	0.84
1:A:98:ASN:HD21	1:B:222[B]:THR:CG2	1.88	0.83
1:C:87:ARG:HH11	1:C:87:ARG:HB3	1.42	0.83
1:D:275:SER:O	1:D:279:ARG:HG3	1.78	0.83
1:B:272:ARG:HG2	1:B:272:ARG:NH1	1.92	0.82
1:A:146:SER:O	1:A:270:PRO:HG3	1.81	0.81
1:D:190:PRO:HG3	1:D:218:ASN:HD22	1.45	0.80
1:C:104:ALA:O	1:C:107:GLU:HG2	1.81	0.79
1:C:222:THR:HG22	1:D:98:ASN:HD21	1.51	0.74
1:B:272:ARG:O	1:B:274:LEU:HD13	1.91	0.70
1:A:121:VAL:HG21	1:B:222[B]:THR:HG21	1.72	0.70
1:C:189:LEU:HD13	1:C:191:LYS:HD3	1.74	0.69
1:D:155:MSE:HE3	1:D:157:PHE:CZ	2.29	0.68
1:C:222:THR:HG22	1:D:98:ASN:ND2	2.11	0.66
1:D:127:ASP:HB3	1:D:143:THR:OG1	1.95	0.66
1:C:98:ASN:HD21	1:D:222:THR:CG2	2.04	0.66
1:C:175:PRO:HG2	1:C:207:ILE:HD11	1.78	0.66
1:C:199:HIS:CD2	1:C:208:GLU:HB2	2.31	0.65
1:B:126:ILE:HD12	3:B:379:HOH:O	1.97	0.65
1:A:107:GLU:HG3	1:A:108:PHE:N	2.11	0.64
1:B:219:GLU:OE1	1:B:222[B]:THR:HG23	1.98	0.64
1:D:190:PRO:HG3	1:D:218:ASN:ND2	2.13	0.62
1:D:151[A]:ARG:HH21	1:D:154:GLU:CG	2.11	0.62
1:A:224:LEU:HD22	1:A:240:MSE:HE2	1.81	0.61
1:A:104:ALA:O	1:A:107:GLU:HG2	2.02	0.59
1:A:125:THR:HG22	1:A:189:LEU:HD23	1.85	0.59
1:C:87:ARG:HB3	1:C:87:ARG:NH1	2.17	0.59
1:C:172:PRO:HG3	1:C:182[A]:CYS:SG	2.43	0.58



2	$\Lambda /$	7	1
J	TAT		Т

		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:B:94:SER:OG	1:B:219:GLU:OE2	2.21	0.58
1:D:189:LEU:HB2	1:D:194:GLN:HG2	1.84	0.58
1:C:212:ARG:HB3	1:C:212:ARG:NH2	2.18	0.57
1:C:85:LYS:NZ	1:C:85:LYS:HB3	2.19	0.57
1:A:222:THR:HG23	1:B:102:ILE:HD11	1.86	0.57
1:A:158:VAL:HG21	1:A:238:LEU:HD13	1.89	0.55
1:D:238:LEU:O	1:D:242[A]:ARG:HG3	2.07	0.55
1:D:224:LEU:HD11	1:D:240:MSE:HB3	1.89	0.55
1:A:189:LEU:HG	1:A:190:PRO:HD2	1.89	0.55
1:A:187:TYR:HE2	1:A:189:LEU:HD12	1.71	0.54
1:B:272:ARG:CG	1:B:272:ARG:NH1	2.56	0.54
1:C:121:VAL:HG21	1:D:222:THR:HG21	1.88	0.54
1:B:122:SER:OG	1:B:124:ARG:HG2	2.07	0.54
1:B:130:ALA:HB3	1:B:133:VAL:HG23	1.90	0.53
1:C:219:GLU:OE1	1:C:222:THR:HG23	2.08	0.53
1:D:199:HIS:HE1	1:D:208:GLU:OE1	1.91	0.53
1:C:189:LEU:HB3	1:C:191:LYS:HG3	1.91	0.53
1:C:212:ARG:HB3	1:C:212:ARG:CZ	2.40	0.52
1:D:142:LEU:HD11	1:D:149:ALA:CB	2.40	0.51
1:C:107:GLU:HG3	1:C:108:PHE:N	2.25	0.51
1:A:274:LEU:HG	1:A:279:ARG:CG	2.41	0.51
1:C:163:ARG:HG3	1:C:248:GLY:O	2.11	0.51
1:C:222:THR:HG21	1:D:121:VAL:HG11	1.93	0.50
1:A:222:THR:HG23	1:B:102:ILE:CD1	2.41	0.50
1:D:194:GLN:O	1:D:194:GLN:HG3	2.12	0.50
1:C:129:LEU:HD12	1:C:129:LEU:H	1.77	0.49
1:B:143:THR:HG23	1:B:144:ASP:OD1	2.13	0.49
1:C:222:THR:HA	1:D:98:ASN:HD21	1.77	0.49
1:A:254:LEU:N	1:A:255:PRO:HD3	2.27	0.49
1:B:172:PRO:HG3	1:B:182:CYS:SG	2.53	0.48
1:D:244:ASP:HB3	1:D:250:MSE:HG2	1.95	0.48
1:A:154:GLU:HG2	1:A:261:PRO:HB2	1.95	0.48
1:D:142:LEU:HD11	1:D:149:ALA:HB2	1.95	0.48
1:C:272:ARG:HD3	1:C:272:ARG:HA	1.69	0.47
1:D:243:GLU:HB3	3:D:333:HOH:O	2.14	0.47
1:D:199:HIS:CE1	1:D:208:GLU:OE1	2.68	0.47
1:B:155:MSE:HE2	1:B:264:ILE:HD11	1.97	0.47
1:A:111[B]:LYS:HG3	3:A:328:HOH:O	2.14	0.46
1:C:212:ARG:CZ	1:C:212:ARG:CB	2.93	0.46
1:D:155:MSE:HE3	1:D:157:PHE:HZ	1.78	0.46
1:C:151:ARG:NH1	1:C:154:GLU:HG3	2.31	0.46



3MZ1
------

	<b>h</b>   <b>0</b>	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:B:126:ILE:O	1:B:138:ARG:NH2	2.47	0.46
1:A:185:VAL:HB	1:A:234:ILE:HG22	1.98	0.45
1:C:189:LEU:HB3	1:C:191:LYS:CG	2.46	0.45
1:D:126:ILE:HG13	1:D:138:ARG:HB2	1.98	0.45
1:D:275:SER:O	1:D:279:ARG:CG	2.57	0.45
1:A:224:LEU:CD2	1:A:240:MSE:HE2	2.48	0.44
1:A:213:TYR:O	1:B:116:GLN:HG2	2.16	0.44
1:D:194:GLN:O	1:D:194:GLN:CG	2.66	0.44
1:A:102:ILE:HD11	1:A:119:LEU:CD2	2.47	0.44
1:B:126:ILE:O	1:B:126:ILE:HG22	2.17	0.44
1:D:85:LYS:HD2	1:D:114:ASP:HB3	1.99	0.44
1:D:247[B]:ASN:HD22	1:D:249:THR:H	1.66	0.44
1:A:137:ILE:HD12	1:A:137:ILE:N	2.33	0.43
1:C:183:TYR:HE1	1:C:212:ARG:HH22	1.57	0.43
1:B:243:GLU:HA	1:B:246:ARG:NH2	2.34	0.43
1:A:142:LEU:HD13	1:A:142:LEU:HA	1.67	0.43
1:A:196:MSE:HA	1:A:197:PRO:HD3	1.94	0.43
1:A:238:LEU:HG	1:A:242[A]:ARG:HE	1.84	0.43
1:B:132:ASN:OD1	1:B:132:ASN:N	2.52	0.43
1:C:200:PHE:HB2	1:C:207:ILE:CD1	2.49	0.43
1:D:254:LEU:N	1:D:255:PRO:HD3	2.34	0.43
1:A:158:VAL:CG2	1:A:238:LEU:HD13	2.48	0.42
1:A:85:LYS:HG2	1:A:114:ASP:HB2	2.01	0.42
1:C:189:LEU:CD1	1:C:191:LYS:HD3	2.44	0.42
1:C:274:LEU:HG	1:C:279:ARG:HG3	2.00	0.42
1:C:95:ALA:O	1:C:99:LEU:HB2	2.19	0.42
1:D:96:PHE:CD1	1:D:289:MSE:HE1	2.54	0.42
1:D:197:PRO:HB2	1:D:208:GLU:HG3	2.02	0.42
1:C:182[B]:CYS:HA	3:C:305:HOH:O	2.18	0.42
1:B:124:ARG:O	1:B:124:ARG:HG3	2.18	0.42
1:C:183:TYR:CE1	1:C:212:ARG:NH2	2.73	0.42
1:C:85:LYS:HB3	1:C:85:LYS:HZ2	1.83	0.42
1:B:126:ILE:HG21	1:B:129:LEU:HD21	2.02	0.42
1:C:222:THR:CG2	1:D:98:ASN:HD21	2.26	0.42
1:A:131:GLU:OE2	1:A:271:ASN:ND2	2.53	0.42
1:A:222:THR:CG2	1:B:102:ILE:HD11	2.49	0.42
1:B:243:GLU:HG2	1:B:247:ASN:HD21	1.84	0.42
1:B:91:GLU:OE2	1:B:120:GLY:HA3	2.20	0.41
1:B:164:ASP:OD2	1:B:168:ARG:NH1	2.53	0.41
1:D:131:GLU:O	1:D:132:ASN:HB2	2.20	0.41
1:C:94:SER:OG	1:C:219:GLU:OE2	2.38	0.41



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:105:LEU:N	1:C:106:PRO:CD	2.83	0.41
1:B:137:ILE:HD12	1:B:137:ILE:N	2.35	0.41
1:C:200:PHE:HB2	1:C:207:ILE:HD11	2.01	0.41
1:A:138:ARG:HD3	1:A:138:ARG:HA	1.84	0.41
1:A:274:LEU:HG	1:A:279:ARG:HD3	2.02	0.41
1:C:157:PHE:HE2	1:C:262:MSE:HB2	1.85	0.41
1:C:197:PRO:HB2	1:C:208:GLU:HG3	2.03	0.41
1:A:144:ASP:HB3	1:A:147:LEU:CD1	2.51	0.41
1:B:199:HIS:CE1	1:B:208:GLU:HB2	2.56	0.41
1:A:215:MSE:HE3	1:A:217:ALA:HB2	2.03	0.41
1:B:143:THR:HG23	1:B:144:ASP:N	2.36	0.41
1:D:247[B]:ASN:ND2	1:D:249:THR:H	2.19	0.41
1:A:244:ASP:HB3	1:A:250:MSE:HG2	2.02	0.40
1:B:105:LEU:N	1:B:106:PRO:CD	2.84	0.40
1:A:269:PRO:HG2	1:A:272:ARG:HH21	1.86	0.40
1:C:88:LEU:HD21	1:C:277:ARG:HG3	2.02	0.40
1:C:182[A]:CYS:HA	3:C:305:HOH:O	2.20	0.40
1:C:189:LEU:HD23	1:C:189:LEU:HA	1.72	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	205/300~(68%)	200~(98%)	5(2%)	0	100 100
1	В	213/300~(71%)	209~(98%)	4 (2%)	0	100 100
1	С	198/300~(66%)	193~(98%)	5(2%)	0	100 100
1	D	215/300~(72%)	212~(99%)	3(1%)	0	100 100
All	All	831/1200 (69%)	814 (98%)	17 (2%)	0	100 100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	182/246~(74%)	174 (96%)	8 (4%)	24 9
1	В	186/246~(76%)	181 (97%)	5(3%)	40 24
1	С	176/246~(72%)	162~(92%)	14 (8%)	10 2
1	D	188/246~(76%)	180 (96%)	8 (4%)	25 9
All	All	732/984~(74%)	697~(95%)	35~(5%)	21 7

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	102	ILE
1	А	125	THR
1	А	142	LEU
1	А	163	ARG
1	А	189	LEU
1	А	219	GLU
1	А	268	TYR
1	А	271	ASN
1	В	125	THR
1	В	138	ARG
1	В	151	ARG
1	В	272	ARG
1	В	274	LEU
1	С	85	LYS
1	С	87	ARG
1	С	99	LEU
1	С	121	VAL
1	С	128	TYR
1	С	129	LEU
1	С	131	GLU
1	С	132	ASN
1	С	141	THR
1	С	150	ARG
1	Ċ	151	ARG



Mol	Chain	Res	Type
1	С	196	MSE
1	С	224	LEU
1	С	256	ASP
1	D	83	LEU
1	D	123	ASP
1	D	129	LEU
1	D	191	LYS
1	D	195	GLN
1	D	196	MSE
1	D	219	GLU
1	D	224	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	98	ASN
1	А	145	GLN
1	А	181	ASN
1	А	271	ASN
1	В	98	ASN
1	В	247	ASN
1	В	258	GLN
1	В	293	GLN
1	С	98	ASN
1	С	194	GLN
1	D	98	ASN
1	D	116	GLN
1	D	199	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



### 5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	201/300~(67%)	0.28	19 (9%) 15 18	16, 30, 75, 119	3(1%)
1	В	205/300~(68%)	0.20	14 (6%) 25 26	11, 30, 73, 108	3 (1%)
1	С	197/300~(65%)	0.40	16 (8%) 19 22	20, 36, 81, 99	2(1%)
1	D	205/300~(68%)	0.16	12 (5%) 29 30	14, 28, 68, 104	5 (2%)
All	All	808/1200 (67%)	0.26	61 (7%) 22 23	11, 31, 77, 119	13 (1%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res   Type		RSRZ
1	А	274	LEU	7.1
1	С	128	TYR	5.5
1	D	189	LEU	5.5
1	А	189	LEU	5.4
1	С	190	PRO	5.2
1	D	83	LEU	4.9
1	С	129	LEU	4.7
1	С	124	ARG	4.7
1	А	273	HIS	4.6
1	В	128	TYR	4.6
1	С	130	ALA	4.3
1	D	193	GLY	4.0
1	А	125	THR	4.0
1	С	189	LEU	3.8
1	D	190	PRO	3.5
1	С	123	ASP	3.4
1	D	126	ILE	3.4
1	D	123	ASP	3.3
1	С	141	THR	3.3
1	D	128	TYR	3.3
1	D	192	THR	3.2



Mol	Chain	Res	Type	RSRZ	
1	В	189	LEU	3.2	
1	В	192	THR	3.1	
1	А	272	ARG	3.1	
1	В	126	ILE	3.0	
1	А	128	TYR	3.0	
1	А	83	LEU	3.0	
1	В	271	ASN	2.9	
1	А	190	PRO	2.8	
1	А	194	GLN	2.8	
1	С	171	VAL	2.7	
1	В	130	ALA	2.7	
1	В	193	GLY	2.6	
1	D	127	ASP	2.6	
1	С	204	ASN	2.5	
1	В	131	GLU	2.5	
1	С	194	GLN	2.5	
1	D	124	ARG	2.4	
1	А	143	THR	2.4	
1	В	143	THR	2.4	
1	С	131	GLU	2.4	
1	С	247	ASN	2.4	
1	В	83	LEU	2.4	
1	А	270	PRO	2.4	
1	А	154	GLU	2.4	
1	В	274	LEU	2.3	
1	С	163	ARG	2.3	
1	В	107	GLU	2.3	
1	А	85	LYS	2.3	
1	А	144	ASP	2.3	
1	С	207	ILE	2.2	
1	A	130	ALA	2.2	
1	В	194	GLN	2.2	
1	В	272	ARG	2.2	
1	D	85	LYS	2.1	
1	A	271	ASN	2.1	
1	С	212	ARG	2.1	
1	A	124	ARG	2.1	
1	A	142	LEU	2.0	
1	D	194	GLN	2.0	
1	А	294	ASN	2.0	

Continued from previous page...



#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q<0.9
2	CL	D	299	1/1	0.99	0.06	24,24,24,24	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

