



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 07:51 pm GMT

PDB ID : 5MZV
Title : IL-23:IL-23R:Nb22E11 complex
Authors : Bloch, Y.; Savvides, S.N.
Deposited on : 2017-02-01
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2792 (2.80-2.80)
Clashscore	122126	3209 (2.80-2.80)
Ramachandran outliers	120053	3158 (2.80-2.80)
Sidechain outliers	120020	3160 (2.80-2.80)
RSRZ outliers	108989	2726 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	 91% 8%
2	B	198	 67% 31%
3	C	330	 84% 11%
4	D	156	 76% 21%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 13889 atoms, of which 6773 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-12 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	303	4689	1515	2290	395	477	12	0	0	0

- Molecule 2 is a protein called Interleukin-23 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	137	2108	674	1047	190	192	5	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	190	GLY	-	expression tag	UNP Q9NPF7
B	191	THR	-	expression tag	UNP Q9NPF7
B	192	LYS	-	expression tag	UNP Q9NPF7
B	193	HIS	-	expression tag	UNP Q9NPF7
B	194	HIS	-	expression tag	UNP Q9NPF7
B	195	HIS	-	expression tag	UNP Q9NPF7
B	196	HIS	-	expression tag	UNP Q9NPF7
B	197	HIS	-	expression tag	UNP Q9NPF7
B	198	HIS	-	expression tag	UNP Q9NPF7

- Molecule 3 is a protein called Interleukin-23 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
3	C	293	4729	1538	2331	392	448	20	0	4	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	318	ILE	-	expression tag	UNP Q5VWK5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	319	GLU	-	expression tag	UNP Q5VWK5
C	320	GLY	-	expression tag	UNP Q5VWK5
C	321	ARG	-	expression tag	UNP Q5VWK5
C	322	GLY	-	expression tag	UNP Q5VWK5
C	323	THR	-	expression tag	UNP Q5VWK5
C	324	LYS	-	expression tag	UNP Q5VWK5
C	325	HIS	-	expression tag	UNP Q5VWK5
C	326	HIS	-	expression tag	UNP Q5VWK5
C	327	HIS	-	expression tag	UNP Q5VWK5
C	328	HIS	-	expression tag	UNP Q5VWK5
C	329	HIS	-	expression tag	UNP Q5VWK5
C	330	HIS	-	expression tag	UNP Q5VWK5

- Molecule 4 is a protein called Nanobody 22E11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
4	D	123	1845	589	900	168	184	4	0	1	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	C	2	Total	Cl	0	0
			2	2		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Na	0	0
			1	1		

- Molecule 7 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
7	A	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
7	A	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
7	C	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
7	C	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
7	C	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
7	C	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
7	C	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
7	C	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
7	C	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
7	C	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
7	C	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

- Molecule 8 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
8	A	1	19	6	8	5	0	0
8	C	1	21	6	10	5	0	0

- Molecule 9 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
9	A	1	20	6	9	5	0	0
9	A	1	20	6	9	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	H	O	0	0
			21	6	10	5		
9	A	1	Total	C	H	O	0	0
			21	6	10	5		



- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	29	Total	O	0	0
			29	29		
10	B	2	Total	O	0	0
			2	2		
10	C	38	Total	O	0	0
			38	38		
10	D	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Interleukin-12 subunit beta




- Molecule 2: Interleukin-23 subunit alpha



- Molecule 3: Interleukin-23 receptor



- Molecule 4: Nanobody 22E11



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.22Å 112.19Å 109.87Å 90.00° 106.24° 90.00°	Depositor
Resolution (Å)	76.85 – 2.80 76.85 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.5 (76.85-2.80) 97.9 (76.85-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.82Å)	Xtrriage
Refinement program	PHENIX (dev_2614: ???), BUSTER	Depositor
R, R_{free}	0.212 , 0.250 0.212 , 0.250	Depositor DCC
R_{free} test set	1594 reflections (4.19%)	wwPDB-VP
Wilson B-factor (Å ²)	68.7	Xtrriage
Anisotropy	0.212	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13889	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NA, BMA, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2457	0.46	0/3339
2	B	0.23	0/1084	0.37	0/1468
3	C	0.26	0/2473	0.46	0/3361
4	D	0.29	0/970	0.46	0/1315
All	All	0.26	0/6984	0.45	0/9483

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2399	2290	2288	2	0
2	B	1061	1047	1046	2	0
3	C	2398	2331	2328	7	0
4	D	945	900	901	3	0
5	A	1	0	0	0	0
5	C	2	0	0	0	0
6	A	1	0	0	0	0
7	A	28	24	24	0	0
7	C	140	125	125	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	11	8	8	0	0
8	C	11	10	10	0	0
9	A	44	38	38	0	0
10	A	29	0	0	0	0
10	B	2	0	0	0	0
10	C	38	0	0	0	0
10	D	6	0	0	0	0
All	All	7116	6773	6768	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:ARG:NH1	1:A:186:GLU:OE1	2.34	0.60
3:C:287:ASN:N	3:C:315:THR:OG1	2.37	0.58
4:D:6:GLU:OE1	4:D:6:GLU:N	2.40	0.55
1:A:85:ALA:O	4:D:106:ARG:NH2	2.42	0.53
2:B:85:ASN:OD1	2:B:87:GLN:N	2.40	0.53
7:C:408:NAG:O7	7:C:408:NAG:O3	2.25	0.52
4:D:62:ASP:OD1	4:D:63:SER:N	2.46	0.49
3:C:234:THR:OG1	3:C:235:VAL:N	2.47	0.47
3:C:221:SER:O	3:C:305:GLN:NE2	2.48	0.47
3:C:139:SER:O	7:C:408:NAG:H82	2.18	0.44
7:C:408:NAG:H62	7:C:409:NAG:N2	2.34	0.43
2:B:112:GLU:O	2:B:158:ARG:NH1	2.51	0.42
3:C:262:ASN:OD1	3:C:263:GLN:N	2.48	0.42
3:C:273:ASN:O	3:C:278:GLN:NE2	2.53	0.41
3:C:141:ASN:HB3	3:C:182:SER:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/328 (91%)	282 (94%)	17 (6%)	0	100	100
2	B	129/198 (65%)	123 (95%)	6 (5%)	0	100	100
3	C	295/330 (89%)	270 (92%)	25 (8%)	0	100	100
4	D	122/156 (78%)	118 (97%)	4 (3%)	0	100	100
All	All	845/1012 (84%)	793 (94%)	52 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/297 (91%)	269 (100%)	0	100	100
2	B	115/167 (69%)	115 (100%)	0	100	100
3	C	268/298 (90%)	265 (99%)	3 (1%)	76	93
4	D	100/122 (82%)	99 (99%)	1 (1%)	78	94
All	All	752/884 (85%)	748 (100%)	4 (0%)	90	97

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	82	LYS
3	C	141	ASN
3	C	210	LEU
4	D	95	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	125	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	266	HIS
2	B	154	GLN
4	D	39	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 4 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	NAG	A	403	1,7	14,14,15	0.34	0	17,19,21	0.44	0
7	NAG	A	404	8,7	14,14,15	0.26	0	17,19,21	0.44	0
8	BMA	A	405	9,7	11,11,12	0.75	0	15,15,17	0.90	0
9	MAN	A	406	9,8	11,11,12	0.69	0	15,15,17	1.10	2 (13%)
9	MAN	A	407	9,8	11,11,12	0.64	0	15,15,17	1.03	2 (13%)
9	MAN	A	408	9	11,11,12	0.60	0	15,15,17	1.01	2 (13%)
9	MAN	A	409	9	11,11,12	0.72	0	15,15,17	0.89	1 (6%)
7	NAG	C	403	3,7	14,14,15	0.38	0	17,19,21	0.51	0
7	NAG	C	404	7	14,14,15	0.25	0	17,19,21	0.58	0
7	NAG	C	405	3,7	14,14,15	0.61	1 (7%)	17,19,21	0.80	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	C	406	8,7	14,14,15	0.24	0	17,19,21	0.77	1 (5%)
8	BMA	C	407	7	11,11,12	0.58	0	15,15,17	0.77	0
7	NAG	C	408	3,7	14,14,15	0.73	1 (7%)	17,19,21	0.83	0
7	NAG	C	409	7	14,14,15	0.29	0	17,19,21	0.60	0
7	NAG	C	410	3,7	14,14,15	0.37	0	17,19,21	0.49	0
7	NAG	C	411	7	14,14,15	0.27	0	17,19,21	0.54	0
7	NAG	C	412	3	14,14,15	0.22	0	17,19,21	0.52	0
7	NAG	C	413	3	14,14,15	0.37	0	17,19,21	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	403	1,7	-	0/6/23/26	0/1/1/1
7	NAG	A	404	8,7	-	0/6/23/26	0/1/1/1
8	BMA	A	405	9,7	-	0/2/19/22	0/1/1/1
9	MAN	A	406	9,8	-	0/2/19/22	0/1/1/1
9	MAN	A	407	9,8	-	0/2/19/22	0/1/1/1
9	MAN	A	408	9	-	0/2/19/22	0/1/1/1
9	MAN	A	409	9	-	0/2/19/22	0/1/1/1
7	NAG	C	403	3,7	-	0/6/23/26	0/1/1/1
7	NAG	C	404	7	-	0/6/23/26	0/1/1/1
7	NAG	C	405	3,7	-	0/6/23/26	0/1/1/1
7	NAG	C	406	8,7	-	0/6/23/26	0/1/1/1
8	BMA	C	407	7	-	0/2/19/22	0/1/1/1
7	NAG	C	408	3,7	-	0/6/23/26	0/1/1/1
7	NAG	C	409	7	-	0/6/23/26	0/1/1/1
7	NAG	C	410	3,7	-	0/6/23/26	0/1/1/1
7	NAG	C	411	7	-	0/6/23/26	0/1/1/1
7	NAG	C	412	3	-	0/6/23/26	0/1/1/1
7	NAG	C	413	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	408	NAG	O5-C1	-2.54	1.39	1.43
7	C	405	NAG	O5-C1	-2.01	1.40	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	406	MAN	O2-C2-C3	-2.79	104.76	110.19
9	A	407	MAN	O2-C2-C3	-2.25	105.79	110.19
9	A	408	MAN	O2-C2-C3	-2.22	105.85	110.19
9	A	409	MAN	C1-O5-C5	2.06	115.02	112.19
9	A	408	MAN	C1-O5-C5	2.45	115.56	112.19
9	A	407	MAN	C1-O5-C5	2.53	115.67	112.19
9	A	406	MAN	C1-O5-C5	2.56	115.71	112.19
7	C	406	NAG	C1-O5-C5	2.74	115.95	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	408	NAG	3	0
7	C	409	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/328 (92%)	0.17	2 (0%) 87 83	43, 77, 128, 161	0
2	B	137/198 (69%)	0.33	5 (3%) 42 32	67, 108, 136, 156	0
3	C	293/330 (88%)	0.42	15 (5%) 28 19	48, 76, 152, 194	0
4	D	123/156 (78%)	0.31	0 100 100	42, 58, 93, 108	0
All	All	856/1012 (84%)	0.30	22 (2%) 56 45	42, 79, 136, 194	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	255	MET	3.1
2	B	48	HIS	3.1
3	C	228	ALA	3.1
3	C	270	PHE	2.9
2	B	187	LEU	2.8
1	A	248	ASN	2.7
2	B	131	LEU	2.7
3	C	281	GLU	2.7
3	C	282	PHE	2.7
3	C	266	ASN	2.7
3	C	264	THR	2.6
3	C	232	ASN	2.6
3	C	241	TYR	2.5
3	C	109	PHE	2.5
3	C	271	ASP	2.5
2	B	47	ALA	2.4
3	C	229	GLU	2.4
1	A	166	GLN	2.4
2	B	135	LEU	2.3
3	C	292	PHE	2.3
3	C	291	VAL	2.0
3	C	265	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	C	413	14/15	0.76	0.28	131,141,148,149	0
7	NAG	C	412	14/15	0.76	0.28	160,167,184,184	0
9	MAN	A	407	11/12	0.80	0.18	124,134,157,158	0
8	BMA	C	407	11/12	0.80	0.14	127,132,161,161	0
7	NAG	C	409	14/15	0.81	0.15	154,163,165,170	0
9	MAN	A	408	11/12	0.82	0.17	160,171,173,177	0
7	NAG	C	410	14/15	0.82	0.15	132,139,149,153	0
7	NAG	C	404	14/15	0.83	0.21	134,142,148,152	0
7	NAG	C	406	14/15	0.84	0.12	111,114,134,136	0
7	NAG	C	411	14/15	0.84	0.26	158,165,176,179	0
9	MAN	A	409	11/12	0.85	0.19	150,159,184,184	0
9	MAN	A	406	11/12	0.86	0.12	113,121,128,129	0
8	BMA	A	405	11/12	0.86	0.16	109,117,134,137	0
7	NAG	C	405	14/15	0.89	0.18	99,111,126,126	0
7	NAG	C	403	14/15	0.89	0.23	105,108,110,114	0
7	NAG	C	408	14/15	0.92	0.15	118,126,130,131	0
5	CL	C	401	1/1	0.94	0.12	57,57,57,57	0
6	NA	A	402	1/1	0.96	0.09	57,57,57,57	0
7	NAG	A	404	14/15	0.97	0.17	34,52,59,61	0
7	NAG	A	403	14/15	0.97	0.18	34,46,56,56	0
5	CL	C	402	1/1	0.97	0.13	57,57,57,57	0
5	CL	A	401	1/1	0.99	0.16	57,57,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.