

# Full wwPDB X-ray Structure Validation Report (i)

#### Jun 22, 2024 – 11:35 PM EDT

PDB ID	:	6N1Z
Title	:	Importin-9 bound to H2A-H2B
Authors	:	Tomchick, D.R.; Chook, Y.M.; Padavannil, A.
Deposited on	:	2018-11-12
Resolution	:	2.70  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity		4 02h-467
Vtria na (Dhanim)	·	1.025 101
Atriage (Phenix)	:	1.20.1
$\mathrm{EDS}$	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069(2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Qualit	y of chain			
1	А	1041	3%			7%	8%
	 	10.11	3%			1.10	0.0
	D	1041	9%			7%	9%
2	В	130	62%	5%		34%	
2	E	130	3% 62%			35%	
-		100	6%	•		0078	
3	С	123	67%		7% •	25%	



Mol	Chain	Length		Quality of chain		
			10%			
3	$\mathbf{F}$	123		74%	5%	21%



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 36201 atoms, of which 18040 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	Λ	061	Total	С	Η	Ν	Ο	S	0	1	0
	A	901	15105	4818	7575	1251	1415	46	0		0
1	л	040	Total	С	Η	Ν	Ο	S	0	1	0
		949	14945	4766	7503	1242	1388	46		1	U

• Molecule 1 is a protein called Importin-9.

• Molecule 2 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	86	Total	С	Η	Ν	Ο	0	0	0
	D	80	1381	421	708	134	118	0	0	0
9	F	85	Total	С	Н	Ν	Ο	0	0	0
	Ľ	85	1367	417	701	133	116	0	U	

• Molecule 3 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	С	92	Total 1471	C 456	Н 746	N 132	0 135	$\begin{array}{c} \mathrm{S} \\ \mathrm{2} \end{array}$	0	0	0
3	F	97	Total 1577	C 484	H 807	N 143	0 141	${S \over 2}$	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	1	MET	-	initiating methionine	UNP P02281
С	30	THR	SER	engineered mutation	UNP P02281
F	1	MET	-	initiating methionine	UNP P02281
F	30	THR	SER	engineered mutation	UNP P02281

• Molecule 4 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	163	Total O 163 163	0	0
4	В	2	Total O 2 2	0	0
4	С	5	Total O 5 5	0	0
4	D	179	Total O 179 179	0	0
4	Ε	3	Total O 3 3	0	0
4	F	3	Total O 3 3	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Importin-9







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	127.42Å 223.29Å 131.83Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	45.44 - 2.70	Depositor
Resolution (A)	45.44 - 2.70	EDS
% Data completeness	90.5 (45.44-2.70)	Depositor
(in resolution range)	90.6 (45.44-2.70)	EDS
R <sub>merge</sub>	0.09	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.81 (at 2.69 \text{\AA})$	Xtriage
Refinement program	PHENIX	Depositor
D D.	0.209 , $0.238$	Depositor
$\Lambda, \Lambda_{free}$	0.209 , $0.233$	DCC
$R_{free}$ test set	1978 reflections $(2.11\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	38.6	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, $39.7$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.004 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	36201	wwPDB-VP
Average B, all atoms $(Å^2)$	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 27.34 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2394e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Ullalli	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.25	0/7685	0.41	0/10454	
1	D	0.25	0/7595	0.41	0/10326	
2	В	0.26	0/681	0.45	0/918	
2	Ε	0.26	0/674	0.41	0/908	
3	С	0.24	0/734	0.40	0/984	
3	F	0.25	0/781	0.42	0/1047	
All	All	0.25	0/18150	0.41	0/24637	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	7530	7575	7574	41	0
1	D	7442	7503	7503	48	0
2	В	673	708	708	3	0
2	Е	666	701	701	9	0
3	С	725	746	746	6	0
3	F	770	807	807	8	0
4	А	163	0	0	0	0
4	В	2	0	0	0	0
4	С	5	0	0	0	0
4	D	179	0	0	0	0



	Jerry							
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes		
4	Е	3	0	0	0	0		
4	F	3	0	0	0	0		
All	All	18161	18040	18039	104	0		

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:E:101:VAL:HG11	3:F:66:ASP:HB3	1.40	1.01
1:D:999:LEU:HD23	1:D:1002:ILE:HD12	1.61	0.82
1:D:701:THR:HG21	1:D:713:GLY:HA3	1.72	0.70
1:D:71:LEU:CD2	1:D:121:ILE:HD13	2.23	0.68
1:D:237:ILE:O	1:D:240:VAL:HG22	1.94	0.67
1:D:33:LEU:HD21	1:D:76:LEU:HD22	1.77	0.67
1:A:725:LEU:HD22	1:A:773:LYS:HB3	1.77	0.66
1:A:52:VAL:HG13	1:A:99:PRO:HD3	1.78	0.65
1:A:753:PRO:O	1:A:795:GLN:NE2	2.30	0.64
1:D:345:GLU:OE2	3:F:28:ARG:NH2	2.32	0.63
1:D:407:LEU:O	1:D:411:THR:HG23	2.00	0.62
1:D:114:PRO:HB3	1:D:150:LEU:HD21	1.81	0.61
1:D:725:LEU:HD22	1:D:773:LYS:HB3	1.83	0.60
1:D:71:LEU:HD21	1:D:121:ILE:CD1	2.32	0.60
1:D:386:VAL:HG21	1:D:579:VAL:HG22	1.84	0.59
1:D:71:LEU:HD21	1:D:121:ILE:HD13	1.85	0.59
1:A:22:LYS:NZ	1:A:62:GLU:OE1	2.35	0.59
2:E:101:VAL:HG12	2:E:101:VAL:O	2.01	0.59
2:E:101:VAL:HG22	3:F:70:ARG:HD3	1.85	0.58
1:A:886:GLU:OE2	1:A:919:LYS:NZ	2.36	0.58
1:D:236:LEU:HD23	1:D:236:LEU:O	2.03	0.58
2:E:100:ARG:NH1	3:F:99:LEU:O	2.38	0.57
2:E:101:VAL:CG1	3:F:66:ASP:HB3	2.27	0.57
1:A:128:VAL:O	1:A:132:VAL:HG23	2.04	0.57
1:D:999:LEU:HD23	1:D:1002:ILE:CD1	2.34	0.57
1:D:697:VAL:O	1:D:701:THR:HG23	2.04	0.57
1:A:757:GLU:OE2	1:A:797:GLU:N	2.36	0.56
1:A:724:THR:HG23	1:A:724:THR:O	2.06	0.55
1:D:1039:ILE:HG22	1:D:1039:ILE:O	2.07	0.55
2:E:101:VAL:HG22	3:F:70:ARG:CD	2.37	0.54
1:D:757:GLU:OE2	1:D:797:GLU:N	2.38	0.54



	A 4 ama 2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:D:997:ASP:N	1:D:997:ASP:OD1	2.40	0.54
1:D:1009:THR:HG23	1:D:1039:ILE:HG23	1.89	0.54
2:B:51:TYR:HB3	3:C:92:ILE:HD11	1.91	0.53
1:A:62:GLU:HG2	1:A:112:LEU:HD21	1.90	0.53
1:D:925:LEU:HD21	1:D:1039:ILE:HG13	1.91	0.53
1:D:238:PHE:N	1:D:239:PRO:HD2	2.23	0.53
1:D:416:GLU:OE1	1:D:416:GLU:N	2.36	0.53
1:A:275:LEU:HB3	1:A:283:MET:SD	2.49	0.53
1:D:52:VAL:HG12	1:D:52:VAL:O	2.09	0.53
1:D:1009:THR:HG23	1:D:1039:ILE:CG2	2.39	0.53
1:A:874:ASP:O	1:A:876:LYS:HG3	2.09	0.52
1:A:237:ILE:HG12	1:A:241:VAL:HG23	1.92	0.52
1:A:525:LEU:HD11	1:A:541:ILE:CD1	2.39	0.52
1:D:33:LEU:CD2	1:D:76:LEU:HD22	2.39	0.52
1:D:735:GLN:N	1:D:735:GLN:OE1	2.42	0.52
1:D:75:GLN:O	1:D:79:VAL:HG23	2.10	0.51
1:A:867:LEU:HD12	1:A:914:LEU:HD22	1.92	0.51
1:D:386:VAL:HG22	1:D:535:ILE:HG21	1.91	0.51
1:D:757:GLU:HG2	1:D:801:VAL:HG21	1.93	0.51
1:D:364:ILE:HG22	1:D:368:MET:HE3	1.92	0.51
1:A:854:TYR:HH	1:A:997:ASP:N	2.09	0.50
1:A:1009:THR:HG23	1:A:1039:ILE:CG2	2.41	0.50
2:B:48:ALA:HB3	2:B:49:PRO:HD3	1.94	0.49
1:D:138:TRP:O	2:E:76:LYS:NZ	2.45	0.49
1:A:33:LEU:HD21	1:A:76:LEU:HD13	1.95	0.49
1:D:724:THR:O	1:D:724:THR:HG23	2.12	0.49
1:A:757:GLU:HG2	1:A:801:VAL:HG21	1.95	0.49
3:C:97:ARG:NE	3:C:105:ALA:HB1	2.28	0.49
3:C:92:ILE:O	3:C:96:VAL:HG23	2.12	0.48
1:A:155:LEU:HA	1:A:163:VAL:HG12	1.95	0.48
1:A:393:THR:O	3:C:27:ARG:NH1	2.47	0.47
2:B:97:LEU:HD12	2:B:97:LEU:O	2.15	0.47
1:A:81:LEU:HD21	1:A:113:LEU:HD21	1.98	0.46
1:D:81:LEU:HD21	1:D:113:LEU:HD21	1.98	0.46
1:D:201:GLU:HG2	1:D:202:VAL:HG23	1.98	0.46
1:A:62:GLU:CG	1:A:112:LEU:HD21	2.45	0.46
3:C:88:THR:OG1	3:C:89:SER:N	2.49	0.46
1:A:52:VAL:HG12	1:A:52:VAL:O	2.16	0.45
1:A:783:ASP:O	1:A:787:ARG:HG3	2.16	0.45
1:A:928:VAL:HG21	1:A:1004:LEU:HD23	1.97	0.45
1:D:52:VAL:HG13	1:D:99:PRO:HD3	1.98	0.45



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1009:THR:HG23	1:A:1039:ILE:HG21	1.97	0.45
2:E:48:ALA:HB1	3:F:92:ILE:HD12	1.97	0.45
1:D:368:MET:HE1	1:D:406:LEU:HD22	1.98	0.45
1:A:75:GLN:O	1:A:79:VAL:HG23	2.17	0.44
1:D:778:LEU:HD23	1:D:778:LEU:N	2.32	0.44
1:D:457:VAL:HG23	1:D:457:VAL:O	2.17	0.44
1:D:513:LEU:HD22	1:D:517:PHE:CE2	2.52	0.44
1:D:64:THR:HG21	1:D:78:SER:HA	1.99	0.44
1:D:701:THR:CG2	1:D:713:GLY:HA3	2.45	0.44
1:A:903:LYS:O	1:A:903:LYS:HG2	2.17	0.44
1:A:186:ALA:HB3	1:A:187:PRO:HD3	2.00	0.43
1:A:735:GLN:OE1	1:A:735:GLN:N	2.52	0.43
2:E:48:ALA:HB3	2:E:49:PRO:HD3	1.99	0.43
1:D:338:GLU:OE1	3:F:28:ARG:NH1	2.51	0.42
1:A:325:GLU:N	1:A:325:GLU:OE1	2.53	0.42
1:D:695:PRO:HG3	1:D:731:TRP:CZ2	2.54	0.42
1:A:782:LEU:C	1:A:782:LEU:HD23	2.40	0.42
1:D:71:LEU:HD22	1:D:121:ILE:HD13	2.00	0.42
1:A:386:VAL:HG21	1:A:579:VAL:HG22	2.02	0.42
3:C:28:ARG:O	3:C:32:LYS:NZ	2.44	0.42
1:A:541:ILE:HD13	1:A:564:ILE:HG23	2.02	0.41
1:D:237:ILE:O	1:D:237:ILE:HG22	2.19	0.41
1:A:757:GLU:OE2	1:A:798:THR:N	2.31	0.41
1:D:246:GLU:O	1:D:250:GLN:HG2	2.20	0.41
1:A:71:LEU:HD23	1:A:121:ILE:HD13	2.03	0.41
1:A:386:VAL:HG22	1:A:535:ILE:HG21	2.02	0.41
1:A:814:HIS:NE2	1:A:861:VAL:HG23	2.35	0.41
1:D:128:VAL:O	1:D:132:VAL:HG23	2.20	0.40
1:A:773:LYS:HA	1:A:773:LYS:HD2	1.92	0.40
1:A:836:LEU:HD23	1:A:881:ILE:HD11	2.02	0.40
1:A:525:LEU:HD11	1:A:541:ILE:HD11	2.02	0.40
1:D:895:ILE:O	1:D:900:LYS:CE	2.69	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	956/1041~(92%)	932~(98%)	23~(2%)	1 (0%)	51	78
1	D	940/1041~(90%)	912~(97%)	26~(3%)	2~(0%)	47	73
2	В	84/130~(65%)	81 (96%)	3 (4%)	0	100	100
2	Ε	83/130~(64%)	81 (98%)	2(2%)	0	100	100
3	С	86/123~(70%)	84 (98%)	2(2%)	0	100	100
3	F	95/123~(77%)	90~(95%)	5(5%)	0	100	100
All	All	2244/2588 (87%)	2180 (97%)	61 (3%)	3~(0%)	51	78

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	875	ASP
1	D	238	PHE
1	D	828	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	832/894~(93%)	824 (99%)	8 (1%)	76	91
1	D	821/894~(92%)	812 (99%)	9 (1%)	73	90
2	В	68/102~(67%)	66~(97%)	2(3%)	42	71
2	Ε	67/102~(66%)	67~(100%)	0	100	100
3	С	79/103~(77%)	76~(96%)	3~(4%)	33	62
3	F	84/103 (82%)	83~(99%)	1 (1%)	71	88
All	All	1951/2198 (89%)	1928 (99%)	23 (1%)	71	88

All (23) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	256	ASP
1	А	430	GLN
1	А	458	LYS
1	А	558	GLN
1	А	776	ARG
1	А	827	LEU
1	А	874	ASP
1	А	899	SER
2	В	98	LEU
2	В	102	THR
3	С	26	LYS
3	С	28	ARG
3	С	88	THR
1	D	145	PRO
1	D	412	ASP
1	D	414	GLN
1	D	458	LYS
1	D	460	ILE
1	D	513	LEU
1	D	558	GLN
1	D	600	SER
1	D	904	ASN
3	F	27	ARG

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



### 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	961/1041 (92%)	0.15	36 (3%) 41 41	12, 47, 98, 153	0
1	D	949/1041 (91%)	0.11	33 (3%) 44 44	12, 42, 99, 151	0
2	В	86/130 (66%)	0.71	12 (13%) 2 1	39, 62, 114, 162	0
2	Е	85/130 (65%)	0.24	4 (4%) 31 30	29, 49, 100, 149	0
3	С	92/123~(74%)	0.53	7 (7%) 13 12	36, 70, 126, 154	0
3	F	97/123 (78%)	0.55	12 (12%) 4 3	24, 61, 137, 178	0
All	All	2270/2588 (87%)	0.19	104 (4%) 32 31	12, 47, 105, 178	0

All (104) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	1002	ILE	9.1
1	D	1000	TYR	8.3
2	В	42	GLU	6.8
1	D	327	LEU	6.5
1	D	38	GLU	5.6
3	F	81	TYR	5.2
3	F	31	ARG	5.2
1	А	1000	TYR	5.1
1	А	237	ILE	4.8
1	D	36	VAL	4.7
1	А	323	ASP	4.7
1	А	734	GLU	4.7
1	D	999	LEU	4.7
1	D	935	ARG	4.6
1	А	324	GLY	4.5
3	C	31	ARG	4.4
3	С	77	ARG	4.4
1	D	1002	ILE	4.4
1	А	234	LYS	4.3



Continued from previous page						
Mol	Chain	Res	Type	RSRZ		
1	D	778	LEU	4.3		
2	В	43	ARG	4.2		
1	А	997	ASP	4.2		
1	А	232	ALA	4.2		
3	С	81	TYR	4.1		
3	F	83	LYS	4.1		
1	D	892	ASP	3.9		
2	В	46	ALA	3.9		
1	А	325	GLU	3.8		
3	F	30	THR	3.7		
1	А	778	LEU	3.6		
1	D	905	PRO	3.6		
1	D	1001	GLN	3.6		
1	D	40	ARG	3.6		
1	D	829	GLY	3.5		
1	А	1001	GLN	3.5		
1	А	38	GLU	3.4		
2	Е	43	ARG	3.4		
3	F	82	ASN	3.4		
3	С	86	THR	3.4		
2	В	41	ALA	3.4		
1	D	37	GLN	3.3		
1	D	471	HIS	3.2		
1	D	68	GLN	3.2		
1	D	1022	ILE	3.1		
1	D	15	GLY	3.1		
1	D	908	TRP	3.1		
1	D	734	GLU	3.0		
1	А	737	HIS	3.0		
1	D	232	ALA	3.0		
1	А	887	GLU	3.0		
3	F	77	ARG	3.0		
1	D	39	VAL	2.9		
1	А	999	LEU	2.8		
2	В	40	TYR	2.7		
1	А	891	MET	2.7		
1	А	1003	ASP	2.7		
1	А	71	LEU	2.7		
1	D	236	LEU	2.7		
2	В	102	THR	2.6		
1	D	1037	GLN	2.6		
3	F	84	ARG	2.6		



Mol	Chain	Res	Type	RSRZ
1	А	829	GLY	2.6
3	F	26	LYS	2.6
1	А	236	LEU	2.5
1	А	754	ARG	2.5
1	А	233	ALA	2.5
1	D	1041	ILE	2.5
3	F	29	LYS	2.5
1	А	75	GLN	2.4
1	А	797	GLU	2.4
1	А	86	GLU	2.4
2	В	39	ASN	2.3
2	Е	96	LYS	2.3
2	В	21	ARG	2.3
1	А	466	LYS	2.3
1	D	71	LEU	2.3
1	А	36	VAL	2.3
2	В	44	VAL	2.3
3	С	90	ARG	2.3
1	D	67	PRO	2.3
1	А	68	GLN	2.3
1	А	1019	PRO	2.3
3	F	91	GLU	2.3
1	А	825	CYS	2.2
1	А	1022	ILE	2.2
2	В	33	ARG	2.2
2	В	36	ARG	2.2
1	А	785	ILE	2.2
3	F	25	LYS	2.2
2	В	99	GLY	2.2
1	D	754	ARG	2.1
3	С	75	ALA	2.1
1	А	904	ASN	2.1
1	D	241	VAL	2.1
1	А	780	GLU	2.1
1	D	884	LYS	2.1
2	Е	39	ASN	2.1
3	С	28	ARG	2.1
3	F	63	PHE	2.1
1	A	471	HIS	2.0
2	E	41	ALA	2.0
1	D	16	PRO	2.0
1	D	934	ALA	2.0



Continued from previous page...

Mol	Chain	$\mathbf{Res}$	Type	RSRZ
1	D	49	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

There are no ligands in this entry.

## 6.5 Other polymers (i)

There are no such residues in this entry.

