



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 27, 2023 – 05:49 AM EDT

PDB ID : 3N2X  
Title : Crystal structure of YagE, a prophage protein belonging to the dihydrodipicolinic acid synthase family from E. coli K12 in complex with pyruvate  
Authors : Bhaskar, V.; Kumar, P.M.; Manicka, S.; Krishnaswamy, S.  
Deposited on : 2010-05-19  
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	 90% 7% ..
1	B	298	 92% 5% ..
1	C	298	 86% 11% ..
1	D	298	 88% 9% ..

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 9576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein yagE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	Total 2263	C 1449	N 385	O 423	S 6	0	0	0
1	B	298	Total 2265	C 1450	N 388	O 421	S 6	0	0	0
1	C	298	Total 2269	C 1454	N 388	O 421	S 6	0	1	0
1	D	298	Total 2268	C 1451	N 388	O 423	S 6	0	0	0

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	123	Total	O	0	0
			123	123		
3	B	136	Total	O	0	0
			136	136		
3	C	117	Total	O	0	0
			117	117		
3	D	119	Total	O	0	0
			119	119		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein yagE

Chain A:  90% 7% ..





- Molecule 1: Uncharacterized protein yagE

Chain B:  92% 5% ..





- Molecule 1: Uncharacterized protein yagE

Chain C:  86% 11% ..



- Molecule 1: Uncharacterized protein yagE

Chain D:  88% 9% ..



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.13Å 153.55Å 55.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.98 – 2.20 48.98 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.98-2.20) 99.3 (48.98-2.20)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.37 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.179 , 0.222 0.177 , 0.220	Depositor DCC
$R_{free}$ test set	3106 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.4	Xtrriage
Anisotropy	0.634	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 33.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9576	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.91 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2019e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, KPI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.03	0/2296	1.00	9/3129 (0.3%)
1	B	0.96	0/2298	0.91	8/3131 (0.3%)
1	C	1.00	2/2303 (0.1%)	1.04	10/3138 (0.3%)
1	D	1.10	4/2301 (0.2%)	0.98	16/3135 (0.5%)
All	All	1.02	6/9198 (0.1%)	0.98	43/12533 (0.3%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	134	VAL	CB-CG2	-12.41	1.26	1.52
1	D	15	THR	CB-OG1	-12.06	1.19	1.43
1	D	134	VAL	CB-CG1	-11.99	1.27	1.52
1	D	99	GLU	CG-CD	5.44	1.60	1.51
1	C	106	GLN	CD-NE2	-5.42	1.19	1.32
1	C	134	VAL	CB-CG2	-5.04	1.42	1.52

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	73	ARG	NE-CZ-NH1	-19.66	110.47	120.30
1	A	73	ARG	NE-CZ-NH1	-19.13	110.73	120.30
1	C	73	ARG	NE-CZ-NH2	13.94	127.27	120.30
1	D	77	ASP	CB-CG-OD1	12.33	129.40	118.30
1	A	73	ARG	NE-CZ-NH2	12.14	126.37	120.30
1	A	77	ASP	CB-CG-OD1	8.87	126.28	118.30
1	A	45	LYS	CD-CE-NZ	-8.52	92.10	111.70
1	C	77	ASP	CB-CG-OD1	8.43	125.89	118.30
1	D	134	VAL	CG1-CB-CG2	-8.40	97.46	110.90
1	C	122	VAL	CG1-CB-CG2	7.56	123.00	110.90
1	B	122	VAL	CG1-CB-CG2	7.47	122.86	110.90

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	292	ASP	CB-CG-OD1	7.38	124.94	118.30
1	D	122	VAL	CG1-CB-CG2	7.24	122.49	110.90
1	D	15	THR	CA-CB-OG1	6.97	123.64	109.00
1	C	292	ASP	CB-CG-OD1	6.96	124.56	118.30
1	D	77	ASP	OD1-CG-OD2	-6.85	110.29	123.30
1	D	73	ARG	NE-CZ-NH1	-6.52	117.04	120.30
1	B	77	ASP	CB-CG-OD1	6.50	124.15	118.30
1	A	213	LEU	CB-CG-CD1	6.49	122.04	111.00
1	B	252	LEU	CB-CG-CD1	6.44	121.94	111.00
1	A	122	VAL	CG1-CB-CG2	6.01	120.51	110.90
1	D	110	ASP	CB-CG-OD2	5.91	123.62	118.30
1	C	213	LEU	CB-CG-CD1	5.78	120.83	111.00
1	B	213	LEU	CB-CG-CD1	5.72	120.72	111.00
1	B	45	LYS	CD-CE-NZ	-5.71	98.58	111.70
1	A	252	LEU	CB-CG-CD1	5.65	120.61	111.00
1	D	121	LYS	CD-CE-NZ	-5.55	98.93	111.70
1	C	154	ASP	CB-CG-OD2	5.43	123.19	118.30
1	C	200	LEU	CB-CG-CD1	5.37	120.12	111.00
1	D	45	LYS	CD-CE-NZ	-5.36	99.37	111.70
1	D	200	LEU	CB-CG-CD1	5.34	120.09	111.00
1	A	213	LEU	CA-CB-CG	5.32	127.54	115.30
1	C	205	ASP	CB-CG-OD1	5.23	123.01	118.30
1	D	252	LEU	CB-CG-CD1	5.20	119.84	111.00
1	D	183	LEU	CB-CG-CD1	5.17	119.79	111.00
1	D	122	VAL	N-CA-CB	-5.16	100.16	111.50
1	D	45	LYS	CB-CG-CD	-5.13	98.27	111.60
1	B	183	LEU	CB-CG-CD1	5.12	119.71	111.00
1	C	213	LEU	CA-CB-CG	5.10	127.03	115.30
1	D	213	LEU	CB-CG-CD1	5.10	119.67	111.00
1	D	73	ARG	CG-CD-NE	-5.09	101.11	111.80
1	B	292	ASP	CB-CG-OD1	5.04	122.84	118.30
1	B	122	VAL	N-CA-CB	-5.02	100.45	111.50

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2263	0	2280	18	0
1	B	2265	0	2287	7	0
1	C	2269	0	2284	24	1
1	D	2268	0	2289	18	0
2	A	4	0	6	0	0
2	B	4	0	6	0	0
2	C	4	0	6	0	0
2	D	4	0	6	2	0
3	A	123	0	0	5	0
3	B	136	0	0	1	1
3	C	117	0	0	4	0
3	D	119	0	0	4	0
All	All	9576	0	9164	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:129:ARG:HD2	3:D:374:HOH:O	1.77	0.85
1:C:122:VAL:HG22	1:D:264:PRO:HD3	1.65	0.77
1:C:264:PRO:HD3	1:D:122:VAL:HG22	1.66	0.77
1:A:264:PRO:HD3	1:B:122:VAL:HG22	1.68	0.76
1:A:122:VAL:HG22	1:B:264:PRO:HD3	1.70	0.74
1:C:281:SER:OG	1:C:283[B]:HIS:HD2	1.72	0.72
1:A:209:ASN:ND2	3:A:489:HOH:O	2.23	0.70
1:B:188:HIS:CE1	3:B:362:HOH:O	2.45	0.70
1:D:20:PRO:HG3	1:D:174:KPI:H1A	1.75	0.69
1:B:105:GLN:HB2	1:B:140:LEU:HD22	1.78	0.66
1:C:281:SER:OG	1:C:283[B]:HIS:CD2	2.50	0.65
1:A:188:HIS:HB3	3:A:452:HOH:O	1.97	0.63
1:A:213:LEU:HD13	1:C:253:LEU:HB3	1.85	0.58
1:D:145:TYR:HD1	2:D:4:EDO:H21	1.69	0.58
1:D:188:HIS:CE1	3:D:445:HOH:O	2.57	0.57
1:A:188:HIS:CE1	3:A:475:HOH:O	2.56	0.56
1:D:105:GLN:HB2	1:D:140:LEU:HD22	1.86	0.56
1:A:28:ASP:OD1	1:A:28:ASP:N	2.37	0.55
1:D:174:KPI:CX1	3:D:444:HOH:O	2.55	0.55
1:A:105:GLN:HB2	1:A:140:LEU:HD22	1.89	0.54
1:A:174:KPI:CX1	3:A:354:HOH:O	2.56	0.54
1:D:12:ALA:N	3:D:442:HOH:O	2.41	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:174:KPI:O1	3:C:380:HOH:O	2.18	0.51
1:A:20:PRO:HG3	1:A:174:KPI:H1A	1.93	0.51
1:A:253:LEU:HB3	1:C:213:LEU:HD13	1.92	0.50
1:A:188:HIS:HE1	1:C:254:GLN:OE1	1.93	0.50
1:D:89:THR:HG21	1:D:101:SER:HA	1.94	0.49
1:D:145:TYR:CD1	2:D:4:EDO:H21	2.48	0.49
1:C:176:THR:HG21	3:C:381:HOH:O	2.14	0.48
1:C:94:ALA:HB3	1:D:286:PRO:HB2	1.96	0.47
1:A:232:ASN:HB3	1:A:248:TYR:CZ	2.49	0.47
1:A:233:LEU:C	1:A:233:LEU:HD23	2.35	0.47
1:D:143:MET:HE2	1:D:200:LEU:HD22	1.97	0.46
1:B:20:PRO:HG3	1:B:174:KPI:H1A	1.96	0.46
1:C:20:PRO:HG3	1:C:174:KPI:H1A	1.98	0.45
1:D:143:MET:CE	1:D:200:LEU:HD22	2.47	0.45
1:A:254:GLN:OE1	1:C:188:HIS:CE1	2.70	0.45
1:D:233:LEU:C	1:D:233:LEU:HD23	2.37	0.45
1:C:262:ASP:OD1	3:C:485:HOH:O	2.21	0.45
1:B:17:ILE:HD12	1:B:230:SER:HB3	1.99	0.45
1:C:105:GLN:HB2	1:C:140:LEU:HD22	2.00	0.44
1:C:229:VAL:HG11	1:C:252:LEU:HD13	2.00	0.44
1:A:176:THR:HG21	3:A:354:HOH:O	2.19	0.43
1:C:233:LEU:C	1:C:233:LEU:HD23	2.39	0.42
1:C:232:ASN:HB3	1:C:248:TYR:CZ	2.55	0.42
1:D:202:GLY:HA2	1:D:219:ILE:O	2.19	0.42
1:A:188:HIS:CE1	1:C:254:GLN:OE1	2.72	0.42
1:A:254:GLN:OE1	1:C:188:HIS:HE1	2.02	0.41
1:C:287:PRO:HG2	1:D:122:VAL:HG21	2.01	0.41
1:B:213:LEU:HD13	1:D:253:LEU:HB3	2.03	0.41
1:C:154:ASP:OD2	1:C:182:HIS:NE2	2.54	0.41
1:C:187:ILE:O	1:C:191:LYS:HB2	2.20	0.41
1:C:101:SER:HB3	1:C:112:ILE:HD13	2.04	0.40
1:C:174:KPI:CX1	3:C:381:HOH:O	2.69	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:73:ARG:NH2	3:B:430:HOH:O[4_454]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/298 (99%)	288 (98%)	7 (2%)	0	100	100
1	B	295/298 (99%)	290 (98%)	5 (2%)	0	100	100
1	C	296/298 (99%)	291 (98%)	5 (2%)	0	100	100
1	D	295/298 (99%)	289 (98%)	6 (2%)	0	100	100
All	All	1181/1192 (99%)	1158 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/242 (100%)	227 (94%)	14 (6%)	20	23
1	B	241/242 (100%)	226 (94%)	15 (6%)	18	21
1	C	240/242 (99%)	227 (95%)	13 (5%)	22	26
1	D	242/242 (100%)	227 (94%)	15 (6%)	18	21
All	All	964/968 (100%)	907 (94%)	57 (6%)	19	23

All (57) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LEU
1	A	28	ASP

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	43	LEU
1	A	45	LYS
1	A	122	VAL
1	A	140	LEU
1	A	145	TYR
1	A	147	PHE
1	A	183	LEU
1	A	200	LEU
1	A	213	LEU
1	A	252	LEU
1	A	275	LEU
1	A	308	LEU
1	B	13	LEU
1	B	28	ASP
1	B	43	LEU
1	B	45	LYS
1	B	122	VAL
1	B	140	LEU
1	B	145	TYR
1	B	147	PHE
1	B	183	LEU
1	B	188	HIS
1	B	200	LEU
1	B	213	LEU
1	B	252	LEU
1	B	275	LEU
1	B	308	LEU
1	C	13	LEU
1	C	43	LEU
1	C	122	VAL
1	C	140	LEU
1	C	145	TYR
1	C	147	PHE
1	C	183	LEU
1	C	200	LEU
1	C	213	LEU
1	C	252	LEU
1	C	275	LEU
1	C	302	LEU
1	C	308	LEU
1	D	13	LEU
1	D	28	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	43	LEU
1	D	45	LYS
1	D	122	VAL
1	D	134	VAL
1	D	140	LEU
1	D	145	TYR
1	D	147	PHE
1	D	183	LEU
1	D	200	LEU
1	D	213	LEU
1	D	252	LEU
1	D	275	LEU
1	D	308	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	102	GLN
1	A	106	GLN
1	A	188	HIS
1	A	254	GLN
1	A	257	GLN
1	B	62	GLN
1	B	102	GLN
1	B	254	GLN
1	B	257	GLN
1	C	62	GLN
1	C	102	GLN
1	C	188	HIS
1	D	62	GLN
1	D	102	GLN
1	D	188	HIS

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	KPI	A	174	1	11,13,14	1.09	0	10,15,17	2.57	3 (30%)
1	KPI	D	174	1	11,13,14	0.80	0	10,15,17	2.63	2 (20%)
1	KPI	C	174	1	11,13,14	1.06	1 (9%)	10,15,17	2.38	5 (50%)
1	KPI	B	174	1	11,13,14	1.48	2 (18%)	10,15,17	2.23	4 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KPI	A	174	1	-	5/13/14/16	-
1	KPI	D	174	1	-	3/13/14/16	-
1	KPI	C	174	1	-	4/13/14/16	-
1	KPI	B	174	1	-	3/13/14/16	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	174	KPI	O1-CX2	-2.33	1.23	1.30
1	C	174	KPI	O1-CX2	-2.28	1.23	1.30
1	B	174	KPI	CX2-CX1	2.11	1.52	1.49

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	174	KPI	C1-CX1-CX2	-7.33	111.04	118.17
1	A	174	KPI	C1-CX1-CX2	-6.84	111.52	118.17
1	B	174	KPI	C1-CX1-CX2	-4.23	114.05	118.17

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	174	KPI	C1-CX1-CX2	-4.02	114.25	118.17
1	C	174	KPI	O2-CX2-CX1	-3.96	116.32	121.38
1	B	174	KPI	O2-CX2-CX1	-3.64	116.73	121.38
1	C	174	KPI	C1-CX1-NZ	-3.02	115.21	123.11
1	A	174	KPI	C1-CX1-NZ	-2.80	115.80	123.11
1	D	174	KPI	CX2-CX1-NZ	-2.63	108.57	114.98
1	C	174	KPI	O1-CX2-CX1	2.33	121.42	116.35
1	B	174	KPI	CX2-CX1-NZ	-2.31	109.35	114.98
1	C	174	KPI	CG-CD-CE	-2.15	106.08	113.57
1	B	174	KPI	CG-CD-CE	-2.15	106.10	113.57
1	A	174	KPI	CX2-CX1-NZ	-2.12	109.83	114.98

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	174	KPI	C1-CX1-NZ-CE
1	A	174	KPI	NZ-CX1-CX2-O1
1	A	174	KPI	C1-CX1-CX2-O2
1	B	174	KPI	C1-CX1-NZ-CE
1	B	174	KPI	NZ-CX1-CX2-O1
1	B	174	KPI	NZ-CX1-CX2-O2
1	C	174	KPI	C1-CX1-NZ-CE
1	C	174	KPI	NZ-CX1-CX2-O2
1	D	174	KPI	C1-CX1-NZ-CE
1	D	174	KPI	NZ-CX1-CX2-O1
1	D	174	KPI	C1-CX1-CX2-O2
1	A	174	KPI	C1-CX1-CX2-O1
1	C	174	KPI	C1-CX1-CX2-O1
1	C	174	KPI	NZ-CX1-CX2-O1
1	A	174	KPI	NZ-CX1-CX2-O2

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	174	KPI	2	0
1	D	174	KPI	2	0
1	C	174	KPI	3	0
1	B	174	KPI	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EDO	A	1	-	3,3,3	0.37	0	2,2,2	0.72	0
2	EDO	D	4	-	3,3,3	0.48	0	2,2,2	0.58	0
2	EDO	C	3	-	3,3,3	0.61	0	2,2,2	0.67	0
2	EDO	B	2	-	3,3,3	0.33	0	2,2,2	1.92	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1	-	-	0/1/1/1	-
2	EDO	D	4	-	-	1/1/1/1	-
2	EDO	C	3	-	-	0/1/1/1	-
2	EDO	B	2	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	EDO	O2-C2-C1	-2.52	93.81	111.91

There are no chirality outliers.

All (1) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	D	4	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	4	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	297/298 (99%)	-0.60	0 100 100	7, 12, 23, 35	0
1	B	297/298 (99%)	-0.58	0 100 100	7, 12, 22, 34	0
1	C	297/298 (99%)	-0.71	1 (0%) 94 93	7, 12, 22, 33	0
1	D	297/298 (99%)	-0.72	0 100 100	7, 12, 23, 34	0
All	All	1188/1192 (99%)	-0.65	1 (0%) 95 95	7, 12, 23, 35	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	28	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KPI	D	174	14/15	0.95	0.09	5,9,19,23	0
1	KPI	B	174	14/15	0.96	0.12	6,9,19,22	0
1	KPI	C	174	14/15	0.97	0.12	6,9,19,23	0
1	KPI	A	174	14/15	0.97	0.10	5,9,19,22	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	C	3	4/4	0.95	0.15	17,17,18,24	0
2	EDO	B	2	4/4	0.96	0.15	13,14,17,20	0
2	EDO	A	1	4/4	0.96	0.12	16,20,20,26	0
2	EDO	D	4	4/4	0.96	0.14	9,18,22,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.