

wwPDB EM Validation Summary Report (i)

Oct 19, 2024 – 07:25 PM EDT

PDB ID : 6N7H

EMDB ID : EMD-0356

Title : Cryo-EM structure of the 2:1 hPtch1-Shhp complex

Authors: Yan, N.; Gong, X.; Qian, H.W.

Deposited on : 2018-11-27

Resolution : 3.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113

Mogul : 2022.3.0, CSD as543be (2022)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : 1.9.13

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM~structures} \ (\#{ m Entries})$	
Clashscore	210492	15764	
Ramachandran outliers	207382	16835	
Sidechain outliers	206894	16415	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
			11%				
1	A	1349	59%	13%	•	27%	
			11%				
1	В	1349	61%	12%	•	27%	
			9%				
2	С	174	72%			26%	•
			50%				
3	D	2	100%				
			50%				
3	E	2	50%		50%		

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CLR	A	1808	-	-	X	-
5	CLR	В	1508	-	-	X	-
8	PLM	С	205	-	-	X	-



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 17242 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein patched homolog 1.

Mol	Chain	Residues	${f Atoms}$				AltConf	Trace	
1	A	991	Total 7807	C 5091	N 1282	O 1392	S 42	0	0
1	В	988	Total 7708	C 5029	N 1267	O 1372	S 40	0	0

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP Q13635
A	-19	ALA	-	expression tag	UNP Q13635
A	-18	ASP	-	expression tag	UNP Q13635
A	-17	TYR	-	expression tag	UNP Q13635
A	-16	LYS	-	expression tag	UNP Q13635
A	-15	ASP	-	expression tag	UNP Q13635
A	-14	ASP	-	expression tag	UNP Q13635
A	-13	ASP	-	expression tag	UNP Q13635
A	-12	ASP	-	expression tag	UNP Q13635
A	-11	LYS	-	expression tag	UNP Q13635
A	-10	SER	-	expression tag	UNP Q13635
A	-9	GLY	-	expression tag	UNP Q13635
A	-8	PRO	-	expression tag	UNP Q13635
A	-7	ASP	-	expression tag	UNP Q13635
A	-6	GLU	-	expression tag	UNP Q13635
A	-5	VAL	-	expression tag	UNP Q13635
A	-4	ASP	-	expression tag	UNP Q13635
A	-3	ALA	-	expression tag	UNP Q13635
A	-2	SER	-	expression tag	UNP Q13635
A	-1	GLY	-	expression tag	UNP Q13635
A	0	ARG	-	expression tag	UNP Q13635
A	1306	LEU	-	expression tag	UNP Q13635
A	1307	GLU	-	expression tag	UNP Q13635
A	1308	GLY	-	expression tag	UNP Q13635
A	1309	SER	-	expression tag	UNP Q13635
A	1310	ASP	_	expression tag	UNP Q13635

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1311	GLU	-	expression tag	UNP Q13635
A	1312	VAL	-	expression tag	UNP Q13635
A	1313	ASP	-	expression tag	UNP Q13635
A	1314	ALA	-	expression tag	UNP Q13635
A	1315	VAL	-	expression tag	UNP Q13635
A	1316	GLU	-	expression tag	UNP Q13635
A	1317	GLY	-	expression tag	UNP Q13635
A	1318	SER	-	expression tag	UNP Q13635
A	1319	HIS	-	expression tag	UNP Q13635
A	1320	HIS	-	expression tag	UNP Q13635
A	1321	HIS	-	expression tag	UNP Q13635
A	1322	HIS	-	expression tag	UNP Q13635
A	1323	HIS	_	expression tag	UNP Q13635
A	1324	HIS	-	expression tag	UNP Q13635
A	1325	HIS	-	expression tag	UNP Q13635
A	1326	HIS	-	expression tag	UNP Q13635
A	1327	HIS	-	expression tag	UNP Q13635
A	1328	HIS	-	expression tag	UNP Q13635
В	-20	MET	-	initiating methionine	UNP Q13635
В	-19	ALA	-	expression tag	UNP Q13635
В	-18	ASP	-	expression tag	UNP Q13635
В	-17	TYR	-	expression tag	UNP Q13635
В	-16	LYS	-	expression tag	UNP Q13635
В	-15	ASP	-	expression tag	UNP Q13635
В	-14	ASP	-	expression tag	UNP Q13635
В	-13	ASP	-	expression tag	UNP Q13635
В	-12	ASP	-	expression tag	UNP Q13635
В	-11	LYS	-	expression tag	UNP Q13635
В	-10	SER	-	expression tag	UNP Q13635
В	-9	GLY	-	expression tag	UNP Q13635
В	-8	PRO	-	expression tag	UNP Q13635
В	-7	ASP	-	expression tag	UNP Q13635
В	-6	GLU	-	expression tag	UNP Q13635
В	-5	VAL	-	expression tag	UNP Q13635
В	-4	ASP	-	expression tag	UNP Q13635
В	-3	ALA	-	expression tag	UNP Q13635
В	-2	SER	-	expression tag	UNP Q13635
В	-1	GLY	-	expression tag	UNP Q13635
В	0	ARG	-	expression tag	UNP Q13635
В	1306	LEU	-	expression tag	UNP Q13635
В	1307	GLU	-	expression tag	UNP Q13635
В	1308	GLY	-	expression tag	UNP Q13635

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	1309	SER	-	expression tag	UNP Q13635
В	1310	ASP	-	expression tag	UNP Q13635
В	1311	GLU	-	expression tag	UNP Q13635
В	1312	VAL	-	expression tag	UNP Q13635
В	1313	ASP	-	expression tag	UNP Q13635
В	1314	ALA	-	expression tag	UNP Q13635
В	1315	VAL	-	expression tag	UNP Q13635
В	1316	GLU	-	expression tag	UNP Q13635
В	1317	GLY	-	expression tag	UNP Q13635
В	1318	SER	-	expression tag	UNP Q13635
В	1319	HIS	-	expression tag	UNP Q13635
В	1320	HIS	-	expression tag	UNP Q13635
В	1321	HIS	-	expression tag	UNP Q13635
В	1322	HIS	-	expression tag	UNP Q13635
В	1323	HIS	-	expression tag	UNP Q13635
В	1324	HIS	-	expression tag	UNP Q13635
В	1325	HIS	-	expression tag	UNP Q13635
В	1326	HIS	-	expression tag	UNP Q13635
В	1327	HIS	-	expression tag	UNP Q13635
В	1328	HIS	-	expression tag	UNP Q13635

• Molecule 2 is a protein called Sonic hedgehog protein.

Mol	Chain	Residues	Atoms			AltConf	Trace		
2	С	174	Total 1371	C 853	N 253	O 259	S 6	0	0


• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	AltConf	Trace
3	D	2	Total C N O 28 16 2 10	0	0
3	E	2	Total C N O 28 16 2 10	0	0

 \bullet Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\rm C_8H_{15}NO_6).$






Mol	Chain	Residues	A	Atoms				
4	A	1	Total	С	N	О	0	
4	Λ	1	14	8	1	5	U	
4	A	1	Total	С	N	Ο	0	
	11	1	14	8	1	5	O	
4	A	1	Total	С	N	O	0	
		1	14	8	1	5	Ü	
4	A	1	Total	С	N	О	0	
1	11	1	14	8	1	5	O	
4	A	1	Total	С	N	O	0	
1	11	1	14	8	1	5	Ů	
4	В	1	Total	С	N	О	0	
		1	14	8	1	5	Ŭ.	
4	В	1	Total	С	N	O	0	
		-	14	8	1	5	Ŭ	
4	В	1	Total	С	N	O	0	
		-	14	8	1	5	Ŭ	
4	В	1	Total	С	N	O	0	
_		1	14	8	1	5		
4	В	1	Total	С	N	О	0	
1		1	14	8	1	5		

 \bullet Molecule 5 is CHOLESTEROL (three-letter code: CLR) (formula: $\mathrm{C_{27}H_{46}O}).$





Mol	Chain	Residues	Atoms	AltConf
5	A	1	Total C O 28 27 1	0
5	A	1	Total C O 28 27 1	0
5	A	1	Total C O 28 27 1	0
5	В	1	Total C O 28 27 1	0
5	С	1	Total C O 28 27 1	0

• Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	AltConf
6	С	1	Total Zn 1 1	0

• Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	AltConf	
7	С	2	Total Ca 2 2	0	

 \bullet Molecule 8 is PALMITIC ACID (three-letter code: PLM) (formula: $\mathrm{C_{16}H_{32}O_2}).$






Mol	Chain	Residues	Atoms	AltConf
8	С	1	Total C O 17 16 1	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein patched homolog 1










• Molecule 2: Sonic hedgehog protein



 \bullet Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 100%



 \bullet Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 50% 50%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	171590	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.265	Depositor
Minimum map value	-0.136	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.045	Depositor
Map size (Å)	311.91998, 311.91998, 311.91998	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.114, 1.114, 1.114	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CLR, ZN, PLM, NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.45	0/8000	0.71	6/10883 (0.1%)	
1	В	0.48	0/7897	0.70	5/10749 (0.0%)	
2	С	0.56	1/1401 (0.1%)	0.76	3/1886 (0.2%)	
All	All	0.47	$1/17298 \; (0.0\%)$	0.71	14/23518 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	$\# ext{Chirality outliers}$	#Planarity outliers
1	A	0	7
1	В	0	6
2	С	0	1
All	All	0	14

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
2	С	98	MET	C-N	8.17	1.52	1.34

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	83	ASP	O-C-N	-7.32	110.99	122.70
1	В	297	LEU	CA-CB-CG	7.01	131.43	115.30
1	A	323	LEU	CA-CB-CG	6.32	129.84	115.30
1	В	262	LEU	CA-CB-CG	6.27	129.71	115.30
1	A	508	LEU	CA-CB-CG	6.21	129.59	115.30



There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	130	VAL	Peptide
1	A	224	TYR	Peptide
1	A	317	LEU	Peptide
1	A	835	GLU	Peptide
1	A	836	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7807	0	7828	108	0
1	В	7708	0	7664	121	0
2	С	1371	0	1329	33	0
3	D	28	0	25	0	0
3	Е	28	0	25	0	0
4	A	70	0	65	0	0
4	В	70	0	65	1	0
5	A	84	0	138	34	0
5	В	28	0	46	32	0
5	С	28	0	45	18	0
6	С	1	0	0	0	0
7	С	2	0	0	0	0
8	С	17	0	31	16	0
All	All	17242	0	17261	300	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 300 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:1147:PHE:HE2	8:C:205:PLM:CG	1.16	1.55
1:A:220:ILE:CD1	5:A:1808:CLR:C6	1.84	1.50
1:B:1147:PHE:CE2	8:C:205:PLM:CG	1.93	1.49

Continued on next page...



Continued from previous page...

Atom-1	Atom-1 Atom-2		$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$	
1:A:220:ILE:CD1	5:A:1808:CLR:H6	0.96	1.40	
1:B:216:MET:SD	5:C:201:CLR:H41	1.65	1.36	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	987/1349 (73%)	858 (87%)	122 (12%)	7 (1%)	19	53
1	В	984/1349 (73%)	856 (87%)	125 (13%)	3 (0%)	37	67
2	С	172/174 (99%)	153 (89%)	17 (10%)	2 (1%)	11	43
All	All	2143/2872 (75%)	1867 (87%)	264 (12%)	12 (1%)	24	55

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	939	ALA
2	С	188	GLU
1	A	424	THR
1	В	389	GLU
1	A	834	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	832/1147 (72%)	820 (99%)	12 (1%)	62	79	
1	В	808/1147 (70%)	795 (98%)	13 (2%)	58	76	
2	С	142/144~(99%)	138 (97%)	4 (3%)	38	64	
All	All	1782/2438 (73%)	1753 (98%)	29 (2%)	58	76	

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	220	ILE
2	С	129	ASP
1	В	349	ASN
1	В	845	LEU
1	В	310	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	409	GLN
1	В	853	GLN
2	С	107	ASN
1	В	846	HIS
1	В	929	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	1	3,1	14,14,15	0.25	0	17,19,21	0.58	0
3	NAG	D	2	3	14,14,15	0.27	0	17,19,21	0.55	0
3	NAG	Е	1	3,1	14,14,15	0.45	0	17,19,21	0.67	0
3	NAG	Е	2	3	14,14,15	0.27	0	17,19,21	0.71	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	4/6/23/26	0/1/1/1
3	NAG	Е	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	Ε	2	NAG	C1-O5-C5	2.54	115.60	112.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2	NAG	C4-C5-C6-O6
3	Е	2	NAG	C4-C5-C6-O6
3	Е	2	NAG	O5-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
3	D	2	NAG	C8-C7-N2-C2

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 3 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain Res			Link	Вс	Bond lengths			Bond angles		
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	NAG	A	1801	1	14,14,15	0.88	1 (7%)	17,19,21	2.39	4 (23%)	
4	NAG	В	1505	1	14,14,15	0.25	0	17,19,21	0.50	0	
8	PLM	С	205	2	15,16,17	0.32	0	14,15,17	0.66	0	



Mol	Type	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1807	1	14,14,15	0.40	0	17,19,21	0.42	0
4	NAG	В	1501	1	14,14,15	0.70	1 (7%)	17,19,21	0.70	0
4	NAG	A	1804	1	14,14,15	0.23	0	17,19,21	0.59	0
5	CLR	A	1808	-	31,31,31	1.04	2 (6%)	48,48,48	1.67	8 (16%)
5	CLR	A	1809	-	31,31,31	0.73	0	48,48,48	1.55	9 (18%)
5	CLR	A	1810	-	31,31,31	0.68	0	48,48,48	1.28	6 (12%)
4	NAG	A	1803	1	14,14,15	0.79	1 (7%)	17,19,21	1.28	2 (11%)
4	NAG	В	1503	1	14,14,15	0.94	1 (7%)	17,19,21	1.40	3 (17%)
5	CLR	С	201	2	31,31,31	1.02	1 (3%)	48,48,48	1.98	9 (18%)
4	NAG	В	1504	1	14,14,15	0.75	1 (7%)	17,19,21	0.77	1 (5%)
4	NAG	A	1802	1	14,14,15	0.24	0	17,19,21	0.57	0
4	NAG	В	1502	1	14,14,15	0.54	0	17,19,21	0.39	0
5	CLR	В	1508	_	31,31,31	0.90	2 (6%)	48,48,48	1.58	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1801	1	-	6/6/23/26	0/1/1/1
4	NAG	В	1505	1	-	4/6/23/26	0/1/1/1
8	PLM	С	205	2	-	7/14/14/15	-
4	NAG	A	1807	1	-	3/6/23/26	0/1/1/1
4	NAG	В	1501	1	-	1/6/23/26	0/1/1/1
4	NAG	A	1804	1	-	1/6/23/26	0/1/1/1
5	CLR	A	1808	-	-	9/10/68/68	0/4/4/4
5	CLR	A	1809	-	-	2/10/68/68	0/4/4/4
5	CLR	A	1810	-	-	6/10/68/68	0/4/4/4
4	NAG	A	1803	1	-	4/6/23/26	0/1/1/1
4	NAG	В	1503	1	-	2/6/23/26	0/1/1/1
5	CLR	С	201	2	-	3/10/68/68	0/4/4/4
4	NAG	В	1504	1	-	4/6/23/26	0/1/1/1
4	NAG	A	1802	1	-	1/6/23/26	0/1/1/1
4	NAG	В	1502	1	-	0/6/23/26	0/1/1/1
5	CLR	В	1508	-	-	4/10/68/68	0/4/4/4

The worst 5 of 10 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
5	С	201	CLR	C13-C14	-2.78	1.49	1.55
4	A	1801	NAG	C1-C2	2.66	1.56	1.52
5	A	1808	CLR	C13-C14	-2.61	1.50	1.55
4	A	1803	NAG	O5-C1	-2.61	1.39	1.43
4	В	1503	NAG	O5-C1	-2.59	1.39	1.43

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
4	A	1801	NAG	C2-N2-C7	8.38	134.14	122.90
5	С	201	CLR	C13-C17-C20	-5.64	110.79	119.50
5	С	201	CLR	C13-C14-C8	-5.60	106.46	114.41
5	С	201	CLR	C8-C7-C6	-4.92	105.94	112.76
5	В	1508	CLR	C13-C17-C20	-4.77	112.12	119.50

There are no chirality outliers.

5 of 57 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	С	201	CLR	C17-C20-C22-C23
5	A	1810	CLR	C13-C17-C20-C21
5	С	201	CLR	C22-C23-C24-C25
5	A	1810	CLR	C16-C17-C20-C21
5	A	1810	CLR	C13-C17-C20-C22

There are no ring outliers.


7 monomers are involved in 101 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	С	205	PLM	16	0
4	В	1501	NAG	1	0
5	A	1808	CLR	31	0
5	A	1809	CLR	1	0
5	A	1810	CLR	2	0
5	С	201	CLR	18	0
5	В	1508	CLR	32	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.




Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Rings

Torsions














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-0356. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map





The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map








Y Index: 140 Z Index: 140



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.045. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $100~\mathrm{nm}^3$; this corresponds to an approximate mass of $91~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



^{*}Reported resolution corresponds to spatial frequency of 0.278 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-0356 and PDB model 6N7H. Per-residue inclusion information can be found in section 3 on page 10.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.045 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.045).



9.4 Atom inclusion (i)



At the recommended contour level, 78% of all backbone atoms, 65% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.045) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6470	0.4080
A	0.6300	0.4040
В	0.6490	0.4070
С	0.7400	0.4400
D	0.3570	0.3380
E	0.4290	0.4030



