



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 10:25 pm BST

PDB ID : 1N8E  
Title : Fragment Double-D from Human Fibrin  
Authors : Yang, Z.; Pandi, L.; Doolittle, R.F.  
Deposited on : 2002-11-20  
Resolution : 4.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibrin alpha/alpha-E chain.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	74	Total C 74 74	0	0	74
1	D	74	Total C 74 74	0	0	74

- Molecule 2 is a protein called Fibrin beta chain.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	B	307	Total C 307 307	0	0	307
2	E	307	Total C 307 307	0	0	307

- Molecule 3 is a protein called Fibrin gamma chain.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	C	295	Total C 295 295	0	0	295
3	F	295	Total C 295 295	0	0	295

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: Fibrin alpha/alpha-E chain

Chain A:  80% 17%




- Molecule 1: Fibrin alpha/alpha-E chain

Chain D:  83% 17%




- Molecule 2: Fibrin beta chain

Chain B:  94% 6%




- Molecule 2: Fibrin beta chain

Chain E:  93% 6%




- Molecule 3: Fibrin gamma chain

Chain C:  91% 9%



- Molecule 3: Fibrin gamma chain

Chain F:  91% 9%

LYS	MET	LEU	GLU	GLU	ILE	MET	LYS	TYR	E97	R991	LEU	THR	ILE	GLY	GLU	GLY	GLN	GLN	HIS	HIS	LEU	LEU	GLY	GLY	ALA	ALA	LYS	GLN	GLN	ALA	ALA	GLY	GLY	ASP	VAL
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.00Å 125.35Å 312.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 4.50	Depositor
% Data completeness (in resolution range)	(Not available) (30.00-4.50)	Depositor
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.415 , 0.416	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1352	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	74	0	0	1	1
1	D	74	0	0	0	0
2	B	307	0	0	0	0
2	E	307	0	0	1	1
3	C	295	0	0	0	0
3	F	295	0	0	0	0
All	All	1352	0	0	2	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (2) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:191:GLU:CA	2:E:192:TYR:CA	2.89	0.50
1:A:166:SER:CA	1:A:167:ARG:CA	2.99	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:ALA:CA	2:E:348:THR:CA[2_655]	1.71	0.49

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

#### 5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

#### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.