



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2023 – 01:07 AM EDT

PDB ID : 6N9J
Title : X-ray structure of a secreted C11 cysteine protease from *Bacteroides thetaio-*
taomicron in complex with an irreversible peptide inhibitor
Authors : Wolan, D.W.; Gonzalez-Paez, G.E.; Roncase, E.J.
Deposited on : 2018-12-03
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	85% 8% 7%
1	B	377	84% 10% 6% 1%
2	C	5	100%
2	D	5	80% 20%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Clostripain-related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	352	2753	1741	456	542	14	0	0	0
1	B	356	2764	1747	458	545	14	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLY	-	expression tag	UNP Q8A866
B	23	GLY	-	expression tag	UNP Q8A866

- Molecule 2 is a protein called Covalently bound peptide inhibitor.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	5	35	24	5	6	0	0	0
2	D	5	35	24	5	6	0	0	0


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	175	Total	O	0	0
			175	175		
3	C	4	Total	O	0	0
			4	4		
3	B	165	Total	O	0	0
			165	165		
3	D	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Clostripain-related protein

Chain A:  85% 8% 7%



- Molecule 1: Clostripain-related protein

Chain B:  84% 10% 6%




- Molecule 2: Covalently bound peptide inhibitor

Chain C:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: Covalently bound peptide inhibitor

Chain D:  80% 20%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.96Å 98.96Å 68.33Å 90.00° 89.90° 90.00°	Depositor
Resolution (Å)	49.48 – 2.17 49.48 – 2.17	Depositor EDS
% Data completeness (in resolution range)	93.3 (49.48-2.17) 92.9 (49.48-2.17)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.05 (at 2.18Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.181 , 0.231 0.184 , 0.230	Depositor DCC
R_{free} test set	2097 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtrriage
Anisotropy	0.850	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 23.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.316 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5934	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CKC, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2823	0.48	0/3849
1	B	0.36	0/2833	0.50	0/3863
2	C	0.41	0/22	0.69	0/30
2	D	1.13	0/22	1.47	0/30
All	All	0.36	0/5700	0.50	0/7772

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	403	LEU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2753	0	2538	19	0
1	B	2764	0	2544	21	0
2	C	35	0	38	0	0
2	D	35	0	38	0	0
3	A	175	0	0	3	0
3	B	165	0	0	4	0
3	C	4	0	0	0	0
3	D	3	0	0	0	0
All	All	5934	0	5158	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:320:GLN:HG3	1:A:326:PHE:HB2	1.54	0.87
1:B:346:ASN:OD1	1:B:360:MET:HG3	1.90	0.69
1:B:166:ASN:O	1:B:197:ARG:NH2	2.26	0.68
1:B:70:LEU:HD23	1:B:85:LEU:HD12	1.76	0.66
1:B:387:GLU:OE1	3:B:401:HOH:O	2.14	0.65
1:B:191:VAL:HG12	1:B:270:LEU:HD11	1.79	0.64
1:B:380:LYS:NZ	3:B:407:HOH:O	2.29	0.64
1:B:60:MET:HE3	1:B:70:LEU:HD22	1.81	0.63
1:B:31:GLY:N	3:B:408:HOH:O	2.33	0.62
1:A:84:ARG:HH12	1:A:86:ARG:HG3	1.65	0.61
1:A:383:ARG:NH1	1:A:389:TYR:OH	2.37	0.58
1:A:57:ILE:HG23	1:A:92:VAL:HB	1.86	0.57
1:B:383:ARG:HG3	1:B:389:TYR:CZ	2.41	0.55
1:B:191:VAL:HG13	1:B:201:PHE:CD1	2.44	0.52
1:A:70:LEU:HD23	1:A:85:LEU:HD12	1.91	0.51
1:B:71:LEU:HD23	1:B:118:ALA:HB1	1.93	0.51
1:B:266:LYS:NZ	1:B:268:SER:OG	2.43	0.50
1:A:278:ARG:O	1:A:282:GLN:HG2	2.11	0.50
1:B:320:GLN:HG3	1:B:326:PHE:HB2	1.94	0.49
1:A:191:VAL:HG12	1:A:270:LEU:HD11	1.94	0.49
1:B:316:MET:HA	1:B:316:MET:HE2	1.95	0.49
1:A:278:ARG:NH2	3:A:417:HOH:O	2.46	0.49
1:A:320:GLN:HG2	1:A:322:ASN:H	1.80	0.46
1:B:68:ASN:HB3	1:B:223:PHE:CD1	2.51	0.46
1:B:106:VAL:HB	1:B:160:MET:HB2	1.98	0.45
1:A:88:ASP:OD1	3:A:401:HOH:O	2.21	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:ARG:NH1	3:A:407:HOH:O	2.36	0.44
1:A:310:HIS:HE1	1:A:372:ARG:HH22	1.65	0.43
1:A:340:LYS:HE2	1:A:364:GLU:OE2	2.17	0.43
1:A:363:PHE:CZ	1:A:365:GLY:HA2	2.54	0.43
1:A:131:PHE:HB3	1:A:149:TRP:CZ2	2.52	0.43
1:A:274:ALA:O	1:A:278:ARG:HG3	2.18	0.43
1:B:260:VAL:C	1:B:346:ASN:HD21	2.21	0.42
1:A:39:ILE:HG23	1:A:132:TRP:CE3	2.55	0.42
1:B:383:ARG:NE	3:B:415:HOH:O	2.45	0.42
1:B:310:HIS:HE1	1:B:372:ARG:HH22	1.68	0.42
1:B:129:VAL:CG1	1:B:178:ILE:HG12	2.51	0.41
1:A:363:PHE:CE1	1:A:365:GLY:HA2	2.56	0.41
1:A:191:VAL:HG11	1:A:265:ILE:HG21	2.04	0.40
1:B:88:ASP:O	1:B:91:VAL:HG12	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/377 (92%)	333 (96%)	12 (4%)	1 (0%)	41	43
1	B	350/377 (93%)	339 (97%)	10 (3%)	1 (0%)	41	43
2	C	3/5 (60%)	3 (100%)	0	0	100	100
2	D	3/5 (60%)	3 (100%)	0	0	100	100
All	All	702/764 (92%)	678 (97%)	22 (3%)	2 (0%)	41	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	78	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	78	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/327 (90%)	288 (98%)	5 (2%)	60	72
1	B	293/327 (90%)	289 (99%)	4 (1%)	67	78
2	C	3/3 (100%)	3 (100%)	0	100	100
2	D	3/3 (100%)	3 (100%)	0	100	100
All	All	592/660 (90%)	583 (98%)	9 (2%)	65	76

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	187	SER
1	A	225	VAL
1	A	236	TYR
1	A	246	SER
1	A	320	GLN
1	B	50	SER
1	B	187	SER
1	B	236	TYR
1	B	283	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/377 (93%)	-0.20	0 100 100	12, 24, 47, 71	0
1	B	356/377 (94%)	-0.20	2 (0%) 89 89	12, 24, 44, 64	0
2	C	3/5 (60%)	-0.08	0 100 100	25, 25, 26, 31	0
2	D	3/5 (60%)	-0.39	0 100 100	23, 23, 26, 32	0
All	All	714/764 (93%)	-0.20	2 (0%) 94 94	12, 24, 45, 71	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	146	THR	2.4
1	B	283	THR	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.