



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 11:01 PM EDT

PDB ID : 7N9H
Title : Structure of the mammalian importin $\alpha 1$ bound to the TDP-43 NLS
Authors : Doll, S.G.; Lokareddy, R.K.; Cingolani, G.
Deposited on : 2021-06-17
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	428	 4% 91% 7%
2	A	24	 63% 75% 21%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3554 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	422	3220	2052	546	612	10	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	79	GLU	ASP	conflict	UNP P52292
C	88	ASN	SER	conflict	UNP P52292
C	90	LEU	VAL	conflict	UNP P52292
C	92	SER	ASN	conflict	UNP P52292
C	130	LYS	ARG	conflict	UNP P52292
C	195	ALA	VAL	conflict	UNP P52292
C	203	HIS	TYR	conflict	UNP P52292
C	206	ILE	VAL	conflict	UNP P52292
C	218	LEU	MET	conflict	UNP P52292
C	220	THR	SER	conflict	UNP P52292
C	246	LEU	ILE	conflict	UNP P52292
C	263	ASN	ASP	conflict	UNP P52292
C	271	SER	THR	conflict	UNP P52292
C	287	GLU	GLY	conflict	UNP P52292
C	292	LYS	THR	conflict	UNP P52292
C	305	THR	SER	conflict	UNP P52292
C	330	LYS	VAL	conflict	UNP P52292
C	384	GLY	SER	conflict	UNP P52292
C	398	ALA	VAL	conflict	UNP P52292
C	401	ILE	VAL	conflict	UNP P52292
C	430	SER	THR	conflict	UNP P52292
C	438	GLN	LEU	conflict	UNP P52292
C	478	ARG	ASN	conflict	UNP P52292
C	490	ASN	SER	conflict	UNP P52292

- Molecule 2 is a protein called TAR DNA-binding protein 43.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	23	171	101	35	34	1	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	162	Total 162	O 162	0	0
3	A	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Importin subunit alpha-1



- Molecule 2: TAR DNA-binding protein 43



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.37Å 91.31Å 97.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.97 – 2.20 14.97 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (14.97-2.20) 99.7 (14.97-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.19.2	Depositor
R, R_{free}	0.190 , 0.215 0.189 , 0.215	Depositor DCC
R_{free} test set	1994 reflections (5.53%)	wwPDB-VP
Wilson B-factor (Å ²)	37.8	Xtrriage
Anisotropy	0.385	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3554	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.39	0/3278	0.55	0/4467
2	A	0.40	0/170	0.70	0/223
All	All	0.39	0/3448	0.56	0/4690

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3220	0	3296	20	0
2	A	171	0	167	4	0
3	A	1	0	0	1	0
3	C	162	0	0	1	0
All	All	3554	0	3463	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:458:GLU:HG2	1:C:459:LYS:HE3	1.81	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:288:MET:HA	1:C:291:LYS:HE3	1.84	0.58
1:C:425:LEU:HD22	1:C:440:ILE:HG23	1.90	0.54
1:C:474:GLU:OE2	1:C:478:ARG:NH2	2.41	0.53
1:C:272:CYS:HB3	1:C:312:PRO:HB2	1.94	0.50
1:C:484:VAL:O	1:C:488:SER:OG	2.29	0.49
1:C:287:GLU:O	1:C:291:LYS:HG2	2.12	0.49
1:C:471:ASP:OD2	1:C:471:ASP:N	2.43	0.48
2:A:94:VAL:O	2:A:95:LYS:HG2	2.14	0.46
1:C:465:GLU:HG3	1:C:470:LEU:HD23	1.98	0.46
1:C:231:TRP:CE2	2:A:98:ARG:HD2	2.51	0.45
1:C:458:GLU:HG2	1:C:459:LYS:CE	2.46	0.45
1:C:429:LEU:HD23	1:C:429:LEU:HA	1.82	0.45
1:C:95:GLN:NE2	3:C:509:HOH:O	2.51	0.44
1:C:494:LYS:HG2	1:C:495:TYR:CZ	2.52	0.44
1:C:142:TRP:HE1	2:A:100:VAL:HB	1.83	0.43
1:C:273:TRP:CD2	1:C:312:PRO:HB3	2.52	0.43
1:C:474:GLU:O	1:C:477:GLN:HG2	2.18	0.43
1:C:238:ARG:HA	1:C:277:TYR:CD2	2.54	0.43
1:C:434:THR:O	1:C:438:GLN:HG3	2.21	0.41
1:C:287:GLU:HA	1:C:290:VAL:HG22	2.02	0.41
2:A:95:LYS:NZ	3:A:201:HOH:O	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	420/428 (98%)	414 (99%)	6 (1%)	0	100 100
2	A	21/24 (88%)	15 (71%)	5 (24%)	1 (5%)	2 1
All	All	441/452 (98%)	429 (97%)	11 (2%)	1 (0%)	47 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	93	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	354/359 (99%)	350 (99%)	4 (1%)	73	85
2	A	17/21 (81%)	17 (100%)	0	100	100
All	All	371/380 (98%)	367 (99%)	4 (1%)	73	85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	432	LYS
1	C	442	ASP
1	C	485	TYR
1	C	488	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	241	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	422/428 (98%)	-0.05	18 (4%) 35 33	27, 40, 87, 121	0
2	A	23/24 (95%)	3.30	15 (65%) 0 0	52, 92, 140, 142	0
All	All	445/452 (98%)	0.12	33 (7%) 14 13	27, 40, 98, 142	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	92	SER	8.4
2	A	88	THR	8.0
1	C	485	TYR	8.0
2	A	91	SER	7.1
2	A	89	ASP	6.3
2	A	90	ALA	6.2
2	A	94	VAL	4.9
2	A	86	ASP	4.7
1	C	482	GLU	4.6
2	A	102	LYS	4.4
2	A	93	ALA	4.3
1	C	486	LYS	4.2
1	C	108	LYS	4.1
2	A	96	VAL	4.1
1	C	75	ASN	3.9
2	A	101	GLN	3.9
1	C	471	ASP	3.7
1	C	493	GLU	3.5
1	C	107	GLU	3.4
1	C	478	ARG	3.2
1	C	483	SER	3.1
1	C	432	LYS	3.1
2	A	87	GLU	3.0
1	C	88	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	100	VAL	2.9
2	A	80	ASP	2.8
1	C	109	GLN	2.8
1	C	480	GLU	2.7
1	C	468	GLY	2.4
1	C	496	PHE	2.3
2	A	95	LYS	2.2
1	C	476	LEU	2.2
1	C	474	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.