



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2024 – 10:33 AM EDT

PDB ID : 1NE2
Title : Crystal Structure of Thermoplasma acidophilum 1320 (APC5513)
Authors : Kim, Y.; Joachimiak, A.; Edwards, A.; Xu, X.; Christendat, D.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2002-12-10
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	 14% 67% 19% 12%
1	B	200	 18% 71% 20% 8%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein ta1320.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	176	1386	879	239	262	2	4	0	0	0
1	B	184	1451	917	255	273	2	4	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	cloning artifact	UNP Q9HIL9
A	-1	SER	-	cloning artifact	UNP Q9HIL9
A	0	HIS	-	cloning artifact	UNP Q9HIL9
A	1	MSE	MET	modified residue	UNP Q9HIL9
A	97	MSE	MET	modified residue	UNP Q9HIL9
A	113	MSE	MET	modified residue	UNP Q9HIL9
A	137	MSE	MET	modified residue	UNP Q9HIL9
B	-2	GLY	-	cloning artifact	UNP Q9HIL9
B	-1	SER	-	cloning artifact	UNP Q9HIL9
B	0	HIS	-	cloning artifact	UNP Q9HIL9
B	1	MSE	MET	modified residue	UNP Q9HIL9
B	97	MSE	MET	modified residue	UNP Q9HIL9
B	113	MSE	MET	modified residue	UNP Q9HIL9
B	137	MSE	MET	modified residue	UNP Q9HIL9

- Molecule 2 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 3 1 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	124	Total O 124 124	0	0
3	B	126	Total O 126 126	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: hypothetical protein ta1320



- Molecule 1: hypothetical protein ta1320



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.09Å 53.36Å 149.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.47 – 1.75 40.47 – 1.75	Depositor EDS
% Data completeness (in resolution range)	92.5 (40.47-1.75) 92.5 (40.47-1.75)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 1.74Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.209 , 0.236 0.199 , 0.224	Depositor DCC
R_{free} test set	3710 reflections (10.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtrriage
Anisotropy	0.161	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3090	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/1408	0.57	0/1890
1	B	0.33	0/1477	0.57	0/1985
All	All	0.33	0/2885	0.57	0/3875

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1386	0	1349	26	0
1	B	1451	0	1412	41	0
2	A	3	0	1	1	0
3	A	124	0	0	0	0
3	B	126	0	0	4	0
All	All	3090	0	2762	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:MSE:HE2	1:B:115:PRO:HG3	1.39	1.01
1:A:1:MSE:HG2	1:A:2:GLY:H	1.37	0.88
1:B:137:MSE:HE3	1:B:138:TRP:HE1	1.40	0.87
1:B:104:ILE:HB	1:B:131:LYS:HD3	1.73	0.71
1:B:144:ASN:HD22	1:B:145:ALA:N	1.90	0.70
1:A:1:MSE:HG2	1:A:2:GLY:N	2.07	0.68
1:B:167:VAL:HG23	1:B:187:ALA:HB3	1.77	0.67
1:A:35:ALA:O	1:A:39:ILE:HG12	1.98	0.64
1:B:113:MSE:CE	1:B:115:PRO:HG3	2.22	0.64
1:A:27:TYR:OH	1:A:175:TYR:HB2	1.96	0.64
1:A:152:ARG:HD3	1:A:164:GLU:OE1	1.97	0.63
1:B:46:ASN:CG	3:B:298:HOH:O	2.37	0.62
1:B:113:MSE:HE2	1:B:115:PRO:CG	2.23	0.62
1:A:81:PRO:O	1:A:85:GLU:HG3	2.01	0.60
1:B:171:VAL:HG21	1:B:185:ILE:HG23	1.85	0.58
1:A:6:ASP:O	1:A:9:ILE:HG12	2.04	0.58
1:A:72:GLU:O	1:B:97:MSE:HE1	2.03	0.57
1:B:46:ASN:N	1:B:46:ASN:OD1	2.37	0.57
1:A:41:ILE:HG22	1:A:47:ILE:HG12	1.87	0.55
1:A:126:ARG:HD3	1:A:154:GLU:OE1	2.06	0.55
1:A:153:ARG:HD3	1:A:153:ARG:C	2.27	0.55
1:B:115:PRO:HB3	1:B:117:PHE:CE2	2.42	0.55
1:B:137:MSE:SE	1:B:193:ARG:NH2	2.90	0.54
1:B:137:MSE:HE3	1:B:138:TRP:NE1	2.18	0.54
1:B:75:THR:HG23	1:B:97:MSE:HE3	1.89	0.53
1:B:144:ASN:HD22	1:B:145:ALA:H	1.55	0.53
1:A:17:GLN:HG2	1:A:86:THR:OG1	2.08	0.53
1:B:137:MSE:CE	1:B:138:TRP:HE1	2.17	0.53
1:B:166:LYS:HG3	1:B:186:GLU:HG3	1.91	0.51
1:A:97:MSE:HE1	1:B:72:GLU:O	2.10	0.50
1:B:193:ARG:NH1	1:B:194:ASN:O	2.46	0.49
1:A:115:PRO:HB2	1:A:117:PHE:CD2	2.47	0.49
1:B:164:GLU:HG3	1:B:190:PHE:CE1	2.48	0.49
1:A:17:GLN:HB3	1:A:86:THR:HG21	1.95	0.48
1:A:11:LEU:O	1:A:14:LEU:HB2	2.13	0.47
1:B:46:ASN:ND2	3:B:298:HOH:O	2.47	0.47
1:B:57:THR:OG1	1:B:60:GLY:HA2	2.15	0.47
1:B:144:ASN:ND2	1:B:146:LYS:H	2.12	0.46
1:B:176:ARG:CZ	1:B:183:ALA:HA	2.46	0.46
1:B:145:ALA:O	1:B:148:ARG:HB2	2.15	0.46
1:B:121:VAL:HG13	1:B:125:ASP:HB2	1.98	0.46
1:B:178:HIS:N	1:B:181:ASP:OD2	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:MSE:HE2	3:B:219:HOH:O	2.15	0.45
1:B:138:TRP:CH2	1:B:193:ARG:HD3	2.52	0.45
1:B:166:LYS:HG3	1:B:186:GLU:CG	2.47	0.45
1:A:3:ILE:HG23	1:A:4:LYS:N	2.33	0.44
1:A:117:PHE:N	1:A:117:PHE:CD1	2.85	0.44
1:B:171:VAL:CG2	1:B:185:ILE:HG23	2.47	0.44
1:B:124:SER:O	1:B:127:ALA:HB3	2.18	0.44
1:A:61:ILE:HG23	1:A:62:LEU:N	2.33	0.43
1:A:94:VAL:HB	1:A:96:PHE:CE2	2.53	0.43
1:B:1:MSE:N	1:B:1:MSE:SE	3.02	0.42
1:B:131:LYS:HE3	1:B:131:LYS:HA	2.02	0.42
1:B:144:ASN:OD1	1:B:146:LYS:HE2	2.19	0.42
1:A:57:THR:OG1	1:A:60:GLY:HA2	2.20	0.42
1:B:170:THR:HA	1:B:184:ARG:HG2	2.02	0.42
1:B:144:ASN:ND2	1:B:145:ALA:N	2.63	0.41
1:B:115:PRO:HB2	3:B:259:HOH:O	2.19	0.41
1:A:129:ILE:HD12	1:A:155:PHE:CE1	2.55	0.41
1:B:87:ALA:HB1	1:B:96:PHE:CE1	2.56	0.41
1:A:28:PRO:HD2	2:A:301:FMT:H	2.02	0.41
1:A:113:MSE:HE3	1:A:115:PRO:HB3	2.03	0.41
1:A:53:ILE:HA	1:A:75:THR:O	2.21	0.41
1:A:30:ASP:OD2	1:A:169:ILE:HG23	2.21	0.41
1:B:138:TRP:CZ3	1:B:193:ARG:HD3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/200 (84%)	160 (95%)	7 (4%)	1 (1%)	22	9
1	B	180/200 (90%)	172 (96%)	7 (4%)	1 (1%)	22	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	348/400 (87%)	332 (95%)	14 (4%)	2 (1%)	22 9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	196	SER
1	A	17	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	143/161 (89%)	139 (97%)	4 (3%)	38 18
1	B	151/161 (94%)	147 (97%)	4 (3%)	41 21
All	All	294/322 (91%)	286 (97%)	8 (3%)	40 19

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LYS
1	A	117	PHE
1	A	137	MSE
1	A	153	ARG
1	B	46	ASN
1	B	144	ASN
1	B	178	HIS
1	B	197	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	114	ASN
1	B	15	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	123	HIS
1	B	144	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FMT	A	301	-	2,2,2	0.64	0	1,1,1	0.38	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	FMT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/200 (86%)	0.82	29 (16%) 5 5	15, 25, 54, 79	0
1	B	180/200 (90%)	0.83	36 (20%) 3 4	14, 24, 69, 85	0
All	All	352/400 (88%)	0.82	65 (18%) 4 4	14, 25, 61, 85	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	116	PRO	7.6
1	B	197	PHE	6.7
1	A	18	GLY	6.4
1	B	120	VAL	5.9
1	A	116	PRO	5.5
1	B	117	PHE	5.5
1	B	121	VAL	5.0
1	A	197	PHE	5.0
1	B	180	TYR	4.8
1	A	16	GLN	4.6
1	A	175	TYR	4.6
1	B	185	ILE	4.5
1	A	117	PHE	4.4
1	B	118	GLY	4.3
1	B	179	SER	4.3
1	B	14	LEU	3.9
1	A	174	ILE	3.9
1	B	125	ASP	3.9
1	B	115	PRO	3.8
1	B	15	GLN	3.8
1	A	3	ILE	3.8
1	B	124	SER	3.7
1	B	13	LYS	3.6
1	A	182	ARG	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	182	ARG	3.4
1	B	181	ASP	3.4
1	B	122	LYS	3.4
1	A	17	GLN	3.3
1	A	125	ASP	3.2
1	A	9	ILE	3.2
1	B	178	HIS	3.1
1	B	123	HIS	3.1
1	A	157	ALA	3.1
1	B	183	ALA	3.1
1	B	177	HIS	3.0
1	A	185	ILE	2.9
1	A	27	TYR	2.9
1	B	5	ASN	2.9
1	A	28	PRO	2.8
1	A	126	ARG	2.8
1	B	126	ARG	2.7
1	B	119	SER	2.6
1	A	152	ARG	2.6
1	B	193	ARG	2.6
1	B	148	ARG	2.6
1	A	169	ILE	2.6
1	B	114	ASN	2.6
1	A	153	ARG	2.6
1	B	184	ARG	2.6
1	B	196	SER	2.6
1	A	85	GLU	2.5
1	B	46	ASN	2.4
1	A	149	ASP	2.4
1	B	167	VAL	2.4
1	B	2	GLY	2.3
1	A	146	LYS	2.3
1	A	15	GLN	2.3
1	A	2	GLY	2.3
1	B	29	THR	2.3
1	B	158	ARG	2.2
1	A	115	PRO	2.2
1	B	91	CYS	2.1
1	A	170	THR	2.0
1	A	89	ARG	2.0
1	A	6	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FMT	A	301	3/3	0.78	0.18	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.