



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 27, 2024 – 10:35 PM EDT

PDB ID : 1NEZ
Title : The Crystal Structure of a TL/CD8aa Complex at 2.1A resolution:Implications for Memory T cell Generation, Co-receptor Preference and Affinity
Authors : Liu, Y.; Xiong, Y.; Naidenko, O.V.; Liu, J.H.; Zhang, R.; Joachimiak, A.; Kronenberg, M.; Cheroutre, H.; Reinherz, E.L.; Wang, J.H.
Deposited on : 2002-12-12
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	274	63% (green), 32% (yellow), 5% (orange), 5% (red), 2% (grey)
2	B	99	51% (green), 41% (yellow), 8% (orange)
3	G	128	62% (green), 27% (yellow), 5% (orange), 5% (red), 2% (grey)
3	H	128	56% (green), 36% (yellow), 6% (orange)

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5194 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, TLA(C) alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	2237	1421	376	432	8	0	0	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	821	524	138	152	7	0	0	0

- Molecule 3 is a protein called T-cell surface glycoprotein CD8 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	122	962	616	156	183	7	0	0	0
3	H	120	954	612	155	180	7	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	123	SER	-	cloning artifact	UNP P01731
G	124	SER	-	cloning artifact	UNP P01731
G	125	ALA	-	cloning artifact	UNP P01731
G	126	LEU	-	cloning artifact	UNP P01731
G	127	VAL	-	cloning artifact	UNP P01731
G	128	PRO	-	cloning artifact	UNP P01731
H	123	SER	-	cloning artifact	UNP P01731
H	124	SER	-	cloning artifact	UNP P01731
H	125	ALA	-	cloning artifact	UNP P01731
H	126	LEU	-	cloning artifact	UNP P01731
H	127	VAL	-	cloning artifact	UNP P01731
H	128	PRO	-	cloning artifact	UNP P01731

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	H	1	Total	C	N	O	0	0
			14	8	1	5		
4	H	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	72	Total	O	0	0
			72	72		
5	B	15	Total	O	0	0
			15	15		
5	G	51	Total	O	0	0
			51	51		
5	H	26	Total	O	0	0
			26	26		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: H-2 class I histocompatibility antigen, TLA(C) alpha chain



- Molecule 2: Beta-2-microglobulin




- Molecule 3: T-cell surface glycoprotein CD8 alpha chain



- Molecule 3: T-cell surface glycoprotein CD8 alpha chain





4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	77.02Å 77.02Å 176.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-2.10)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.217 , 0.279	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5194	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/2304	0.74	4/3138 (0.1%)
2	B	0.51	0/847	0.74	0/1148
3	G	0.60	0/983	0.80	1/1328 (0.1%)
3	H	0.49	0/976	0.74	0/1319
All	All	0.54	0/5110	0.75	5/6933 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	272	LEU	CA-CB-CG	6.40	130.02	115.30
1	A	20	PRO	N-CA-C	-6.11	96.21	112.10
1	A	19	GLU	N-CA-C	6.07	127.39	111.00
3	G	54	VAL	CB-CA-C	-5.63	100.70	111.40
1	A	28	LEU	N-CA-C	-5.37	96.50	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2237	0	2090	99	0
2	B	821	0	798	58	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	962	0	957	45	0
3	H	954	0	952	45	0
4	G	28	0	26	2	0
4	H	28	0	26	2	0
5	A	72	0	0	3	0
5	B	15	0	0	1	0
5	G	51	0	0	1	0
5	H	26	0	0	0	0
All	All	5194	0	4849	237	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 237 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:ILE:HD11	3:G:105:ILE:HG21	1.24	1.07
1:A:14:ARG:HH22	1:A:20:PRO:HD2	1.27	0.99
1:A:58:GLU:HB3	1:A:62:ARG:HH21	1.28	0.97
2:B:54:MET:CE	2:B:64:ILE:HD11	1.97	0.94
1:A:238:ASP:OD2	1:A:240:THR:HG22	1.68	0.93

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:17:ASN:OD1	2:B:17:ASN:OD1[4_641]	1.41	0.79

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/274 (99%)	249 (92%)	16 (6%)	7 (3%)	4	1
2	B	97/99 (98%)	91 (94%)	4 (4%)	2 (2%)	5	2
3	G	120/128 (94%)	115 (96%)	4 (3%)	1 (1%)	16	13
3	H	118/128 (92%)	103 (87%)	14 (12%)	1 (1%)	16	13
All	All	607/629 (96%)	558 (92%)	38 (6%)	11 (2%)	7	3

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	18	SER
1	A	108	SER
2	B	48	LYS
3	H	59	SER
1	A	51	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/234 (100%)	224 (96%)	10 (4%)	25	25
2	B	94/94 (100%)	88 (94%)	6 (6%)	14	12
3	G	113/119 (95%)	104 (92%)	9 (8%)	10	7
3	H	112/119 (94%)	108 (96%)	4 (4%)	30	32
All	All	553/566 (98%)	524 (95%)	29 (5%)	19	18

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	77	THR
3	H	96	ASN
3	G	10	PHE
3	G	88	THR
3	G	7	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	96	ASN
3	H	3	GLN
3	H	120	GLN
3	H	90	ASN
3	H	96	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	H	2070	3	14,14,15	0.81	1 (7%)	17,19,21	0.66	0
4	NAG	G	1042	3	14,14,15	0.87	1 (7%)	17,19,21	0.71	0
4	NAG	H	2042	3	14,14,15	0.66	0	17,19,21	0.74	0
4	NAG	G	1070	3	14,14,15	0.82	0	17,19,21	0.79	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	H	2070	3	-	6/6/23/26	0/1/1/1
4	NAG	G	1042	3	-	4/6/23/26	0/1/1/1
4	NAG	H	2042	3	-	2/6/23/26	0/1/1/1
4	NAG	G	1070	3	-	4/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	2070	NAG	C1-C2	2.32	1.55	1.52
4	G	1042	NAG	C1-C2	2.20	1.55	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	1070	NAG	C8-C7-N2	2.17	119.72	116.12

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	1042	NAG	C1-C2-N2-C7
4	G	1042	NAG	C8-C7-N2-C2
4	G	1042	NAG	O7-C7-N2-C2
4	H	2042	NAG	C8-C7-N2-C2
4	H	2042	NAG	O7-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	2070	NAG	1	0
4	G	1042	NAG	2	0
4	H	2042	NAG	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.