



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 07:52 am GMT

PDB ID : 1NTB  
Title : 2.9 Å crystal structure of Streptomycin RNA-aptamer complex  
Authors : Tereshko, V.; Skripkin, E.; Patel, D.J.  
Deposited on : 2003-01-29  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1716 (2.90-2.90)
Clashscore	122126	1924 (2.90-2.90)
RNA backbone	2636	1059 (3.20-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	22	 64% 9% 27%
2	B	18	 44% 33% 11% 11%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(\*GP\*GP\*AP\*UP\*CP\*GP\*CP\*AP\*UP\*UP\*UP\*GP\*GP\*AP\*CP\*UP\*UP\*CP\*UP\*GP\*CP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	22	461	207	77	156	21	0	0	0


- Molecule 2 is a RNA chain called 5'-R(\*CP\*GP\*GP\*CP\*AP\*CP\*CP\*AP\*CP\*GP\*GP\*UP\*CP\*GP\*GP\*AP\*UP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	18	381	171	70	123	17	0	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		

- Molecule 4 is STREPTOMYCIN (three-letter code: SRY) (formula: C<sub>21</sub>H<sub>39</sub>N<sub>7</sub>O<sub>12</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	40	0
			40	21	7	12		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	O	0	0
			2	2		
6	B	4	Total	O	1	0
			4	4		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-R(\*GP\*GP\*AP\*UP\*CP\*GP\*CP\*AP\*UP\*UP\*UP\*GP\*GP\*AP\*CP\*UP\*UP\*CP\*UP\*GP\*CP\*C)-3'

Chain A: 



- Molecule 2: 5'-R(\*CP\*GP\*GP\*CP\*AP\*CP\*CP\*AP\*CP\*GP\*GP\*UP\*CP\*GP\*GP\*AP\*UP\*CP)-3'

Chain B: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.63Å 82.63Å 49.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.90 29.54 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-2.90) 99.3 (29.54-2.70)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 2.72Å)	Xtrriage
Refinement program	REFMAC 5.1.13	Depositor
R, $R_{free}$	0.204 , 0.255 0.304 , 0.335	Depositor DCC
$R_{free}$ test set	692 reflections (7.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.0	Xtrriage
Anisotropy	0.117	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 31.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.034 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	891	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, SRY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.99	3/513 (0.6%)	1.19	2/797 (0.3%)
2	B	1.54	4/425 (0.9%)	2.05	29/661 (4.4%)
All	All	1.27	7/938 (0.7%)	1.64	31/1458 (2.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	3	0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	112	U	P-OP2	12.75	1.70	1.49
2	B	117	U	P-OP2	12.66	1.70	1.49
2	B	106	C	O5'-C5'	11.53	1.62	1.44
1	A	11	U	C4-O4	9.33	1.31	1.23
1	A	11	U	C1'-N1	7.14	1.59	1.48
2	B	117	U	N1-C6	-5.53	1.32	1.38
1	A	19	U	C3'-O3'	-5.29	1.34	1.42

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	U	C5-C4-O4	-12.21	118.57	125.90
2	B	106	C	O5'-C5'-C4'	-11.56	89.73	111.70
2	B	112	U	N1-C1'-C2'	10.87	128.13	114.00
2	B	117	U	O4'-C1'-C2'	10.59	117.13	107.60

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	U	N1-C2-N3	10.50	121.20	114.90
2	B	117	U	C4'-C3'-C2'	10.21	112.81	102.60
2	B	112	U	C1'-O4'-C4'	-9.82	102.05	109.90
2	B	112	U	C2-N3-C4	-9.64	121.21	127.00
2	B	117	U	O4'-C1'-N1	9.25	115.60	108.20
2	B	112	U	O4'-C1'-C2'	9.06	115.76	107.60
2	B	117	U	N1-C2-N3	9.01	120.31	114.90
2	B	117	U	C2-N3-C4	-8.25	122.05	127.00
2	B	112	U	C2-N1-C1'	8.06	127.37	117.70
2	B	117	U	C2-N1-C1'	7.92	127.20	117.70
1	A	6	G	P-O3'-C3'	7.88	129.16	119.70
2	B	112	U	O4'-C1'-N1	7.66	114.33	108.20
2	B	117	U	C6-N1-C1'	-7.10	111.27	121.20
2	B	117	U	C5-C4-O4	-7.02	121.69	125.90
2	B	106	C	P-O5'-C5'	-6.82	109.98	120.90
2	B	112	U	C6-N1-C1'	-6.43	112.19	121.20
2	B	107	C	P-O3'-C3'	6.13	127.06	119.70
2	B	112	U	N3-C4-C5	6.03	118.22	114.60
1	A	8	A	O4'-C1'-N9	5.75	112.80	108.20
2	B	117	U	C2'-C3'-O3'	5.74	122.88	113.70
2	B	112	U	C3'-C2'-C1'	-5.65	96.98	101.50
2	B	117	U	C4'-C3'-O3'	5.62	124.23	113.00
2	B	112	U	N3-C2-O2	-5.50	118.35	122.20
2	B	112	U	N3-C4-O4	5.44	123.21	119.40
2	B	117	U	C3'-C2'-C1'	-5.41	97.18	101.50
2	B	117	U	N1-C1'-C2'	5.40	121.02	114.00
2	B	107	C	O3'-P-O5'	-5.17	94.17	104.00

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	112	U	C3',C1'
2	B	117	U	C3'

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	461	0	236	2	0
2	B	381	0	198	2	0
3	A	2	0	0	0	0
4	A	40	0	37	0	0
5	B	1	0	0	0	0
6	A	2	0	0	0	0
6	B	4	0	0	0	0
All	All	891	0	471	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:U:C6	1:A:9:U:H5''	2.45	0.51
1:A:9:U:H3'	1:A:10:U:C5'	2.46	0.46
2:B:103:G:C6	2:B:104:C:C4	3.09	0.40
2:B:107:C:H4'	2:B:107:C:OP1	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	21/22 (95%)	7 (33%)	3 (14%)
2	B	17/18 (94%)	8 (47%)	4 (23%)
All	All	38/40 (95%)	15 (39%)	7 (18%)

All (15) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	7	C
1	A	8	A
1	A	9	U
1	A	10	U
1	A	11	U
1	A	13	G
1	A	19	U
2	B	106	C
2	B	107	C
2	B	108	A
2	B	109	C
2	B	112	U
2	B	113	C
2	B	117	U
2	B	118	C

All (7) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	6	G
1	A	8	A
1	A	10	U
2	B	107	C
2	B	108	A
2	B	112	U
2	B	117	U

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SRY	A	25	-	39,42,42	1.76	6 (15%)	47,63,63	1.41	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SRY	A	25	-	-	0/20/87/87	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	25	SRY	CD1-N31	-4.52	1.25	1.33
4	A	25	SRY	O13-C22	-3.20	1.35	1.44
4	A	25	SRY	C23-N23	-2.96	1.42	1.47
4	A	25	SRY	O21-C21	2.79	1.49	1.43
4	A	25	SRY	C11-N11	3.66	1.51	1.45
4	A	25	SRY	OG2-CG2	6.45	1.42	1.19

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	25	SRY	C12-O41-C41	-4.75	106.11	117.97
4	A	25	SRY	C13-O13-C22	-2.72	111.52	116.28
4	A	25	SRY	O21-C21-C11	-2.41	104.77	109.62
4	A	25	SRY	C43-C33-C23	-2.29	106.94	110.33
4	A	25	SRY	N31-CD1-NF1	-2.17	116.66	120.50
4	A	25	SRY	C21-C31-N31	2.44	115.26	110.58
4	A	25	SRY	CI3-N23-C23	3.90	120.06	114.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.