



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 11:36 AM EDT

PDB ID : 4NU8
Title : Crystal structure of O-acetylserine sulfhydrylase from Haemophilus influenzae in complex with high affinity inhibitory peptide from serine acetyl transferase of Salmonella typhimurium at 2.0 Å
Authors : Ekka, M.K.; Kaushik, A.; Singh, A.K.; Kumaran, S.
Deposited on : 2013-12-03
Resolution : 2.07 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	322	<p>10% 86% 10% ..</p>
2	A	8	<p>50% 88% 12%</p>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	X	311	2274	1429	394	442	1	8	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	-5	HIS	-	expression tag	UNP P45040
X	-4	HIS	-	expression tag	UNP P45040
X	-3	HIS	-	expression tag	UNP P45040
X	-2	HIS	-	expression tag	UNP P45040
X	-1	HIS	-	expression tag	UNP P45040
X	0	HIS	-	expression tag	UNP P45040

- Molecule 2 is a protein called Peptide from Serine acetyltransferase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	A	8	64	41	8	15	0	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	X	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	X	79	Total O 79 79	0	0
4	A	2	Total O 2 2	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, α , β , γ	112.32Å 112.32Å 45.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.52 – 2.07 35.52 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.8 (35.52-2.07) 99.8 (35.52-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.83 (at 2.06Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.187 , 0.235 0.180 , 0.225	Depositor DCC
R_{free} test set	892 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtrriage
Anisotropy	0.534	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2425	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	X	0.43	0/2282	0.52	0/3102
2	A	0.23	0/65	0.36	0/85
All	All	0.42	0/2347	0.52	0/3187

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2274	0	2277	22	0
2	A	64	0	51	1	0
3	X	6	0	8	1	0
4	A	2	0	0	0	0
4	X	79	0	0	0	0
All	All	2425	0	2336	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:255:LEU:HD21	1:X:310:LEU:HA	1.57	0.87
1:X:227:GLN:HE21	1:X:305:TYR:HE1	1.38	0.70
1:X:97:SER:O	1:X:99:GLU:N	2.27	0.67
1:X:59:LEU:HD13	1:X:139:VAL:HG11	1.78	0.65
1:X:38:SER:HB2	1:X:75:ILE:HG22	1.82	0.59
1:X:292:LYS:HD2	1:X:292:LYS:N	2.18	0.57
1:X:269:ALA:O	1:X:302:SER:HB3	2.05	0.57
1:X:269:ALA:HB1	1:X:273:SER:CB	2.39	0.53
1:X:231:ALA:O	2:A:269:TYR:HA	2.11	0.51
1:X:253:THR:HG21	3:X:401:GOL:H31	1.93	0.49
1:X:149:ASN:HB3	1:X:150:PRO:CD	2.42	0.49
1:X:291:ASP:C	1:X:292:LYS:HD2	2.34	0.48
1:X:227:GLN:NE2	1:X:305:TYR:CE1	2.82	0.46
1:X:42:LLP:H2'1	1:X:72:ASN:ND2	2.31	0.45
1:X:259:ARG:NH2	1:X:309:ALA:O	2.49	0.45
1:X:96:MET:HE1	1:X:100:ARG:HD3	1.99	0.44
1:X:29:VAL:CG2	1:X:295:VAL:HG22	2.47	0.43
1:X:149:ASN:OD1	1:X:180:GLY:HA3	2.19	0.42
1:X:242:ILE:C	1:X:242:ILE:HD12	2.39	0.42
1:X:100:ARG:NH1	1:X:227:GLN:OE1	2.48	0.41
1:X:149:ASN:O	1:X:152:ILE:HG22	2.19	0.41
1:X:269:ALA:HB1	1:X:273:SER:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	309/322 (96%)	295 (96%)	13 (4%)	1 (0%)	41	32
2	A	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
All	All	315/330 (96%)	300 (95%)	14 (4%)	1 (0%)	41	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	98	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	231/253 (91%)	226 (98%)	5 (2%)	52	46
2	A	6/6 (100%)	6 (100%)	0	100	100
All	All	237/259 (92%)	232 (98%)	5 (2%)	53	48

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	69	THR
1	X	79	TYR
1	X	120	MET
1	X	255	LEU
1	X	303	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	X	42	1	23,24,25	1.63	4 (17%)	25,32,34	1.71	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	X	42	1	-	1/16/17/19	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	42	LLP	O3-C3	-5.20	1.25	1.36
1	X	42	LLP	C2-N1	3.13	1.39	1.33
1	X	42	LLP	C4-C4'	2.22	1.51	1.46
1	X	42	LLP	CE-NZ	2.01	1.51	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	42	LLP	OP4-C5'-C5	5.65	119.95	109.36
1	X	42	LLP	C5-C4-C4'	-2.58	117.48	121.47
1	X	42	LLP	CE-NZ-C4'	-2.58	110.45	118.72
1	X	42	LLP	C4-C3-C2	-2.06	118.99	120.14
1	X	42	LLP	C4-C4'-NZ	-2.01	114.75	124.04

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	X	42	LLP	CE-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	X	42	LLP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	X	401	-	5,5,5	0.45	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	X	401	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	X	401	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	310/322 (96%)	0.39	33 (10%) 6 6	24, 32, 70, 83	0
2	A	8/8 (100%)	2.27	4 (50%) 0 0	35, 39, 41, 42	0
All	All	318/330 (96%)	0.44	37 (11%) 4 4	24, 33, 70, 83	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	307	SER	7.6
1	X	98	LEU	7.0
2	A	269	TYR	7.0
1	X	132	ALA	6.9
1	X	130	ILE	5.9
1	X	133	SER	5.7
1	X	122	GLY	5.5
1	X	308	THR	4.9
1	X	94	GLU	4.7
1	X	309	ALA	4.2
1	X	96	MET	3.8
1	X	123	ALA	3.4
1	X	134	ASP	3.4
1	X	62	GLY	3.4
1	X	129	GLU	3.2
1	X	112	VAL	3.1
1	X	117	ALA	3.1
1	X	115	GLU	3.0
2	A	265	THR	2.9
1	X	306	LEU	2.8
2	A	271	ASP	2.8
1	X	1	MET	2.8
1	X	119	GLY	2.7
1	X	138	TYR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	128	GLU	2.5
1	X	102	ARG	2.5
1	X	126	LYS	2.5
2	A	268	GLU	2.5
1	X	131	VAL	2.5
1	X	93	PRO	2.4
1	X	95	THR	2.4
1	X	114	THR	2.4
1	X	113	LEU	2.4
1	X	125	ALA	2.3
1	X	135	PRO	2.2
1	X	35[A]	ARG	2.2
1	X	118	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	X	42	24/25	0.98	0.14	23,27,29,31	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	X	401	6/6	0.88	0.14	32,40,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.