



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 12:19 am GMT

PDB ID : 7NUC
Title : Crystal Structure of Neisseria gonorrhoeae LeuRS L550G mutant in Complex with ATP and L-leucinol
Authors : Pang, L.; Strelkov, S.V.; Weeks, S.D.
Deposited on : 2021-03-11
Resolution : 2.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	877	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leucine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	862	6583	4199	1119	1228	37	0	2	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP B4RNT1
A	454	ASN	ASP	conflict	UNP B4RNT1
A	508	ILE	MET	conflict	UNP B4RNT1
A	550	GLY	LEU	engineered mutation	UNP B4RNT1

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

Continued on next page...



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	8	6	1	1	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	148	148	148	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leucine-tRNA ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.91Å 81.59Å 216.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.37 – 2.77 108.49 – 2.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (76.37-2.77) 100.0 (108.49-2.75)	Depositor EDS
R_{merge}	0.37	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.77Å)	Xtrriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.185 , 0.264 0.187 , 0.265	Depositor DCC
R_{free} test set	1153 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtrriage
Anisotropy	0.240	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6781	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, EDO, ATP, DCL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/6759	0.48	0/9203

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6583	0	6163	80	0
2	A	8	0	12	0	0
3	A	31	0	12	0	0
4	A	2	0	0	0	0
5	A	8	0	12	1	0
6	A	1	0	0	0	0
7	A	148	0	0	0	0
All	All	6781	0	6199	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:587:THR:HG23	1:A:630:SER:HB2	1.70	0.73
1:A:82:PHE:CE1	1:A:128:CYS:HA	2.24	0.73
1:A:312:VAL:HG12	1:A:331:ILE:HB	1.72	0.70
1:A:43:TYR:O	1:A:637:LYS:NZ	2.26	0.69
1:A:87:GLU:OE2	1:A:438:ARG:NH1	2.25	0.68
1:A:52:HIS:CE1	1:A:637:LYS:HD3	2.31	0.66
1:A:52:HIS:HE1	1:A:637:LYS:HD3	1.62	0.64
1:A:83:GLY:HA3	1:A:506:THR:HB	1.82	0.61
1:A:798:LYS:HB2	1:A:801:GLU:HG2	1.82	0.61
1:A:134:ARG:HB3	1:A:463:PRO:HB3	1.85	0.57
1:A:195:ILE:HG22	1:A:431:ASP:HA	1.86	0.57
1:A:241:LYS:HA	1:A:244:LEU:HD23	1.86	0.57
1:A:225:GLY:HA3	1:A:428:ARG:HB2	1.88	0.55
1:A:666:PRO:HB3	1:A:739:GLN:OE1	2.07	0.55
1:A:32:LYS:HB3	1:A:72:ASN:HB2	1.90	0.53
1:A:589:TYR:CZ	1:A:597:LYS:HE3	2.44	0.52
1:A:807:VAL:HG12	1:A:809:GLU:HG3	1.91	0.52
1:A:358:ALA:HB1	1:A:363:LEU:HB2	1.90	0.52
1:A:348:PRO:HB2	1:A:365:LYS:HG3	1.90	0.52
1:A:368:VAL:O	1:A:394:VAL:N	2.33	0.52
1:A:268:VAL:HG23	1:A:329:VAL:HA	1.92	0.52
1:A:263:MET:HG3	1:A:396:SER:HB3	1.92	0.51
1:A:866:VAL:HG13	1:A:872:VAL:HG22	1.90	0.51
1:A:261:THR:HB	1:A:347:VAL:HG11	1.92	0.51
1:A:82:PHE:CZ	1:A:128:CYS:HA	2.46	0.50
1:A:480:ALA:HA	1:A:501:THR:HG21	1.94	0.49
1:A:41:PHE:CD2	1:A:513:TYR:CE2	3.01	0.49
1:A:834:ALA:HB3	1:A:837:ALA:HB2	1.95	0.48
1:A:41:PHE:HD2	1:A:41:PHE:H	1.61	0.48
1:A:177:TRP:CE2	1:A:178:ARG:HG2	2.47	0.48
1:A:258:ARG:NH1	1:A:260:ASP:OD1	2.45	0.48
1:A:40:MET:HG2	5:A:1004:DCL:HD12	1.95	0.48
1:A:255:TYR:CZ	1:A:426:GLN:HG3	2.48	0.48
1:A:744:ILE:HD11	1:A:782:ILE:HD13	1.95	0.47
1:A:82:PHE:CE1	1:A:128:CYS:CA	2.95	0.47
1:A:635:MET:HG2	1:A:642:GLY:HA2	1.95	0.47
1:A:247:ASP:HA	1:A:250:LYS:HD2	1.96	0.47
1:A:267:TYR:OH	1:A:358:ALA:HB2	2.14	0.47
1:A:270:VAL:HG22	1:A:274:HIS:HB3	1.97	0.47
1:A:350:HIS:CE1	1:A:369:ILE:HB	2.50	0.47
1:A:64:ARG:NH2	1:A:116:GLY:O	2.47	0.46
1:A:439:TYR:CZ	1:A:470:VAL:HB	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:543:GLY:O	1:A:579:THR:HA	2.16	0.46
1:A:839:LYS:HA	1:A:842:LEU:HB2	1.98	0.46
1:A:660:PHE:CE1	1:A:677:GLY:HA3	2.51	0.46
1:A:244:LEU:HD11	1:A:276:LEU:HA	1.97	0.46
1:A:146:LYS:NZ	1:A:565:GLU:OE2	2.36	0.45
1:A:360:LYS:HD2	1:A:361:TYR:CZ	2.50	0.45
1:A:587:THR:HG22	1:A:631:GLY:O	2.16	0.45
1:A:589:TYR:HB3	1:A:599:TRP:CE3	2.52	0.45
1:A:35:TYR:CE2	1:A:37:CYS:HB2	2.52	0.45
1:A:479:LEU:HD22	1:A:485:PHE:CD2	2.52	0.44
1:A:361:TYR:HB2	1:A:363:LEU:CD2	2.48	0.44
1:A:206:LEU:HD23	1:A:209:LEU:HD12	2.00	0.44
1:A:41:PHE:CE2	1:A:513:TYR:CZ	3.06	0.43
1:A:82:PHE:CE1	1:A:128:CYS:CB	3.01	0.43
1:A:123:ARG:HA	1:A:520:PRO:HG3	1.99	0.43
1:A:439:TYR:O	1:A:441:GLY:N	2.52	0.43
1:A:798:LYS:HD2	1:A:800:TRP:CZ2	2.54	0.43
1:A:134:ARG:CB	1:A:463:PRO:HB3	2.48	0.42
1:A:150:TYR:OH	1:A:431:ASP:OD2	2.23	0.42
1:A:541:TYR:C	1:A:542:ILE:HG13	2.39	0.42
1:A:142:LYS:HA	1:A:142:LYS:HD3	1.91	0.42
1:A:41:PHE:CD2	1:A:513:TYR:CZ	3.08	0.42
1:A:63:SER:HB3	1:A:73:VAL:HG11	2.00	0.42
1:A:374:ASN:HB2	1:A:384:TRP:CE2	2.55	0.42
1:A:185:LYS:HD3	1:A:185:LYS:HA	1.92	0.42
1:A:348:PRO:HB2	1:A:365:LYS:CG	2.49	0.42
1:A:351:ASP:HB3	1:A:354:ASP:HB2	2.02	0.41
1:A:763:GLN:O	1:A:767:VAL:HG23	2.20	0.41
1:A:47:LYS:O	1:A:637:LYS:HG2	2.19	0.41
1:A:267:TYR:HA	1:A:327:LEU:HB3	2.02	0.41
1:A:448:HIS:O	1:A:497:ALA:HB1	2.21	0.41
1:A:54:ARG:NH1	1:A:581:GLY:O	2.54	0.41
1:A:486:TYR:HA	1:A:499:ARG:HB2	2.02	0.41
1:A:664:ALA:HB3	1:A:671:LEU:HD11	2.03	0.41
1:A:183:ILE:HD13	1:A:183:ILE:HA	1.87	0.40
1:A:240:SER:HB3	1:A:316:ARG:HB3	2.02	0.40
1:A:87:GLU:OE2	1:A:438:ARG:HD2	2.21	0.40
1:A:856:MET:CE	1:A:874:ILE:HG21	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	858/877 (98%)	817 (95%)	38 (4%)	3 (0%)	41 70

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	372	GLY
1	A	337	TRP
1	A	850	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	639/716 (89%)	615 (96%)	24 (4%)	33 64

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	SER
1	A	34	LYS
1	A	36	TYR
1	A	41	PHE
1	A	55	ASN
1	A	82	PHE
1	A	94	ASN
1	A	104	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	186	ARG
1	A	214	GLU
1	A	258	ARG
1	A	290	PHE
1	A	339	TYR
1	A	365	LYS
1	A	422	GLU
1	A	489	SER
1	A	541	TYR
1	A	558	PHE
1	A	610	ASP
1	A	636	SER
1	A	670	SER
1	A	820	MET
1	A	828	ARG
1	A	836	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	454	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ATP	A	1002	4	26,33,33	0.63	0	31,52,52	0.70	1 (3%)
2	EDO	A	1001	-	3,3,3	0.53	0	2,2,2	0.22	0
2	EDO	A	1005	-	3,3,3	0.46	0	2,2,2	0.26	0
5	DCL	A	1004	-	7,7,7	0.10	0	6,8,8	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	1002	4	-	5/18/38/38	0/3/3/3
2	EDO	A	1001	-	-	0/1/1/1	-
2	EDO	A	1005	-	-	0/1/1/1	-
5	DCL	A	1004	-	-	1/6/6/6	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1002	ATP	C5-C6-N6	2.35	123.92	120.35

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1002	ATP	C5'-O5'-PA-O1A
3	A	1002	ATP	C5'-O5'-PA-O2A
3	A	1002	ATP	PB-O3A-PA-O1A
3	A	1002	ATP	C5'-O5'-PA-O3A
3	A	1002	ATP	PG-O3B-PB-O2B
5	A	1004	DCL	O-C-CA-N


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1004	DCL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	862/877 (98%)	0.15	50 (5%) 23 18	20, 37, 118, 210	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	873	ASN	13.9
1	A	868	PRO	7.6
1	A	852	ALA	6.6
1	A	875	VAL	6.3
1	A	865	ILE	6.1
1	A	872	VAL	5.0
1	A	864	ILE	4.8
1	A	361	TYR	4.2
1	A	866	VAL	4.0
1	A	827	LEU	3.9
1	A	874	ILE	3.8
1	A	867	VAL	3.8
1	A	331	ILE	3.7
1	A	243	GLY	3.6
1	A	273	GLU	3.5
1	A	360	LYS	3.4
1	A	869	GLY	3.4
1	A	835	ALA	3.2
1	A	839	LYS	3.2
1	A	870	ARG	3.1
1	A	312	VAL	3.0
1	A	840	ALA	3.0
1	A	838	SER	3.0
1	A	871	LEU	3.0
1	A	311	GLY	3.0
1	A	846	ALA	2.8
1	A	832	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	823	VAL	2.7
1	A	279	ALA	2.7
1	A	847	LEU	2.7
1	A	828	ARG	2.7
1	A	856	MET	2.6
1	A	332	ALA	2.6
1	A	836	ASP	2.5
1	A	282	ALA	2.5
1	A	334	TYR	2.5
1	A	278	THR	2.4
1	A	358	ALA	2.4
1	A	357	PHE	2.4
1	A	419	GLY	2.4
1	A	819	VAL	2.4
1	A	876	VAL	2.4
1	A	363	LEU	2.4
1	A	853	VAL	2.4
1	A	826	LYS	2.3
1	A	851	GLY	2.3
1	A	314	THR	2.3
1	A	376	PHE	2.2
1	A	855	PHE	2.1
1	A	291	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	A	1007	1/1	0.89	0.12	64,64,64,64	0
2	EDO	A	1001	4/4	0.93	0.14	36,40,40,43	0
4	MG	A	1003	1/1	0.96	0.05	12,12,12,12	0
2	EDO	A	1005	4/4	0.97	0.23	27,33,33,35	0
5	DCL	A	1004	8/8	0.97	0.18	11,20,24,26	0
3	ATP	A	1002	31/31	0.98	0.14	15,25,34,39	0
6	ZN	A	1006	1/1	0.98	0.17	67,67,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.