

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 05:44 PM EDT

PDB ID	:	6NX3
Title	:	Structures of the transcriptional regulator BgaR, a lactose sensor.
Authors	:	Peat, T.S.; Newman, J.
Deposited on	:	2019-02-07
Resolution	:	1.87 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 1.87 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	158	Total	С	Ν	0	\mathbf{S}	0	4	0
	A	150	1314	851	205	251	7	0	4	0
1	р	159	Total	С	Ν	0	S	0	4	0
	D	150	1308	846	203	252	7	0	4	0
1	C	158	Total	С	Ν	0	S	0	6	0
			1324	857	207	253	7		0	0
1	П	159	Total	С	Ν	0	S	0	5	0
	D		1325	860	208	250	7	0		
1	F	150	Total	С	Ν	0	S	0	1	0
		109	1289	838	202	242	7	0		0
1	Б	150	Total	С	Ν	0	\mathbf{S}	0	1	0
	Г	159	1317	854	207	249	$\overline{7}$		4	U

• Molecule 1 is a protein called Transcriptional regulator BgaR.

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	171	LEU	-	expression tag	UNP Q8XMB9
А	172	VAL	-	expression tag	UNP Q8XMB9
А	173	PRO	-	expression tag	UNP Q8XMB9
А	174	ARG	-	expression tag	UNP Q8XMB9
А	175	GLY	-	expression tag	UNP Q8XMB9
А	176	SER	-	expression tag	UNP Q8XMB9
А	177	HIS	-	expression tag	UNP Q8XMB9
А	178	HIS	-	expression tag	UNP Q8XMB9
А	179	HIS	-	expression tag	UNP Q8XMB9
А	180	HIS	-	expression tag	UNP Q8XMB9
А	181	HIS	-	expression tag	UNP Q8XMB9
А	182	HIS	-	expression tag	UNP Q8XMB9
В	171	LEU	-	expression tag	UNP Q8XMB9
В	172	VAL	-	expression tag	UNP Q8XMB9
В	173	PRO	-	expression tag	UNP Q8XMB9
В	174	ARG	-	expression tag	UNP Q8XMB9
В	175	GLY	-	expression tag	UNP Q8XMB9

Continued on next page...



Continued from previous page						
Chain	Residue	Modelled	Actual	Comment	Reference	
В	176	SER	-	expression tag	UNP Q8XMB9	
В	177	HIS	-	expression tag	UNP Q8XMB9	
В	178	HIS	-	expression tag	UNP Q8XMB9	
В	179	HIS	-	expression tag	UNP Q8XMB9	
В	180	HIS	-	expression tag	UNP Q8XMB9	
В	181	HIS	-	expression tag	UNP Q8XMB9	
В	182	HIS	-	expression tag	UNP Q8XMB9	
С	171	LEU	-	expression tag	UNP Q8XMB9	
С	172	VAL	-	expression tag	UNP Q8XMB9	
С	173	PRO	-	expression tag	UNP Q8XMB9	
С	174	ARG	-	expression tag	UNP Q8XMB9	
С	175	GLY	-	expression tag	UNP Q8XMB9	
С	176	SER	-	expression tag	UNP Q8XMB9	
С	177	HIS	-	expression tag	UNP Q8XMB9	
С	178	HIS	-	expression tag	UNP Q8XMB9	
С	179	HIS	-	expression tag	UNP Q8XMB9	
С	180	HIS	-	expression tag	UNP Q8XMB9	
С	181	HIS	-	expression tag	UNP Q8XMB9	
С	182	HIS	-	expression tag	UNP Q8XMB9	
D	171	LEU	-	expression tag	UNP Q8XMB9	
D	172	VAL	-	expression tag	UNP Q8XMB9	
D	173	PRO	-	expression tag	UNP Q8XMB9	
D	174	ARG	-	expression tag	UNP Q8XMB9	
D	175	GLY	-	expression tag	UNP Q8XMB9	
D	176	SER	-	expression tag	UNP Q8XMB9	
D	177	HIS	-	expression tag	UNP Q8XMB9	
D	178	HIS	-	expression tag	UNP Q8XMB9	
D	179	HIS	-	expression tag	UNP Q8XMB9	
D	180	HIS	-	expression tag	UNP Q8XMB9	
D	181	HIS	-	expression tag	UNP Q8XMB9	
D	182	HIS	-	expression tag	UNP Q8XMB9	
Е	171	LEU	-	expression tag	UNP Q8XMB9	
Е	172	VAL	-	expression tag	UNP Q8XMB9	
Е	173	PRO	-	expression tag	UNP Q8XMB9	
Е	174	ARG	-	expression tag	UNP Q8XMB9	
Е	175	GLY	-	expression tag	UNP Q8XMB9	
Е	176	SER	-	expression tag	UNP Q8XMB9	
Е	177	HIS	-	expression tag	UNP Q8XMB9	
Е	178	HIS	-	expression tag	UNP Q8XMB9	
Е	179	HIS	-	expression tag	UNP Q8XMB9	
Е	180	HIS	-	expression tag	UNP Q8XMB9	
Е	181	HIS	-	expression tag	UNP Q8XMB9	

6NX3

Continued on next page...



6NX3

Chain	Residue	Modelled	Actual	Comment	Reference
Е	182	HIS	-	expression tag	UNP Q8XMB9
F	171	LEU	-	expression tag	UNP Q8XMB9
F	172	VAL	-	expression tag	UNP Q8XMB9
F	173	PRO	-	expression tag	UNP Q8XMB9
F	174	ARG	-	expression tag	UNP Q8XMB9
F	175	GLY	-	expression tag	UNP Q8XMB9
F	176	SER	-	expression tag	UNP Q8XMB9
F	177	HIS	-	expression tag	UNP Q8XMB9
F	178	HIS	-	expression tag	UNP Q8XMB9
F	179	HIS	-	expression tag	UNP Q8XMB9
F	180	HIS	-	expression tag	UNP Q8XMB9
F	181	HIS	-	expression tag	UNP Q8XMB9
F	182	HIS	-	expression tag	UNP Q8XMB9

Continued from previous page...

• Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace		
9	G	2	Total C O	0	0	0		
2	G	2	23 12 11	0	0	0		
2	н	2	Total C O	0	0	0		
2	11	2	23 12 11	0	0	0		
9	Ι	т	т	0	Total C O	0	0	0
		Δ	23 12 11	0	0	0		
0	т	2	Total C O	0	0	0		
	J		23 12 11	0	0			
0	K	Z 2	Total C O	0	0	0		
	Γ		23 12 11	0	0			
9	т	2	Total C O	0	0	0		
2	L		23 12 11		0	U		

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	69	Total O 69 69	0	0
3	В	85	Total O 85 85	0	0
3	С	79	Total O 79 79	0	0
3	D	94	Total O 94 94	0	0
3	Е	86	Total O 86 86	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	F	86	Total O 86 86	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	99.15Å 46.32Å 118.90Å	Depositor
a, b, c, α , β , γ	90.00° 100.65° 90.00°	Depositor
Resolution (Å)	46.86 - 1.87	Depositor
% Data completeness	99 4 (46 86-1 87)	Depositor
(in resolution range)	33.4 (40.00-1.01)	
R _{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.00 (at 1.87 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.224 , 0.254	Depositor
Wilson B-factor $(Å^2)$	25.1	Xtriage
Anisotropy	0.190	Xtriage
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8514	wwPDB-VP
Average B, all atoms $(Å^2)$	30.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 93.39 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.5046e-09. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for a centric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Iol Turno Chain De		Dog	Tink	Bo	ond leng	$_{\rm ths}$	Bond angles		
10101	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	G	1	2	12,12,12	0.46	0	17,17,17	0.87	0
2	GAL	G	2	2	11,11,12	0.54	0	$15,\!15,\!17$	0.98	0
2	BGC	Н	1	2	12,12,12	0.45	0	17,17,17	1.10	2 (11%)
2	GAL	Н	2	2	11,11,12	0.62	0	15,15,17	1.14	0
2	BGC	Ι	1	2	12,12,12	0.39	0	17,17,17	0.95	0
2	GAL	Ι	2	2	11,11,12	0.59	0	15,15,17	1.14	2 (13%)
2	BGC	J	1	2	12,12,12	0.49	0	17,17,17	1.29	2 (11%)
2	GAL	J	2	2	11,11,12	0.67	0	15,15,17	1.05	1 (6%)
2	BGC	K	1	2	12,12,12	0.72	0	17,17,17	1.08	1 (5%)
2	GAL	K	2	2	11,11,12	0.59	0	15,15,17	0.85	0
2	BGC	L	1	2	12,12,12	0.67	0	17,17,17	1.07	1 (5%)
2	GAL	L	2	2	11,11,12	0.78	0	15,15,17	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
2	BGC	G	1	2	-	0/2/22/22	0/1/1/1
2	GAL	G	2	2	-	0/2/19/22	0/1/1/1
2	BGC	Н	1	2	-	1/2/22/22	0/1/1/1
2	GAL	Н	2	2	-	0/2/19/22	0/1/1/1
2	BGC	Ι	1	2	-	0/2/22/22	0/1/1/1
2	GAL	Ι	2	2	-	0/2/19/22	0/1/1/1
2	BGC	J	1	2	-	1/2/22/22	0/1/1/1
2	GAL	J	2	2	-	0/2/19/22	0/1/1/1
2	BGC	Κ	1	2	-	0/2/22/22	0/1/1/1
2	GAL	Κ	2	2	-	0/2/19/22	0/1/1/1
2	BGC	L	1	2	-	1/2/22/22	0/1/1/1
2	GAL	L	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
2	L	1	BGC	O6-C6-C5	-2.73	101.93	111.29
2	Κ	1	BGC	O6-C6-C5	-2.67	102.13	111.29
2	J	1	BGC	O5-C5-C4	2.61	114.43	109.69
2	J	1	BGC	O6-C6-C5	-2.56	102.51	111.29
2	Н	1	BGC	O3-C3-C2	-2.50	104.57	110.35



There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	1	BGC	C4-C5-C6-O6
2	L	1	BGC	C4-C5-C6-O6
2	Н	1	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











6NX3















4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

